

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 September 2003 (25.09.2003)

PCT

(10) International Publication Number  
**WO 03/077914 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/4184**,  
C07D 235/04, 235/06

(74) Agent: NOE, Jeremy, E.; McDonnell Boehnen Hubert &  
Berghoff, 300 South Wacker Drive, Suite 3200, Chicago,  
IL 60606 (US).

(21) International Application Number: PCT/US03/07864

(22) International Filing Date: 13 March 2003 (13.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/364,007 13 March 2002 (13.03.2002) US

(71) Applicant (for all designated States except US): **ARRAY  
BIOPHARMA, INC** [US/US]; 3200 Walnut Street, Boul-  
der, CO 80301 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WALLACE, Eli**,  
**M.** [US/US]; P.O. Box 1727, Lyons, CO 80540 (US).  
**LYSSIKATOS, Joseph, P.** [US/US]; 1720 Eldorado  
Circle, Superior, CO 80027 (US). **HURLEY, Brian, T.**  
[US/US]; 3725 Birchwood Drive, #22, Boulder, CO 80304  
(US). **MARLOW, Allison, L.** [US/US]; 2722 Juniper  
Avenue, #57, Boulder, CO 80304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,  
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: N3 ALKYLATED BENZIMIDAZOLE DERIVATIVES AS MEK INHIBITORS



(I)

(57) Abstract: Disclosed are compounds of the formula I, and pharmaceutically acceptable salts and prodrugs thereof, wherein W, R<sup>1</sup>, R<sup>2</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup> and R<sup>10</sup> are as defined in the specification. Such compounds are MEK inhibitors and useful in the treatment of hyperproliferative diseases, such as cancer and inflammation, in mammals. Also disclosed is a method of using such compounds in the treatment of hyperproliferative diseases in mammals, and pharmaceutical compositions containing such compounds.

**N3 Alkylated Benzimidazole Derivatives as MEK Inhibitors****BACKGROUND OF THE INVENTION****Field of the Invention**

This invention relates to a series of alkylated (1H-Benzoimidazol-5-yl)-(4-substituted-phenyl)-amine derivatives that are useful in the treatment of hyperproliferative diseases, such as cancer and inflammation, in mammals. This invention also relates to a method of using such compounds in the treatment of hyperproliferative diseases in mammals, especially humans, and to pharmaceutical compositions containing such compounds.

**Summary of the Related Art**

Cell signaling through growth factor receptors and protein kinases is an important regulator of cell growth, proliferation and differentiation. In normal cell growth, growth factors, through receptor activation (i.e. PDGF or EGF and others), activate MAP kinase pathways. One of the most important and most well understood MAP kinase pathways involved in normal and uncontrolled cell growth is the Ras/ Raf kinase pathway. Active GTP-bound Ras results in the activation and indirect phosphorylation of Raf kinase. Raf then phosphorylates MEK1 and 2 on two serine residues (S218 and S222 for MEK1 and S222 and S226 for MEK2) (Ahn *et al.*, *Methods in Enzymology* 2001, 332, 417-431). Activated MEK then phosphorylates its only known substrates, the MAP kinases, ERK1 and 2. ERK phosphorylation by MEK occurs on Y204 and T202 for ERK1 and Y185 and T183 for ERK2 (Ahn *et al.*, *Methods in Enzymology* 2001, 332, 417-431). Phosphorylated ERK dimerizes and then translocates to the nucleus where it accumulates (Khokhlatchev *et al.*, *Cell* 1998, 93, 605-615). In the nucleus, ERK is involved in several important cellular functions, including but not limited to nuclear transport, signal transduction, DNA repair, nucleosome assembly and translocation, and mRNA processing and

translation (Ahn *et al.*, *Molecular Cell* 2000, 6, 1343-1354). Overall, treatment of cells with growth factors leads to the activation of ERK1 and 2 which results in proliferation and, in some cases, differentiation (Lewis *et al.*, *Adv. Cancer Res.* 1998, 74, 49-139).

In proliferative diseases, genetic mutations and/or overexpression of the growth factor receptors, downstream signaling proteins, or protein kinases involved in the ERK kinase pathway lead to uncontrolled cell proliferation and, eventually, tumor formation. For example, some cancers contain mutations which result in the continuous activation of this pathway due to continuous production of growth factors. Other mutations can lead to defects in the deactivation of the activated GTP-bound Ras complex, again resulting in activation of the MAP kinase pathway. Mutated, oncogenic forms of Ras are found in 50% of colon and >90% pancreatic cancers as well as many others types of cancers (Kohl *et al.*, *Science* 1993, 260, 1834-1837). Recently, bRaf mutations have been identified in more than 60% of malignant melanoma (Davies, H. *et al.*, *Nature* 2002, 417, 949-954). These mutations in bRaf result in a constitutively active MAP kinase cascade. Studies of primary tumor samples and cell lines have also shown constitutive or overactivation of the MAP kinase pathway in cancers of pancreas, colon, lung, ovary and kidney (Hoshino, R. *et al.*, *Oncogene* 1999, 18, 813-822). Hence, there is a strong correlation between cancers and an overactive MAP kinase pathway resulting from genetic mutations.

As constitutive or overactivation of MAP kinase cascade plays a pivotal role in cell proliferation and differentiation, inhibition of this pathway is believed to be beneficial in hyperproliferative diseases. MEK is a key player in this pathway as it is downstream of Ras and Raf. Additionally, it is an attractive therapeutic target because the only known substrates for MEK phosphorylation are the MAP kinases, ERK1 and 2. Inhibition of MEK has been shown to have potential therapeutic benefit in several studies. For example, small molecule MEK inhibitors have been shown to inhibit human tumor


growth in nude mouse xenografts, (Sebolt-Leopold *et al.*, *Nature-Medicine* **1999**, *5* (7), 810-816; Trachet *et al.*, AACR April 6-10, 2002, Poster #5426; Tecle, H. IBC 2<sup>nd</sup> International Conference of Protein Kinases, September 9-10, 2002), block static allodynia in animals (WO 01/05390 published January 25, 2001) and inhibit growth of acute myeloid leukemia cells (Milella *et al.*, *J Clin Invest* **2001**, *108* (6), 851-859).

Small molecule inhibitors of MEK have been disclosed. At least thirteen patent applications have appeared in the last several years: US 5,525,625 filed January 24, 1995; WO 98/43960 published October 8, 1998; WO 99/01421 published January 14, 1999; WO 99/01426 published January 14, 1999; WO 00/41505 published July 20, 2000; WO 00/42002 published July 20, 2000; WO 00/42003 published July 20, 2000; WO 00/41994 published July 20, 2000; WO 00/42022 published July 20, 2000; WO 00/42029 published July 20, 2000; WO 00/68201 published November 16, 2000; WO 01/68619 published September 20, 2001; and WO 02/06213 published January 24, 2002.

## SUMMARY OF THE INVENTION

This invention provides for alkylated (1H-benzoimidazol-5-yl)-(4-substituted phenyl)-amine compounds of formula I, and pharmaceutically acceptable salts and prodrugs thereof that are useful in the treatment of hyperproliferative diseases. Specifically, the present invention relates to compounds of formula I that act as MEK inhibitors. Also provided is a method for treatment of cancer. Also provided are formulations containing compounds of formula I and methods of using the compounds to treat a patient in need thereof. In addition, there are described processes for preparing the inhibitory compounds of formula I.

Accordingly, the present invention provides compounds of the formula I:



I

and pharmaceutically accepted salts, prodrugs and solvates thereof, wherein:

— is an optional bond, provided that one and only one nitrogen of the ring is double-bonded;

$R^1$ ,  $R^2$ ,  $R^9$  and  $R^{10}$  are independently selected from hydrogen, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-OR^3$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $NR^4C(O)OR^6$ ,  $-OC(O)R^3$ ,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-NR^4C(O)R^3$ ,  $-C(O)NR^3R^4$ ,  $-NR^5C(O)NR^3R^4$ ,  $-NR^5C(NCN)NR^3R^4$ ,  $-NR^3R^4$ , and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, -S(O)<sub>j</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), -S(O)<sub>j</sub>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heteroaryl, -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heteroaryl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heterocyclyl and -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heterocyclyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

R<sup>3</sup> is selected from hydrogen, trifluoromethyl, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SR', -S(O)R''', -SO<sub>2</sub>R''', -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

R', R'' and R''' independently are selected from hydrogen, lower alkyl, lower alkenyl, aryl and arylalkyl;

R'''' is selected from lower alkyl, lower alkenyl, aryl and arylalkyl; or

Any two of R', R'', R''' or R'''' can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl; or

R<sup>3</sup> and R<sup>4</sup> can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl; or

R<sup>4</sup> and R<sup>5</sup> independently represent hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; or

R<sup>4</sup> and R<sup>5</sup> together with the atom to which they are attached form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R''', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl;

R<sup>6</sup> is selected from trifluoromethyl, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl, where each alkyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

R<sup>7</sup> is selected from hydrogen, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -SO<sub>2</sub>R<sup>6</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

W is selected from heteroaryl, heterocyclyl, -C(O)OR<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -C(O)NR<sup>4</sup>OR<sup>3</sup>, -C(O)R<sup>4</sup>OR<sup>3</sup>, -C(O)(C<sub>3</sub>-C<sub>10</sub> cycloalkyl), -C(O)(C<sub>1</sub>-C<sub>10</sub> alkyl), -C(O)(aryl), -C(O)(heteroaryl) and -C(O)(heterocyclyl), each of which is optionally substituted with 1-5 groups independently selected from

-NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, -R<sup>2</sup>, and



C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, and C<sub>2</sub>-C<sub>10</sub> alkynyl, each of which is optionally substituted with 1 or 2 groups independently selected from -NR<sup>3</sup>R<sup>4</sup> and -OR<sup>3</sup>;

R<sup>8</sup> is selected from hydrogen, -SCF<sub>3</sub>, -Cl, -Br, -F, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -OR<sup>3</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, -S(O)<sub>j</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), -S(O)<sub>j</sub>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heteroaryl, -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heteroaryl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heterocyclyl and -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heterocyclyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

m is 0, 1, 2, 3, 4 or 5; and

j is 1 or 2.

# DETAILED DESCRIPTION OF THE INVENTION

The novel compounds encompassed by the instant invention are those described by the general formula I set forth above, and the pharmaceutically acceptable salts and prodrugs thereof.


The present invention also provides compounds of formula I in which  $R^7$  is  $C_1$ - $C_{10}$  alkyl,  $C_3$ - $C_7$  cycloalkyl,  $C_3$ - $C_7$  cycloalkylalkyl,  $C_3$ - $C_7$  heterocycloalkyl or  $C_3$ - $C_7$  heterocycloalkylalkyl each of which can be optionally substituted with 1 - 3 groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $-OC(O)R^3$ ,  $-SO_2R^3$ ,  $-NR^4C(O)OR^6$ ,  $-NR^4C(O)R^3$ ,  $-C(O)NR^3R^4$ ,  $-NR^3R^4$ ,  $-NR^5C(O)NR^3R^4$ ,  $-NR^5C(NCN)NR^3R^4$ ,  $-OR^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl.

The present invention also provides compounds of formula I wherein  $R^8$  is  $-OCF_3$ ,  $-Br$  or  $-Cl$ ,  $R^2$  is hydrogen, and  $R^1$  is lower alkyl or halogen.

The present invention also provides compounds of formula I wherein  $R^9$  is hydrogen or halogen, and  $R^{10}$  is hydrogen.

The present invention also provides compounds of formula I wherein  $W$  is  $-C(O)OR^3$  or  $-C(O)NR^4OR^3$ .

The present invention also provides compounds of formula II:



II

wherein W, R<sup>1</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup> and R<sup>10</sup> are as defined above for formula I.


The present invention also provides compounds of formula II in which R<sup>7</sup> is C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl or C<sub>3</sub>-C<sub>7</sub> cycloalkylalkyl, each of which can be optionally substituted with 1 - 3 groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -SO<sub>2</sub>R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl.

The present invention also provides compounds of formula II wherein R<sup>8</sup> is -OCF<sub>3</sub>, -Br or -Cl, and R<sup>1</sup> is lower alkyl or halogen.

The present invention also provides compounds of formula II wherein R<sup>9</sup> is hydrogen or halogen, and R<sup>10</sup> is hydrogen.

The present invention also provides compounds of formula II wherein W is -C(O)OR<sup>3</sup> or -C(O)NR<sup>4</sup>OR<sup>3</sup>.

The present invention also provides compounds of formula III:



III

wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined above for formula I, and A is -OR<sup>3</sup> or -NR<sup>4</sup>OR<sup>3</sup>, wherein R<sup>3</sup> and R<sup>4</sup> are as defined above for formula I.


The present invention also provides compounds of formula III in which  $R^7$  is  $C_1$ - $C_{10}$  alkyl,  $C_3$ - $C_7$  cycloalkyl or  $C_3$ - $C_7$  cycloalkylalkyl, each of which can be optionally substituted with 1 - 3 groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $-OC(O)R^3$ ,  $-SO_2R^3$ ,  $-NR^4C(O)OR^6$ ,  $-NR^4C(O)R^3$ ,  $-C(O)NR^3R^4$ ,  $-NR^3R^4$ ,  $-NR^5C(O)NR^3R^4$ ,  $-NR^5C(NCN)NR^3R^4$ ,  $-OR^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl.

The present invention also provides compounds of formula III wherein  $R^8$  is  $-OCF_3$ ,  $-Br$  or  $-Cl$ ,  $R^2$  is hydrogen, and  $R^1$  is lower alkyl or halogen.

The present invention also provides compounds of formula III wherein  $R^9$  is hydrogen or halogen.

The present invention also provides compounds of formula III wherein  $R^3$  is hydrogen or lower alkyl when A is  $-OR^3$ ; and  $R^4$  is hydrogen when A is  $-NR^4OR^3$ .

The present invention also provides compounds of formula IIIa:



IIIa

wherein  $R^1$ ,  $R^2$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined above for formula I, and A is  $-OR^3$  or  $-NR^4OR^3$ , wherein  $R^3$  and  $R^4$  are as defined above for formula I.

The present invention also provides compounds of formula IIIa in which  $R^7$  is  $C_1$ - $C_{10}$  alkyl,  $C_3$ - $C_7$  cycloalkyl or  $C_3$ - $C_7$  cycloalkylalkyl, each of which can be optionally substituted with 1 - 3 groups independently selected from oxo, halogen, cyano, nitro,


trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-\text{NR}^4\text{SO}_2\text{R}^6$ ,  $-\text{SO}_2\text{NR}^3\text{R}^4$ ,  $-\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{OR}^3$ ,  $-\text{OC}(\text{O})\text{R}^3$ ,  $-\text{SO}_2\text{R}^3$ ,  $-\text{NR}^4\text{C}(\text{O})\text{OR}^6$ ,  $-\text{NR}^4\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{NCN})\text{NR}^3\text{R}^4$ ,  $-\text{OR}^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl.

The present invention also provides compounds of formula IIIa wherein  $\text{R}^8$  is  $-\text{OCF}_3$ ,  $-\text{Br}$  or  $-\text{Cl}$ ,  $\text{R}^2$  is hydrogen, and  $\text{R}^1$  is lower alkyl or halogen.

The present invention also provides compounds of formula IIIa wherein  $\text{R}^9$  is hydrogen or halogen.

The present invention also provides compounds of formula IIIa wherein  $\text{R}^3$  is hydrogen or lower alkyl when  $\text{A}$  is  $-\text{OR}^3$ ; and  $\text{R}^4$  is hydrogen when  $\text{A}$  is  $-\text{NR}^4\text{OR}^3$ .

The present invention also provides compounds of formula IIIb:



IIIb

wherein  $\text{R}^1$ ,  $\text{R}^7$ ,  $\text{R}^8$  and  $\text{R}^9$  are as defined above for formula I, and  $\text{A}$  is  $-\text{OR}^3$  or  $-\text{NR}^4\text{OR}^3$ , wherein  $\text{R}^3$  and  $\text{R}^4$  are as defined above for formula I.

The present invention also provides compounds of formula IIIb in which  $\text{R}^7$  is  $\text{C}_1$ - $\text{C}_{10}$  alkyl,  $\text{C}_3$ - $\text{C}_7$  cycloalkyl or  $\text{C}_3$ - $\text{C}_7$  cycloalkylalkyl, each of which can be optionally substituted with 1 - 3 groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-\text{NR}^4\text{SO}_2\text{R}^6$ ,  $-\text{SO}_2\text{NR}^3\text{R}^4$ ,  $-\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{OR}^3$ ,  $-\text{OC}(\text{O})\text{R}^3$ ,  $-\text{SO}_2\text{R}^3$ ,  $-\text{NR}^4\text{C}(\text{O})\text{OR}^6$ ,  $-\text{NR}^4\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ , -

$\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{NCN})\text{NR}^3\text{R}^4$ ,  $-\text{OR}^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl.

The present invention also provides compounds of formula IIIb wherein  $\text{R}^8$  is  $-\text{OCF}_3$ ,  $-\text{Br}$  or  $-\text{Cl}$ , and  $\text{R}^1$  is lower alkyl or halogen.

The present invention also provides compounds of formula IIIb wherein  $\text{R}^9$  is fluoro or chloro.

The present invention also provides compounds of formula IIIb wherein  $\text{R}^3$  is hydrogen or lower alkyl when A is  $-\text{OR}^3$ ; and  $\text{R}^4$  is hydrogen when A is  $-\text{NR}^4\text{OR}^3$ .

Except as expressly defined otherwise, the following definition of terms is employed throughout this specification.

By " $\text{C}_1\text{-C}_{10}$  alkyl", "alkyl" and "lower alkyl" in the present invention is meant straight or branched chain alkyl groups having 1-10 carbon atoms, such as, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, 3-methylpentyl, heptyl, octyl, and the like. Preferred alkyl radicals are  $\text{C}_{1-6}$  alkyl. More preferred alkyl radicals are  $\text{C}_{1-3}$  alkyl.

By " $\text{C}_2\text{-C}_{10}$  alkenyl", "lower alkenyl" and "alkenyl" means straight and branched hydrocarbon radicals having from 2 to 10 carbon atoms and at least one double bond and includes ethenyl, propenyl, 1-but-3-enyl, 1-pent-3-enyl, 1-hex-5-enyl and the like. More preferred are lower alkenyl having 3-5 carbon atoms.

By " $\text{C}_2\text{-C}_{10}$  alkynyl", "lower alkynyl" and "alkynyl" means straight and branched hydrocarbon radicals having from 2 to 10 carbon atoms and at least one triple bond and includes ethynyl, propynyl, butynyl, pentyn-2-yl and the like. More preferred are alkynyl having 3-5 carbon atoms.

By the term "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.

By "aryl" is meant an aromatic carbocyclic group having a single ring (e.g., phenyl), multiple rings (e.g., biphenyl), or multiple condensed rings in which at least one is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl), which is optionally mono-, di-, or trisubstituted with, e.g., halogen, lower alkyl, lower alkoxy, trifluoromethyl, aryl, heteroaryl, and hydroxy.

By "heteroaryl" is meant one or more aromatic ring systems of 5-, 6-, or 7-membered rings which includes fused ring systems (at least one of which is aromatic) of 5-10 atoms containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Examples of heteroaryl groups are pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, triazolyl, thiadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, and furopyridinyl. Spiro moieties are also included within the scope of this definition. Heteroaryl groups are optionally mono-, di-, or trisubstituted with, e.g., halogen, lower alkyl, lower alkoxy, haloalkyl, aryl, heteroaryl, and hydroxy.

As used herein, the term "carbocycle", "carbocyclyl", "cycloalkyl" or "C<sub>3</sub>-C<sub>10</sub> cycloalkyl" refers to saturated carbocyclic radicals having three to ten carbon atoms. The cycloalkyl can be monocyclic, or a polycyclic fused system, and can be fused to an aromatic ring. Examples of such radicals include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The cycloalkyl groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such cycloalkyl groups may be optionally substituted with, for example, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-

C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

By "heterocycle" or "heterocyclyl" is meant one or more carbocyclic ring systems of 5-, 6-, or 7-membered rings which includes fused ring systems of 4-10 atoms containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur, and with the proviso that the ring of the group does not contain two adjacent O or S atoms. A fused system can be a heterocycle fused to an aromatic group. Preferred heterocycles include, but are not limited to, pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidino, morpholino, thiomorpholino, thioxanyl, piperazinyl, homopiperazinyl, azetidiny, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepiny, diazepiny, thiazepiny, 1,2,3,6-tetrahydropyridiny, 2-pyrroliny, 3-pyrroliny, indoliny, 2H-pyrany, 4H-pyrany, dioxany, 1,3-dioxolany, pyrazoliny, dithianyl, dithiolany, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidiny, imidazoliny, 3-azabicyco[3.1.0]hexany, 3-azabicyclo[4.1.0]heptany, azabicyclo[2.2.2]hexany, 3H-indoly and quinoliziny. Spiro moieties are also included within the scope of this definition. The foregoing groups, as derived from the groups listed above, may be C-attached or N-attached where such is possible. For instance, a group derived from pyrrole may be pyrrol-1-yl (N-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole may be imidazol-1-yl (N-attached) or imidazol-3-yl (C-attached). An example of a heterocyclic group wherein 2 ring carbon atoms are substituted with oxo (=O) moieties is 1,1-dioxo-thiomorpholinyl. The heterocycle groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such heterocycle groups



may be optionally substituted with, for example, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

The term "arylalkyl" means an alkyl moiety (as defined above) substituted with one or more aryl moiety (also as defined above). More preferred arylalkyl radicals are aryl-C<sub>1-3</sub>-alkyls. Examples include benzyl, phenylethyl, and the like.

The term "heteroarylalkyl" means an alkyl moiety (as defined above) substituted with a heteroaryl moiety (also as defined above). More preferred heteroarylalkyl radicals are 5- or 6-membered heteroaryl-C<sub>1-3</sub>-alkyls. Examples include, oxazolylmethyl, pyridylethyl and the like.

The term "heterocyclalkyl" means an alkyl moiety (as defined above) substituted with a heterocycl moiety (also defined above). More preferred heterocyclalkyl radicals are 5- or 6-membered heterocycl-C<sub>1-3</sub>-alkyls. Examples include tetrahydropyranylmethyl.

The term "cycloalkylalkyl" means an alkyl moiety (as defined above) substituted with a cycloalkyl moiety (also defined above). More preferred heterocyclalkyl radicals are 5- or 6-membered cycloalkyl-C<sub>1-3</sub>-alkyls. Examples include cyclopropylmethyl.

The term "Me" means methyl, "Et" means ethyl, "Bu" means butyl and "Ac" means acetyl.

The phrase "pharmaceutically acceptable salt(s)", as used herein, unless otherwise indicated, includes salts of acidic and basic groups which may be present in the compounds of the present invention. The compounds of the present invention that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid

addition salts of such basic compounds of the present invention are those that form non-toxic acid addition salts, i.e., salts containing pharmaceutically acceptable anions, such as the acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edisylate, estolate, esylate, ethylsuccinate, fumarate, gluceptate, gluconate, glutamate, glycollylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodode, and valerate salts. Since a single compound of the present invention may include more than one acidic or basic moieties, the compounds of the present invention may include mono, di or tri-salts in a single compound.

In the case of an acidic moiety in a compound of the present invention, a salt may be formed by treatment of a compound of the present invention with a basic compound, particularly an inorganic base. Preferred inorganic salts are those formed with alkali and alkaline earth metals such as lithium, sodium, potassium, barium and calcium. Preferred organic base salts include, for example, ammonium, dibenzylammonium, benzylammonium, 2-hydroxyethylammonium, bis(2-hydroxyethyl)ammonium, phenylethylbenzylamine, dibenzyl-ethylenediamine, and the like salts. Other salts of acidic moieties may include, for example, those salts formed with procaine, quinine and N-methylglucosamine, plus salts formed with basic amino acids such as glycine, ornithine, histidine, phenylglycine, lysine and arginine. An especially preferred salt is a sodium or potassium salt of a compound of the present invention.

With respect to basic moieties, a salt is formed by the treatment of a compound of the present invention with an acidic compound, particularly an inorganic acid. Preferred inorganic salts of this type may include, for example, the hydrochloric, hydrobromic, hydroiodic, sulfuric, phosphoric or the like salts. Preferred organic salts of this type, may include, for example, salts formed with formic, acetic, succinic, citric, lactic, maleic, fumaric, palmitic, cholic, pamoic, mucic, D-glutamic, D-camphoric, glutaric, glycolic, phthalic, tartaric, lauric, stearic, salicylic, methanesulfonic, benzenesulfonic, paratoluenesulfonic, sorbic, puric, benzoic, cinnamic and the like organic acids. An especially preferred salt of this type is a hydrochloride or sulfate salt of a compound of the present invention.

In the compounds of the present invention, where terms such as  $(CR^4R^5)_m$  or  $(CR^4R^5)_t$  are used,  $R^4$  and  $R^5$  may vary with each iteration of  $m$  or  $t$  above 1. For instance, where  $m$  or  $t$  is 2, the terms  $(CR^4R^5)_m$  or  $(CR^4R^5)_t$  may equal  $-CH_2CH_2-$  or  $-CH(CH_3)C(CH_2CH_3)(CH_2CH_2CH_3)-$  or any number of similar moieties falling within the scope of the definitions of  $R^4$  and  $R^5$ .

Certain compounds of the present invention may have asymmetric centers and therefore exist in different enantiomeric forms. All optical isomers and stereoisomers of the compounds of the present invention, and mixtures thereof, are considered to be within the scope of the invention. With respect to the compounds of the present invention, the invention includes the use of a racemate, one or more enantiomeric forms, one or more diastereomeric forms, or mixtures thereof. The compounds of the present invention may also exist as tautomers. This invention relates to the use of all such tautomers and mixtures thereof.

The subject invention also includes isotopically-labeled compounds, which are identical to those recited in the present invention, but for the fact that one or more atoms

are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine and chloride, such as  $^2\text{H}$ ,  $^3\text{H}$ ,  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{18}\text{O}$ ,  $^{17}\text{O}$ ,  $^{31}\text{P}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{18}\text{F}$ , and  $^{36}\text{Cl}$ , respectively. Compounds of the present invention, prodrugs thereof, and pharmaceutically acceptable salts of said compounds or of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labeled compounds of the present invention, for example those into which radioactive isotopes such as  $^3\text{H}$  and  $^{14}\text{C}$  are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e.,  $^3\text{H}$  and carbon-14, i.e.,  $^{14}\text{C}$ , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e.,  $^2\text{H}$ , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labeled compound of the present invention and prodrugs thereof can generally be prepared by carrying out procedures disclosed in the Schemes and/or in the Examples and Preparations below, by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

This invention also encompasses pharmaceutical compositions containing a compound of formulas I-IIIb and methods of treating proliferative disorders, or abnormal cell growth, by administering prodrugs of compounds of the the present invention. Compounds of the present invention having free amino, amido, hydroxy or carboxylic groups can be converted into prodrugs. Prodrugs include compounds wherein an amino acid residue, or a polypeptide chain of two or more (e.g., two, three or four) amino acid

residues is covalently joined through an amide or ester bond to a free amino, hydroxy or carboxylic acid group of compounds of the present invention. The amino acid residues include but are not limited to the 20 naturally occurring amino acids commonly designated by three letter symbols and also includes 4-hydroxyproline, hydroxyllysine, demosine, isodemosine, 3-methylhistidine, norvaline, beta-alanine, gamma-aminobutyric acid, cirtulline, homocysteine, homoserine, ornithine and methionine sulfone. Additional types of prodrugs are also encompassed. For instance, free carboxyl groups can be derivatized as amides or alkyl esters. Free hydroxy groups may be derivatized using groups including but not limited to hemisuccinates, phosphate esters, dimethylaminoacetates, and phosphoryloxymethyloxycarbonyls, as outlined in *Advanced Drug Delivery Reviews* 1996, 19, 115. Carbamate prodrugs of hydroxy and amino groups are also included, as are carbonate prodrugs, sulfonate esters and sulfate esters of hydroxy groups. Derivatization of hydroxy groups as (acyloxy)methyl and (acyloxy)ethyl ethers wherein the acyl group may be an alkyl ester, optionally substituted with groups including but not limited to ether, amine and carboxylic acid functionalities, or where the acyl group is an amino acid ester as described above, are also encompassed. Prodrugs of this type are described in *J. Med. Chem.* 1996, 39, 10. Free amines can also be derivatized as amides, sulfonamides or phosphonamides. All of these prodrug moieties may incorporate groups including but not limited to ether, amine and carboxylic acid functionalities.

It is to be understood that in instances where two or more radicals are used in succession to define a substituent attached to a structure, the first named radical is considered to be terminal and the last named radical is considered to be attached to the structure in question. Thus, for example, the radical arylalkyl is attached to the structure in question by the alkyl group.

The invention also relates to a pharmaceutical composition for the treatment of a hyperproliferative disorder in a mammal which comprises a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a pharmaceutically acceptable carrier. In one embodiment, said pharmaceutical composition is for the treatment of cancer such as brain, lung, squamous cell, bladder, gastric, pancreatic, breast, head, neck, renal, kidney, ovarian, prostate, colorectal, esophageal, testicular, gynecological or thyroid cancer. In another embodiment, said pharmaceutical composition is for the treatment of a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (e.g., psoriasis), restenosis, or prostate (e.g., benign prostatic hypertrophy (BPH)).

The invention also relates to a pharmaceutical composition for the treatment of pancreatitis or kidney disease (including proliferative glomerulonephritis and diabetes-induced renal disease) or the treatment of pain in a mammal which comprises a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a pharmaceutically acceptable carrier.

The invention also relates to a pharmaceutical composition for the prevention of blastocyte implantation in a mammal which comprises a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a pharmaceutically acceptable carrier.

The invention also relates to a pharmaceutical composition for treating a disease related to vasculogenesis or angiogenesis in a mammal which comprises a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a pharmaceutically acceptable carrier. In one embodiment, said pharmaceutical composition is for treating a disease selected from

the group consisting of tumor angiogenesis, chronic inflammatory disease such as rheumatoid arthritis, atherosclerosis, inflammatory bowel disease, skin diseases such as psoriasis, excema, and scleroderma, diabetes, diabetic retinopathy, retinopathy of prematurity, age-related macular degeneration, hemangioma, glioma, melanoma, Kaposi's sarcoma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer.

The invention also relates to a method of treating a hyperproliferative disorder in a mammal that comprises administering to said mammal a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof. In one embodiment, said method relates to the treatment of cancer such as brain, lung, squamous cell, bladder, gastic, pancreatic, breast, head, neck, renal, kidney, ovarian, prostate, colorectal, esophageal, testicular, gynecological or thyroid cancer. In another embodiment, said method relates to the treatment of a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (e.g., psoriasis), restenosis, or prostate (e.g., benign prostatic hypertrophy (BPH)).

The invention also relates to a method for the treatment of a hyperproliferative disorder in a mammal that comprises administering to said mammal a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, in combination with an anti-tumor agent selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzyme inhibitors, topoisomerase inhibitors, biological response modifiers, anti-hormones, angiogenesis inhibitors, and anti-androgens.

The invention also relates to a method of treating pancreatitis or kidney disease in a mammal that comprises administering to said mammal a therapeutically effective

amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof.

The invention also relates to a method of preventing blastocyte implantation in a mammal that comprises administering to said mammal a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof.

The invention also relates to a method of treating diseases related to vasculogenesis or angiogenesis in a mammal that comprises administering to said mammal a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof. In one embodiment, said method is for treating a disease selected from the group consisting of tumor angiogenesis, chronic inflammatory disease such as rheumatoid arthritis, atherosclerosis, inflammatory bowel disease, skin diseases such as psoriasis, excema, and scleroderma, diabetes, diabetic retinopathy, retinopathy of prematurity, age-related macular degeneration, hemangioma, glioma, melanoma, Kaposi's sarcoma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer.

Patients that can be treated with compounds of the present invention, or pharmaceutically acceptable salts, prodrugs and hydrates of said compounds, according to the methods of this invention include, for example, patients that have been diagnosed as having psoriasis, restenosis, atherosclerosis, BPH, lung cancer, bone cancer, CMML, pancreatic cancer, skin cancer, cancer of the head and neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer, testicular, gynecologic tumors (e.g., uterine sarcomas, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina or carcinoma of the vulva), Hodgkin's disease, cancer



of the esophagus, cancer of the small intestine, cancer of the endocrine system (e.g., cancer of the thyroid, parathyroid or adrenal glands), sarcomas of soft tissues, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute leukemia, solid tumors of childhood, lymphocytic lymphomas, cancer of the bladder, cancer of the kidney or ureter (e.g., renal cell carcinoma, carcinoma of the renal pelvis), or neoplasms of the central nervous system (e.g., primary CNS lymphoma, spinal axis tumors, brain stem gliomas or pituitary adenomas).

This invention also relates to a pharmaceutical composition for inhibiting abnormal cell growth in a mammal which comprises an amount of a compound of the present invention, or a pharmaceutically acceptable salt or solvate or prodrug thereof, in combination with an amount of a chemotherapeutic, wherein the amounts of the compound, salt, solvate, or prodrug, and of the chemotherapeutic are together effective in inhibiting abnormal cell growth. Many chemotherapeutics are presently known in the art. In one embodiment, the chemotherapeutic is selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, angiogenesis inhibitors, and anti-androgens.

This invention further relates to a method for inhibiting abnormal cell growth in a mammal or treating a hyperproliferative disorder which method comprises administering to the mammal an amount of a compound of the present invention, or a pharmaceutically acceptable salt or solvate or prodrug thereof, in combination with radiation therapy, wherein the amounts of the compound, salt, solvate, or prodrug, is in combination with the radiation therapy effective in inhibiting abnormal cell growth or treating the hyperproliferative disorder in the mammal. Techniques for administering radiation therapy are known in the art, and these techniques can be used in the combination therapy

described herein. The administration of the compound of the invention in this combination therapy can be determined as described herein.

It is believed that the compounds of the present invention can render abnormal cells more sensitive to treatment with radiation for purposes of killing and/or inhibiting the growth of such cells. Accordingly, this invention further relates to a method for sensitizing abnormal cells in a mammal to treatment with radiation which comprises administering to the mammal an amount of a compound of the present invention or pharmaceutically acceptable salt or solvate or prodrug thereof, which amount is effective is sensitizing abnormal cells to treatment with radiation. The amount of the compound, salt, or solvate in this method can be determined according to the means for ascertaining effective amounts of such compounds described herein.

The invention also relates to a method of and to a pharmaceutical composition of inhibiting abnormal cell growth in a mammal which comprises an amount of a compound of the present invention, or a pharmaceutically acceptable salt or solvate thereof, a prodrug thereof, or an isotopically-labeled derivative thereof, and an amount of one or more substances selected from anti-angiogenesis agents, signal transduction inhibitors, and antiproliferative agents.

Anti-angiogenesis agents, such as MMP-2 (matrix-metalloproteinase 2) inhibitors, MMP-9 (matrix-metalloproteinase 9) inhibitors, and COX-II (cyclooxygenase II) inhibitors, can be used in conjunction with a compound of the present invention and pharmaceutical compositions described herein. Examples of useful COX-II inhibitors include CELEBREX<sup>TM</sup> (alecoxib), valdecoxib, and rofecoxib. Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 (published October 24, 1996), WO 96/27583 (published March 7, 1996), European Patent Application No. 97304971.1 (filed July 8, 1997), European Patent Application No. 99308617.2 (filed October 29,

1999), WO 98/07697 (published February 26, 1998), WO 98/03516 (published January 29, 1998), WO 98/34918 (published August 13, 1998), WO 98/34915 (published August 13, 1998), WO 98/33768 (published August 6, 1998), WO 98/30566 (published July 16, 1998), European Patent Publication 606,046 (published July 13, 1994), European Patent Publication 931,788 (published July 28, 1999), WO 90/05719 (published May 31, 1990), WO 99/52910 (published October 21, 1999), WO 99/52889 (published October 21, 1999), WO 99/29667 (published June 17, 1999), PCT International Application No. PCT/IB98/01113 (filed July 21, 1998), European Patent Application No. 99302232.1 (filed March 25, 1999), Great Britain Patent Application No. 9912961.1 (filed June 3, 1999), United States Provisional Application No. 60/148,464 (filed August 12, 1999), United States Patent 5,863,949 (issued January 26, 1999), United States Patent 5,861,510 (issued January 19, 1999), and European Patent Publication 780,386 (published June 25, 1997), all of which are incorporated herein in their entireties by reference. Preferred MMP-2 and MMP-9 inhibitors are those that have little or no activity inhibiting MMP-1. More preferred, are those that selectively inhibit MMP-2 and/or MMP-9 relative to the other matrix-metalloproteinases (i.e., MMP-1, MMP-3, MMP-4, MMP-5, MMP-6, MMP-7, MMP-8, MMP-10, MMP-11, MMP-12, and MMP-13).

Some specific examples of MMP inhibitors useful in the present invention are AG-3340, RO 32-3555, and RS 13-0830.

The terms "abnormal cell growth" and "hyperproliferative disorder" are used interchangeably in this application.

"Abnormal cell growth", as used herein, unless otherwise indicated, refers to cell growth that is independent of normal regulatory mechanisms (e.g., loss of contact inhibition). This includes, for example, the abnormal growth of: (1) tumor cells (tumors) that proliferate by expressing a mutated tyrosine kinase or overexpression of a receptor

tyrosine kinase; (2) benign and malignant cells of other proliferative diseases in which aberrant tyrosine kinase activation occurs; (3) any tumors that proliferate by receptor tyrosine kinases; (4) any tumors that proliferate by aberrant serine/threonine kinase activation; and (5) benign and malignant cells of other proliferative diseases in which aberrant serine/threonine kinase activation occurs.


The term "treating", as used herein, unless otherwise indicated, means reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment", as used herein, unless otherwise indicated, refers to the act of treating as "treating" is defined immediately above.

Representative compounds of the present invention, which are encompassed by the present invention include, but are not limited to the compounds of the examples and their pharmaceutically acceptable acid or base addition salts or prodrugs thereof.


The examples presented below are intended to illustrate particular embodiments of the invention, and are not intended to limit the scope of the specification or the claims in any way.

An illustration of the preparation of compounds of the present invention is shown in Schemes 1-4.


Scheme 1




Scheme 1a



Scheme 2




Scheme 3






Scheme 4



Scheme 5



General synthetic methods which may be referred to for preparing some of the compounds of the present invention are provided in PCT published application number WO 00/42022 (published July 20, 2000). The foregoing patent application is incorporated herein by reference in its entirety.

The examples presented below are intended to illustrate particular embodiments of the invention, and are not intended to limit the scope of the specification or the claims in any way.

An illustration of the preparation of compounds of the present invention is shown in Schemes 1-4.

Scheme 1 illustrates the synthesis of compounds of the present invention. In step 1, the acid is nitrated using standard conditions preferable fuming nitric acid in  $H_2SO_4$ . In step 2, the aniline is prepared by fluoride displacement with  $NH_4OH$  at room

temperature in water followed by careful acidification with concentrated mineral acid to pH near 0. In step 3, the ester is prepared by standard methods including by not limited to Fisher Esterification (MeOH, H<sub>2</sub>SO<sub>4</sub>), and reaction with TMSCHN<sub>2</sub> in suitable organic solvents like PhMe/MeOH or THF/MeOH. In step 4, the dianilino derivative is prepared by heating (60 to 200 °C) the ester with an excess of the appropriate aniline neat or in an organic solvent like xylenes. For example, when R<sup>1</sup> = Me and R<sup>2</sup> = H the preferred method is stirring the ester with 10 equivalents aniline in xylenes at reflux until complete reaction. In step 5, the nitro arene is reduced to produce the diamine by standard reduction conditions, including by not limited to H<sub>2</sub>, and Pd/C or Pd(OH)<sub>2</sub>/C or Raney Nickel in organic solvent like EtOH or THF, Fe in AcOH, Zn in AcOH or Zn, NH<sub>4</sub>Cl (aq) in MeOH. In step 6, the diamine is cyclization by heating with formic acid neat or formamidinium acetate in an appropriate solvent like EtOH. Alternatively, when R<sup>1</sup> or R<sup>2</sup> does not equal halo the nitro arene can be converted directly to the benzimidazole in step 7 by heating in formic acid with Pd(OH)<sub>2</sub>/C or other palladium source like Pd/C. In step 8, a halide can be incorporated by standard methods, including but not limited to NBS or NCS and pTsOH in organic cosolvents like THF and MeOH. In step 9, the benzimidazole is alkylated to give a near equal mixture of N1 and N3 products which are separable by standard techniques, including, for example, chromatography and trituration. The alkylation is accomplished by use of an alkylating agent like an alkyl halide and base like NaH, or K<sub>2</sub>CO<sub>3</sub> in suitable organic solvent like DMF or THF at temperatures ranging from 0 to 80 °C. R<sup>7</sup> can be further modified by various synthetic methods known in the art, as exemplified below. In step 10, the ester is hydrolyzed by standard saponification methods. The acid is then converted to the desired hydroxamate in step 11 by standard coupling procedures including but not limited to EDCI, HOBt or PyBOP and the

appropriate hydroxylamine in suitable organic solvents like DMF, THF or methylene chloride.

Scheme 2 illustrates an example in which the  $R^8$  substituent is on the aniline prior to the coupling procedure with the nitro ester. The reaction description is exactly like that for Scheme 1 except that there is no need to incorporate  $R^8$  as it is present in the aniline from the beginning.

In Scheme 3, the preparation of N3 alkyl amino benzimidazole derivatives is illustrated. In step 1, the terminal alkene of the N3 alkylated benzimidazole hydroxamate is dihydroxylated using a suitable oxidant like  $OsO_4$  in suitable solvent or  $KMnO_4$  or  $I_2$ ,  $AgOAc$ ,  $AcOH$ , water. The diol is then further oxidized in step 2 by  $NaIO_4$  or  $Pb(OAc)_4$  in suitable biphasic mixture to give the aldehyde. Alternatively (step 3), the alkene can be directly converted to the aldehyde by standard methods including but not limited to ozone/ $Me_2S$ ,  $NaIO_4/OsO_4$  or  $KMnO_4$ . In step 4, the amine is prepared by reductive amination using standard methods such as  $Na(CN)BH_3$ ,  $Na(OAc)_3BH$ ,  $NMe_4BH(OAc)_3$  with or without  $AcOH$  in a suitable solvent such as methylene chloride, acetonitrile or THF. The preferable reduction amination is to treat the aldehyde with amine,  $Me_4NBH(OAc)_3$  and acetic acid in  $MeCN$  at room temperature.

Scheme 4 illustrates the preparation of compounds of the present invention where W is heterocyclic. In step 1, the methyl ester is converted to the hydrazide by stirring with hydrazine in a suitable solvent like  $EtOH$  at temperatures from 50 to 100 °C. The desired heterocyclic derivative is then prepared by cyclization with the appropriate reagent. For oxadiazole 21 the hydrazide is treated with an orthoformate like triethyl orthoformate, and an acid catalyst like  $pTsOH$  in a suitable organic solvent like  $EtOH$  at elevated temperatures (50 – 100 °C). For hydroxy oxadiazole 22 the hydrazide can be cyclized with phosgene or a phosgene equivalent like triphosgene or carbonyl diimidazole

in a suitable organic solvent like toluene at temperatures ranging from 50 to 120 °C. The mercapto oxadiazole 23 can be prepared by reaction with carbon disulfide, and base like KOH in suitable organic solvent like EtOH at elevated temperatures (50 – 100 °C). The amino oxadiazole 24 can be made by reaction with BrCN and base like NaHCO<sub>3</sub>, in a suitable biphasic solvent system like dioxane and water at room temperature. Finally, the substituted amino oxadiazole 25 can be prepared by first reacting the hydrazide with an appropriate isothiocyanate in a suitable organic solvent like DMF or THF at temperatures ranging from 25 to 100 °C. The intermediate can be isolated or can be cyclized directly with the treatment of EDCI or other carbodiimide in suitable organic solvent like THF or DMF at temperatures ranging from room temperature to 80 °C.

In Scheme 5, the preparation of keto benzimidazole derivatives are illustrated. In step 1, the methyl ester is converted to the benzyl alcohol by standard reductive methods, preferably LAH in THF at 0 °C or NaBH<sub>4</sub> in EtOH:THF at room temperature. Oxidation to the aldehyde can be accomplished in step 2 using MnO<sub>2</sub> in acetone:THF at 50 °C. In step 3, organometallic reagents, such as organolithium reagents and Grignard reagents, can be added to the aldehyde in THF at low temperature (*e.g.*, -78 °C) to give the substituted benzyl alcohol. The keto derivatives can be prepared in step 4 by oxidation of the benzyl alcohol under standard conditions such as Swern or Dess-Martin oxidation.

The compounds of the present invention may have asymmetric carbon atoms. Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known to those skilled in the art, for example, by chromatography or fractional crystallization. Enantiomers can be separated by converting the enantiomer mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (*e.g.*, alcohol), separating the diastereomers and converting (*e.g.*, hydrolyzing) the individual diastereomers to the corresponding pure

enantiomers. All such isomers, including diastereomeric mixtures and pure enantiomers are considered as part of the invention.

The activity of the compounds of the present invention may be determined by the following procedure. N-terminal 6 His-tagged, constitutively active MEK1 (2-393) is expressed in *E. coli* and protein is purified by conventional methods (*Ahn et al., Science* 1994, 265, 966-970). The activity of MEK1 is assessed by measuring the incorporation of  $\gamma$ -<sup>33</sup>P-phosphate from  $\gamma$ -<sup>33</sup>P-ATP onto N-terminal His tagged ERK2, which is expressed in *E. coli* and is purified by conventional methods, in the presence of MEK1. The assay is carried out in 96-well polypropylene plate. The incubation mixture (100  $\mu$ L) comprises of 25 mM Hepes, pH 7.4, 10 mM MgCl<sub>2</sub>, 5 mM  $\beta$ -glycerolphosphate, 100  $\mu$ M Na-orthovanadate, 5 mM DTT, 5 nM MEK1, and 1  $\mu$ M ERK2. Inhibitors are suspended in DMSO, and all reactions, including controls are performed at a final concentration of 1% DMSO. Reactions are initiated by the addition of 10  $\mu$ M ATP (with 0.5  $\mu$ Ci  $\gamma$ -<sup>33</sup>P-ATP/well) and incubated at ambient temperature for 45 minutes. Equal volume of 25% TCA is added to stop the reaction and precipitate the proteins. Precipitated proteins are trapped onto glass fiber B filterplates, and excess labeled ATP washed off using a Tomtec MACH III harvester. Plates are allowed to air-dry prior to adding 30  $\mu$ L/well of Packard Microscint 20, and plates are counted using a Packard TopCount. In this assay, compounds of the invention exhibited an IC<sub>50</sub> of less than 50 micromolar.

The following compounds exemplify compounds of such activity.

Compound #
8n
11b
11c
11p
18i
29c
29i

29s
29t
29bb
29lll
29mmm

Administration of the compounds of the present invention (hereinafter the "active compound(s)") can be effected by any method that enables delivery of the compounds to the site of action. These methods include oral routes, intraduodenal routes, parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion), topical, and rectal administration.

The amount of the active compound administered will be dependent on the subject being treated, the severity of the disorder or condition, the rate of administration, the disposition of the compound and the discretion of the prescribing physician. However, an effective dosage is in the range of about 0.001 to about 100 mg per kg body weight per day, preferably about 1 to about 35 mg/kg/day, in single or divided doses. For a 70 kg human, this would amount to about 0.05 to 7 g/day, preferably about 0.05 to about 2.5 g/day. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several small doses for administration throughout the day.

The active compound may be applied as a sole therapy or may involve one or more other anti-tumor substances, for example those selected from, for example, mitotic inhibitors, for example vinblastine; alkylating agents, for example cis-platin, carboplatin and cyclophosphamide; anti-metabolites, for example 5-fluorouracil, cytosine arabinside and hydroxyurea, or, for example, one of the preferred anti-metabolites disclosed in European Patent Application No. 239362 such as N-(5-[N-(3,4-dihydro-2-methyl-4-

oxoquinazolin-6-ylmethyl)-N-methylamino]-2-thenoyl)-L-glutamic acid; growth factor inhibitors; cell cycle inhibitors; intercalating antibiotics, for example adriamycin and bleomycin; enzymes, for example, interferon; and anti-hormones, for example anti-estrogens such as Nolvadex<sup>TM</sup> (tamoxifen) or, for example anti-androgens such as Casodex<sup>TM</sup> (4'-cyano-3-(4-fluorophenylsulphonyl)-2-hydroxy-2-methyl-3'-(trifluoromethyl)propionanilide). Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate dosing of the individual components of treatment.

The pharmaceutical composition may, for example, be in a form suitable for oral administration as a tablet, capsule, pill, powder, sustained release formulations, solution, suspension, for parenteral injection as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository. The pharmaceutical composition may be in unit dosage forms suitable for single administration of precise dosages. The pharmaceutical composition will include a conventional pharmaceutical carrier or excipient and a compound according to the invention as an active ingredient. In addition, it may include other medicinal or pharmaceutical agents, carriers, adjuvants, etc.

Exemplary parenteral administration forms include solutions or suspensions of active compounds in sterile aqueous solutions, for example, aqueous propylene glycol or dextrose solutions. Such dosage forms can be suitably buffered, if desired.

Suitable pharmaceutical carriers include inert diluents or fillers, water and various organic solvents. The pharmaceutical compositions may, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. Thus for oral administration, tablets containing various excipients, such as citric acid may be employed together with various disintegrants such as starch, alginic acid and certain complex silicates and with binding agents such as sucrose, gelatin and acacia. Additionally,



lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tableting purposes. Solid compositions of a similar type may also be employed in soft and hard filled gelatin capsules. Preferred materials, therefore, include lactose or milk sugar and high molecular weight polyethylene glycols. When aqueous suspensions or elixirs are desired for oral administration the active compound therein may be combined with various sweetening or flavoring agents, coloring matters or dyes and, if desired, emulsifying agents or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin, or combinations thereof.

Methods of preparing various pharmaceutical compositions with a specific amount of active compound are known, or will be apparent, to those skilled in this art. For examples, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Ester, Pa., 15<sup>th</sup> Edition (1975).

The examples and preparations provided below further illustrate and exemplify the compounds of the present invention and methods of preparing such compounds. It is to be understood that the scope of the present invention is not limited in any way by the scope of the following examples and preparations. In the following examples molecules with a single chiral center, unless otherwise noted, exist as a racemic mixture. Those molecules with two or more chiral centers, unless otherwise noted, exist as a racemic mixture of diastereomers. Single enantiomers/diastereomers may be obtained by methods known to those skilled in the art.

The disclosures in this application of all articles and references, including patents, are incorporated herein by reference.


The invention is illustrated further by the following examples which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them.

The starting materials and various intermediates may be obtained from commercial sources, prepared from commercially available organic compounds, or prepared using well known synthetic methods.

Representative examples of methods for preparing intermediates of the invention are set forth below.

### Examples

#### Example 1



***7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-3H-benzoimidazole-5-carboxylic acid  
cyclopropylmethoxy-amide (11a)***

***Step A: 2,3,4-Trifluoro-5-nitro-benzoic acid 2***

A 3 liter three neck round bottom flask is charged with 125 ml H<sub>2</sub>SO<sub>4</sub>. Fuming nitric acid is added (8.4 ml, 199 mmol) and the mixture gently stirred. 2,3,4-Trifluorobenzoic acid 1 (25 g, 142 mmol) is added in 5 g portions over 90 minutes. The dark brownish yellow solution is stirred for 60 min at which time the reaction is complete. The reaction mixture is poured into 1 liter of an ice:water mixture and extracted with diethyl ether (3 x 600 ml). The combined organic extracts are dried (MgSO<sub>4</sub>) and concentrated under reduced pressure to give a yellow solid. The solid is suspended in hexanes and stirred for 30 min after which time it is filtered to give 29 g (92%) of clean desired product as an off-yellow solid: MS APCI (-) *m/z* 220 (M-1) detected.

***Step B: 4-Amino-2,3-difluoro-5-nitro-benzoic acid 3***

Ammonium hydroxide solution (~30% in water) (35 ml, 271 mmol) is added to a solution of 2,3,4-trifluoro-5-nitro-benzoic acid 2 (15 g, 67.8 mmol) in 30 ml water at 0 °C with stirring. Upon completion of ammonium hydroxide addition the reaction mixture is

warmed to room temperature with stirring. After 2.5 h, the reaction mixture is cooled to 0 °C and concentrated HCl is carefully added until pH of reaction mixture is near 0. The reaction mixture is diluted with water (30 ml) and extracted with diethyl ether (3 x 50 ml). The combined organic extracts are dried (MgSO<sub>4</sub>) and concentrated under reduced pressure to give 14 g (95%) of pure desired product: MS APCI (-) *m/z* 217 (M-1) detected.

*Step C: 4-Amino-2,3-difluoro-5-nitro-benzoic acid methyl ester 4*

A 2 M solution of TMS diazomethane in hexanes (6.88 ml, 13.75 mmol) is added to a suspension of 4-amino-2,3-difluoro-5-nitro-benzoic acid **3** (2.00 g, 9.17 mmol) in 25 ml of 4:1 THF:MeOH at 0 °C under nitrogen atmosphere. Upon completion of addition, reaction mixture is warmed to room temperature. After 0.5 h, excess TMS diazomethane is destroyed by the careful addition of acetic acid. The reaction is then concentrated under reduced pressure and dried *in vacuo* to give 1.95 g (92%) of pure desired product: MS APCI (-) *m/z* 231 (M-1) detected.

*Step D: 4-Amino-3-fluoro-5-nitro-2-o-tolylamino-benzoic acid methyl ester 5a*

4-Amino-2,3-difluoro-5-nitro-benzoic acid methyl ester **4** (12.0 g, 51.7 mmol) is suspended in xylenes (60 ml) and *ortho*-toluidine is added (55.2 ml, 517 mmol). The reaction mixture is heated to reflux with stirring under a nitrogen atmosphere. After 36 h, the reaction mixture is cooled to room temperature, diluted with diethyl ether and washed with 10% aqueous HCl solution. The aqueous washings are extracted with diethyl ether. The combined organic extracts are concentrated under reduced pressure. The residue is dissolved in methylene chloride and filtered through silica gel in a fritted funnel, rinsing

with methylene chloride. Three fractions are recovered. The first (2 liter) is nearly clean by HPLC. The second (1 liter) and third (1 liter) fractions are only partially pure. The first fraction is concentrated under reduced pressure and triturated with diethyl ether to give 11.2 g (68%) of clean desired product as a bright yellow solid: MS APCI (-)  $m/z$  318 (M-1) detected.

*Step E: 7-Fluoro-6-o-tolylamino-1H-benzoimidazole-5-carboxylic acid methyl ester 7a*

4-Amino-3-fluoro-5-nitro-2-o-tolylamino-benzoic acid methyl ester **5a** (1.57 g, 4.92 mmol), formic acid (25 ml, 26.5 mmol) and 20% Pd(OH)<sub>2</sub>/C (1.57 g, 2.95 mmol) in 25 ml EtOH are heating with stirring to 95 °C. After 16 h, the reaction mixture is cooled to room temperature and 0.5 g 20% Pd(OH)<sub>2</sub>/C and 10 ml formic acid added. The reaction mixture is heated to 95 °C with stirring. After 16 h, the reaction mixture is cooled to room temperature and filtered through Celite rinsing with EtOH. The filtrate is concentrated under reduced pressure until the desired product precipitates. The desired product is collected by filtration. The filtrate is concentrated again until more desired product precipitates. The product is collected by filtration. Repeated EtOH concentration, product filtration several times. Recovered 1.09 g (74%) pure desired product: MS APCI (+)  $m/z$  300 (M+1) detected; MS APCI (-)  $m/z$  298 (M-1) detected.

*Step F: 7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid methyl ester 8a*

7-Fluoro-6-o-tolylamino-1H-benzoimidazole-5-carboxylic acid methyl ester **7a** (2.00 g, 6.68 mmol) is suspended in 1:1 THF:MeOH mixture (60 ml) and cooled to -78 °C under a nitrogen atmosphere. A solution of NBS (1.20 g, 6.75 mmol) in 1:1

THF/MeOH (5 ml) is added followed by a MeOH (5 ml) solution of TsOH·H<sub>2</sub>O (1.9 g, 10.0 mmol). After 30 minutes, the reaction mixture is warmed to 0 °C and then after 1 h warmed to rt. After 16 h, more NBS (0.12 g, 0.67 mmol) is added and the reaction mixture is allowed to stir for 3 h. The reaction mixture is quenched by the addition of 10% Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> solution. After 30 min, the reaction mixture is diluted with water and ethyl acetate and the layers separated. The aqueous layer is extracted with ethyl acetate. The combined organic extracts are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The recovered solid is triturated with methylene chloride to give 2.00 g (79%) pure desired product: MS APCI (+) *m/z* 380, 378 (M+1 Br pattern) detected.

*Step G: 7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid 10a*


7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid methyl ester **8a** (63 mg, 0.167 mmol) is suspended in MeOH (1.5 ml) and 20% NaOH (400 µl) is added. After 16 h, the reaction mixture is cooled to 0 °C and 1 N HCl solution is added dropwise until pH is 2 to 3. The reaction mixture is diluted with ethyl acetate and water and the layers separated. The organic layer is washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give 58 mg (95%) of desired product: MS APCI (+) *m/z* 366, 364 (M+1 Br pattern) detected; MS APCI (-) *m/z* 364, 362 (M-1 Br pattern) detected.

*Step H: 7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide 11a*

7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid **10a**

(48 mg, 0.132 mmol) is dissolved in 1:1 THF: methylene chloride (1 ml) and Hunig's base (0.23  $\mu$ l, 1.32 mmol) is added followed by PyBOP (82 mg, 0.158 mmol). After a few minutes, cyclopropyl methyl hydroxylamine hydrochloride (20 mg, 0.158 mmol) (WO 0042022) is added. After the reaction is complete, the mixture is partitioned between methylene chloride and saturated  $\text{NaHCO}_3$  solution. The layers are separated and the organic layer is washed with saturated  $\text{NaHCO}_3$  and brine. The organic layer is dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. After purification by FCC (elute with 20:1 methylene chloride:MeOH), 25 mg (45%) of pure desired product is isolated: MS ESI (+)  $m/z$  435, 433 (M+1 Br pattern) detected; MS ESI (-)  $m/z$  433, 431 (M-1 Br pattern) detected;  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.15 (s, 1H), 8.02 (s, 1H), 7.28 (s, 1H), 7.43 (d, 1H), 7.07 (dd, 1H), 6.36 (m, 1H), 3.70 (d, 2H), 2.38 (s, 3H), 0.86 (m, 1H), 0.41 (m, 2H), 0.13 (m, 2H);  $^{19}\text{F}$  NMR (376MHz,  $\text{CDCl}_3$ ) -134.05 (s).

### Example 2



#### *7-Fluoro-6-phenylamino-3H-benzimidazole-5-carboxylic acid methyl ester (27a)*

##### *Step A: 4-Amino-3-fluoro-5-nitro-2-phenylamino-benzoic acid methyl ester 26a*


4-Amino-2,3-difluoro-5-nitro-benzoic acid methyl ester 4 (23.48 g, 101.1 mmol), the product of Example 1, Step C, is suspended in xylenes (125 mL) and aniline (92 mL, 1011 mmol) is added. The reaction mixture is stirred at 125  $^{\circ}\text{C}$  for 16 hours under  $\text{N}_2$ . The reaction mixture is cooled to room temperature and solids precipitate out of solution. The solids are collected by filtration and are washed with xylenes and then diethyl ether.

Recovered 22.22 g (72.78 mmol) of yellow solid which is pure desired product. The filtrate is concentrated under reduced pressure, redissolved in methylene chloride and flushed through a plug of silica gel eluting with methylene chloride. The desired fractions are concentrated under reduced pressure to give a brown solid which is triturated with diethyl ether to give 5.47 g (17.91 mmol) of yellow solid which is pure desired product. Combined product yield is 27.69 g (90%). MS APCI (-)  $m/z$  304 (M-1) detected.

*Step B: 7-Fluoro-6-phenylamino-3H-benzoimidazole-5-carboxylic acid methyl ester 27a*

4-Amino-3-fluoro-5-nitro-2-phenylamino-benzoic acid methyl ester 26a (16.70 g, 54.71 mmol), formic acid (250 mL, 6.63 mol) and 20% Pd(OH)<sub>2</sub>/C (9.00 g, 16.91 mmol) in ethanol (250 mL) are stirred at 40 °C for two hours under N<sub>2</sub> and then at 95 °C for 16 hours. The reaction mixture is cooled to room temperature and filtered through Celite rinsing with ethyl acetate. The filtrate is concentrated under reduced pressure to give a yellow solid. The solid is triturated with diethyl ether to give 13.47 g (86%) of the desired product as a tan solid. MS APCI (+)  $m/z$  286 (M+1) detected; MS APCI (-)  $m/z$  284 (M-1) detected.

### Example 3



***6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester (8b)***




*Step A: 6-(4-Bromo-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid methyl ester 28a*

7-Fluoro-6-phenylamino-3H-benzimidazole-5-carboxylic acid methyl ester **27a** (4.99 g, 17.51 mmol) is dissolved in *N,N*-dimethylformamide (275 mL). *N*-bromosuccinimide (3.15 g, 17.70 mmol) is added as a solid and the reaction mixture is stirred at room temperature under N<sub>2</sub>. After 30 min, the reaction mixture is quenched by the addition of aqueous saturated sodium bisulfite solution. The reaction mixture is then poured into a separatory funnel, diluted with water and ethyl acetate and the layers separated. The aqueous layer is extracted with ethyl acetate. The combined organic extracts are washed three times with water, once with brine and then are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to yield 6.38 g (100%) of the pure desired product as a tan solid. MS ESI (+) *m/z* 364, 366 (M+ Br pattern) detected.


*Step B: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid methyl ester 8b*

6-(4-Bromo-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid methyl ester **28a** (6.38 g, 17.51 mmol) is dissolved in *N,N*-dimethylformamide (275 mL). *N*-chlorosuccinimide (2.36 g, 17.70 mmol) is added as a solid and the reaction mixture is stirred at room temperature under N<sub>2</sub> until the reaction is complete (5-6 days). The reaction mixture is quenched by the addition of aqueous saturated sodium bisulfite solution to give a suspension. The resulting solids are collected by filtration, washed with water and diethyl ether and dried under reduced pressure to yield 6.07 g (87%) of the pure desired product as a beige solid. MS ESI (+) *m/z* 398, 400 (M+ Br pattern) detected.

**Example 4**

***6-(2,4-Dichloro-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid  
methyl ester (8c)***

7-Fluoro-6-phenylamino-3H-benzimidazole-5-carboxylic acid methyl ester **27a** (1.00 g, 3.51 mmol) is suspended in 1:1 tetrahydrofuran/methanol (20 mL) and cooled to  $-78^{\circ}\text{C}$  under  $\text{N}_2$ .  $\text{TsOH}\cdot\text{H}_2\text{O}$  (3.00 g, 10.50 mmol) is added followed by N-chlorosuccinimide (0.95 g, 7.08 mmol). After 10 minutes, the reaction mixture is warmed to  $0^{\circ}\text{C}$  to give a solution and then 30 min later warmed to room temperature. After stirring for 16 hours, the reaction is complete. The reaction mixture is quenched by the addition of aqueous saturated sodium bisulfite solution and diluted with ethyl acetate and water and the layers separated. The aqueous layer is extracted with ethyl acetate. The combined organic extracts are washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The resulting solid residue is triturated with methylene chloride to yield a white solid which is collected by filtration to yield 1.05 g (85%) of the pure desired product. MS ESI (+)  $m/z$  355, 357 ( $\text{M}^+$  Cl pattern) detected.

**Example 5**


***6-(4-Bromo-2-fluoro-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid***

**methyl ester (8d)****Step A: 4-Amino-3-fluoro-2-(2-fluoro-phenylamino)-5-nitro-benzoic acid methyl ester 5b**

4-Amino-2,3-difluoro-5-nitro-benzoic acid methyl ester **4** (1.50 g, 6.46 mmol) is suspended in xylenes (7.5 mL) and 2-fluoro-phenylamine (6.24 mL, 64.6 mmol) is added. The reaction mixture is stirred at 140 °C under N<sub>2</sub>. After stirring for 6 days, the reaction is complete. The reaction mixture is cooled to room temperature and diluted with methylene chloride and filtered through a silica gel plug eluting with methylene chloride (1L) to give an orange filtrate. The filtrate is concentrated to dryness and then triturated with diethyl ether to yield a bright yellow solid. The trituration is repeated. The yellow solid is collected to yield 1.08 g (52%) of the pure desired product. MS APCI (-) *m/z* 322 (M-1) detected.

**Step B: 6-(4-Bromo-2-fluoro-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid methyl ester 8d**

4-Amino-3-fluoro-2-(2-fluoro-phenylamino)-5-nitro-benzoic acid methyl ester **5b** is converted by the reduction/cyclization and bromination procedures already described to yield the desired product. MS ESI (+) *m/z* 382, 384 (M<sup>+</sup>, Br pattern) detected.


**Example 6****6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid methyl ester (8e)**

7-Fluoro-6-*o*-tolylamino-3H-benzimidazole-5-carboxylic acid methyl ester **7a** is

converted by the procedure already described for bromination, except *N*-chlorosuccinimide is used instead of *N*-bromosuccinimide, to yield the desired product.

MS ESI (+) *m/z* 334, 336 (*M*<sup>+</sup>, Cl pattern) detected.

### Example 7



#### ***7-Fluoro-6-(2-methyl-4-trifluoromethoxy-phenylamino)-3H-benzimidazole-5-carboxylic acid methyl ester (8f)***

***Step A. 4-Amino-3-fluoro-2-(2-methyl-4-trifluoromethoxy-phenylamino)-5-nitro-benzoic acid methyl ester 12a***

4-Amino-2,3-difluoro-5-nitro-benzoic acid methyl ester **4** (0.50 g, 2.15 mmol) is suspended in xylenes (3 mL) and 2-methyl-4-trifluoromethoxy-phenylamine (1.00 g, 5.23 mmol) is added. The reaction mixture is stirred at 140 °C under N<sub>2</sub>. After stirring for 7 days, the reaction is a mixture of starting material and product. The reaction mixture is cooled to room temperature. The reaction mixture is poured into a separatory funnel and diethyl ether and 10% aqueous HCl are added and the layers separated. The aqueous phase is extracted with three portions of diethyl ether. The combined diethyl ether layers are dried (MgSO<sub>4</sub>) and concentrated under reduced pressure. The residue is redissolved in methylene chloride and flushed through a plug of silica gel eluting with methylene chloride. The filtrate is concentrated under reduced pressure to give a bright yellow solid. The solid is washed with diethyl ether and the filtrate is concentrated under reduced pressure and the residue is further purified by FCC (eluting with 100% methylene chloride) to yield 0.39 g (45%) of the desired pure product as a yellow solid. MS APCI (-

)  $m/z$  402 (M-1) detected.

*Step B. 7-Fluoro-6-(2-methyl-4-trifluoromethoxy-phenylamino)-3H-benzoimidazole-5-carboxylic acid methyl ester 8f*

4-Amino-3-fluoro-2-(2-methyl-4-trifluoromethoxy-phenylamino)-5-nitro-benzoic acid methyl ester **12a** is converted by the reduction/cyclization procedure already described to yield the desired product. MS APCI (+)  $m/z$  384 (M+1) detected; MS APCI (-)  $m/z$  382 (M-1) detected.

### Example 8

#### *Preparation of Hydroxylamines*

Hydroxylamines useful for synthesizing compounds of the present invention may be prepared as follows

##### (i) **O-(2-Methoxy-ethyl)-hydroxylamine**

*Step A: 2-(2-Methoxy-ethoxy)-isoindole-1,3-dione*


DEAD (10 mL, 63 mmol) is added to a mixture of 2-methoxyethanol (5.0 mL, 63 mmol), PPh<sub>3</sub> (17 g, 63 mmol), and N-hydroxyphthalimide (10 g, 62 mmol) in THF (170 mL). The resulting orange solution is stirred 16 h at room temperature. The reaction mixture is concentrated in vacuo, and the solids are filtered washing with CHCl<sub>3</sub>. The filtrate is concentrated again, and the solids are filtered washing with CHCl<sub>3</sub>. This process is repeated until no precipitate forms. The final yellowish solids are recrystallized from EtOH to give the desired product (7.7 g, 55 %).

*Step B: O-(2-Methoxy-ethyl)-hydroxylamine*


To a solution of 2-(2-methoxy-ethoxy)-isoindole-1,3-dione (7.7 g, 35 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (30 mL) at room temperature is added methylhydrazine (2.0 mL, 36 mmol). The

resulting solution is stirred for 16 h at room temperature. The white solids are filtered off. The solvent is carefully distilled off under reduced pressure, then the concentrate is distilled under vacuum (20 torr, 57-58 °C) to afford the desired product (2.2 g, 68%).


(ii) The following hydroxylamines are prepared as described above using the appropriate alcohols. The isoindole-1,3-dione intermediates are purified by flash chromatography.




*O*-(2-Isobutoxy-ethyl)-hydroxylamine is used directly without purification.




*O*-(2-Pyrrolidin-1-yl-ethyl)-hydroxylamine is used directly without purification.




*O*-(2-Piperidin-1-yl-ethyl)-hydroxylamine is purified by Kugelrohr distillation (chamber temperature 140 °C, 1 torr).



*O*-(2-Methylsulfanyl-ethyl)-hydroxylamine is purified by vacuum distillation (76-78 °C, 20 torr).




*O*-(2-Phenylsulfanyl-ethyl)-hydroxylamine is used directly without purification.




*O*-(3-Methylsulfanyl-propyl)-hydroxylamine is used directly without purification.

(iii) The following hydroxylamines are prepared from the appropriate isoindole-1,3-dione by oxidation using oxone (*Tetrahedron Lett.* 1981, 22, 1287), and then


deprotection as described above.




*O*-(2-Methanesulfonyl-ethyl)-hydroxylamine is used directly without purification.




*O*-(2-Benzenesulfonyl-ethyl)-hydroxylamine is purified by flash chromatography (1% MeOH in CH<sub>2</sub>Cl<sub>2</sub>).



*O*-(3-Methanesulfonyl-propyl)-hydroxylamine is used directly without purification.



*O*-(3-Phenylsulfanyl-propyl)-hydroxylamine is prepared from PhSCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Br and N-hydroxyphthalimide by the patent procedure WO 0018790 and then is deprotected by the procedure described above and used directly without purification.



*O*-(3-Benzenesulfonyl-propyl)-hydroxylamine is prepared from the above isoindole-1,3-dione through its oxidation with oxone followed by deprotection as described above and is purified by flash chromatography (100% CH<sub>2</sub>Cl<sub>2</sub> to 2% MeOH in CH<sub>2</sub>Cl<sub>2</sub>).

**(v) *O*-(2-Morpholin-4-yl-ethyl)-hydroxylamine dihydrochloride**

*Step A: O*-(2-Bromo-ethyl)-hydroxylamine hydrobromide

2-(2-Bromo-ethoxy)-isoindole-1,3-dione is prepared from 1,2-dibromoethane and N-hydroxyphthalimide as described in WO 0018790, and is then subjected to the procedure in *J. Org. Chem.* **1963**, 28, 1604 to yield the desired product.

*Step B: (2-Bromo-ethoxy)-carbamic acid tert-butyl ester*

To a solution of O-(2-bromo-ethyl)-hydroxylamine hydrobromide (100 mg, 0.45 mmol) and di-*t*-butyl dicarbonate (110 mg, 0.49 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) at room temperature is added Et<sub>3</sub>N (0.08 mL, 0.56 mmol). The resulting suspension is stirred for 16 h room temperature. The reaction mixture is diluted with EtOAc, washed with 1 N aq HCl and brine, dried over MgSO<sub>4</sub>, filtered, concentrated, and purified by flash chromatography (100% CH<sub>2</sub>Cl<sub>2</sub>) to give the desired product (81 mg, 75%).

*Step C: (2-Morpholin-4-yl-ethoxy)-carbamic acid tert-butyl ester*

To a solution of (2-bromo-ethoxy)-carbamic acid tert-butyl ester (252 mg, 1.05 mmol) in DMF (2 mL) at room temperature is added morpholine (0.14 mL, 1.6 mmol). The reaction mixture is stirred for 7h at 50 °C. The reaction mixture is diluted with EtOAc, and washed with water. The organic layer is dried over MgSO<sub>4</sub>, filtered, concentrated, and purified by flash chromatography (2% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to give the desired product (118 mg, 46%): MS APCI (+) *m/z* 247 detected.

*Step D: O-(2-Morpholin-4-yl-ethyl)-hydroxylamine dihydrochloride*

To a solution of (2-morpholin-4-yl-ethoxy)-carbamic acid tert-butyl ester (118 mg, 0.48 mmol) in MeOH (1 mL) is added 4 M dioxane solution of HCl (2.4 mL, 9.60 mmol) at room temperature. The resulting solution is stirred for 16 h at room temperature. After addition of additional HCl (2.4 mL) followed by stirring for 4 h, the reaction mixture is concentrated *in vacuo* to give yellow solids (82 mg, 78%).


(vi) The isoindole-1,3-dione intermediates of the following hydroxylamines are prepared from the appropriate alkyl halide and N-hydroxyphthalimide by the procedure described within *J. Heterocyclic Chem.* 2000, 37, 827-830. The isoindole-1,3-diones are deprotected by the procedure described above: O-but-3-enyl-hydroxylamine; O-



(tetrahydro-furan-2-ylmethyl)-hydroxylamine; O-(3-methoxy-propyl)-hydroxylamine; and O-(3-benzyloxy-propyl)-hydroxylamine.

(vii) The following hydroxylamines are prepared as described in WO 0206213: O-(2-vinyloxy-ethyl)-hydroxylamine; 2-aminooxy-2-methyl-propan-1-ol; 1-aminooxy-2-methyl-propan-2-ol; 3-aminooxy-propan-1-ol; and (2-aminooxy-ethyl)-methyl-carbamic acid tert-butyl ester.

### Example 9



***6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid  
cyclopropylmethoxy-amide (11b)***

***Step A: 4-Amino-2-(2-chloro-phenylamino)-3-fluoro-5-nitro-benzoic acid methyl ester 5b***

4-Amino-2,3-difluoro-5-nitro-benzoic acid methyl ester **4** (2.00 g, 8.62 mmol) is suspended in xylenes (15 ml) and 2-chloro aniline (9.06 ml, 86.15 mmol) is added. The reaction mixture is heated to 140 °C under a nitrogen atmosphere. After 6 days, the reaction mixture is cooled to room temperature, and diluted with ethyl acetate. The reaction mixture is washed with water, 10% HCl solution and brine. The organic layer is dried (MgSO<sub>4</sub>) and concentrated under reduced pressure. The crude product is triturated with diethyl ether, twice, to give 0.35 g (12%) pure desired product as a brownish solid.

***Step B: 4,5-Diamino-2-(2-chloro-phenylamino)-3-fluoro-benzoic acid methyl ester 6a***

4-Amino-2-(2-chloro-phenylamino)-3-fluoro-5-nitro-benzoic acid methyl ester **5b**

(0.30 g, 0.88 mmol) is suspended in AcOH (5 ml) and zinc dust (0.29 g, 4.42 mmol) is added. After 15 minutes, the reaction is complete. The reaction mixture is diluted with ethyl acetate and filtered through Celite. The filtrate is washed with water, saturated NaHCO<sub>3</sub>, 10% K<sub>2</sub>CO<sub>3</sub> and brine. The organic layer is dried (MgSO<sub>4</sub>) and concentrated under reduced pressure to give 0.13 g (48%) pure desired product as an whitish brown foam.

*Step C: 6-(2-Chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester 7b*

4,5-Diamino-2-(2-chloro-phenylamino)-3-fluoro-benzoic acid methyl ester **6a** (0.125 g, 0.404 mmol) is suspended in EtOH (2 ml) and formamidine acetate (63 mg, 0.605 mmol) is added. The reaction mixture is heated to reflux. After 16 hours, the reaction mixture is cooled to rt and diluted with ethyl acetate. The organic layer is washed with water, saturated NaHCO<sub>3</sub>, 10% K<sub>2</sub>CO<sub>3</sub> and brine. The organic layer is dried (MgSO<sub>4</sub>) and concentrated under reduced pressure to give 0.109 g (85%) pure desired product.

*Step D: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester 8b*

6-(2-Chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester **7b** (55 mg, 0.172 mmol) is dissolved in 1:1 THF:MeOH (2 ml) and cooled to -78 °C under an atmosphere of nitrogen. TsOH·H<sub>2</sub>O (49 mg, 0.258 mmol) is added followed by NBS (31 mg, 0.174 mmol). After 10 minutes, the reaction mixture is warmed to 0 °C and then 2 hours later warmed to rt. After 16 hours, the reaction mixture is quenched by the

addition of 10%  $\text{Na}_2\text{S}_2\text{O}_3$  and diluted with ethyl acetate and water. The layers are separated and the aqueous layer is extracted with ethyl acetate. The combined organic extracts are dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure. The crude product is triturated with methylene chloride to give 58 mg (85%) of pure desired product as a tan solid.

*Step E: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid 10b*

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester **8b** (58 mg, 0.146 mmol) is suspended in EtOH (2 ml) and 1 ml 2 N NaOH is added. After 16 hours, the reaction mixture is diluted with ethyl acetate, water, and 10% HCl solution. The layers are separated and the organic layer is washed with brine. The organic layer is dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure. Trituration with MeOH gives 22 mg (39%) pure desired product.

*Step F: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide (11b)*


6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid **10b** (22 mg, 0.057 mmol) is dissolved in DMF (1 ml) and HOBt (9 mg, 0.062 mmol) followed by triethyl amine (18  $\mu\text{l}$ , 0.132 mmol) is added. Cyclopropyl methyl hydroxylamine hydrochloride (8 mg, 0.062 mmol) is added followed by EDCI (14 mg, 0.074 mmol). After 16 hours, the reaction mixture is diluted with ethyl acetate and water and the layers separated. The organic layer is washed with saturated  $\text{NH}_4\text{Cl}$ , brine, saturated  $\text{NaHCO}_3$ , water and brine. The organic layer is dried ( $\text{MgSO}_4$ ) and

concentrated under reduced pressure to give 23 mg (89%) pure desired product. MS APCI (+)  $m/z$  455, 453 ( $M+Br$  pattern) detected; MS APCI (-)  $m/z$  453, 451 ( $M-Br$  pattern) detected;  $^1H$  NMR (400MHz, DMSO- $d_6$ )  $\delta$  11.69 (broad s, 1H), 8.43 (s, 1H), 7.62 (d, 1H), 7.28 (dd, 1H), 6.42 (m, 1H), 3.63 (d, 2H), 1.03 (m, 1H), 0.48 (m, 2H), 0.19 (m, 2H);  $^{19}F$  NMR (376MHz, DMSO- $d_6$ ) -132.95 (s).

The following compounds are prepared by methods similar to those described in Example 1 and in this Example 9 by using the appropriate carboxylic acid and the appropriate hydroxylamine:

8g		8k	
8h		8l	
8i		8m	
8j		8n	

### Example 10



***6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide (29c)***

*Step A. 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid methyl ester 9a and 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-1-methyl-1H-benzoimidazole-5-carboxylic acid methyl ester*

A solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester **8b** (150 mg, 0.38 mmol), iodomethane (28  $\mu$ L, 0.45 mmol) and potassium carbonate (78 mg, 0.56 mmol) in dimethylformamide (1.5 mL) is stirred at 75 °C for one hour. The reaction mixture is diluted with ethyl acetate, washed with saturated aqueous potassium carbonate (2x), brine, and dried ( $\text{Na}_2\text{SO}_4$ ). Flash column chromatography (20:1 methylene chloride/ethyl acetate) provides 56 mg (36%) of the more mobile 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid methyl ester **9a** as a white solid.  $^{19}\text{F}$  NMR (376 MHz,  $\text{CD}_3\text{OD}$ ) -133.5 (s). MS APCI (+)  $m/z$  412, 414 ( $\text{M}^+$ , Br pattern) detected. Also isolated is 54 mg (35%) of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-1-methyl-1H-benzoimidazole-5-carboxylic acid methyl ester as a white solid.  $^{19}\text{F}$  NMR (376 MHz,  $\text{CD}_3\text{OD}$ ) -139.9 (s). MS APCI (+)  $m/z$  412, 414 ( $\text{M}^+$ , Br pattern) detected.

*Step B. 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid 10c*

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid methyl ester **9a** (56 mg, 0.14 mmol) is dissolved into 2:1 THF/water (3 mL) and NaOH (0.55 mL, 1.0 M aqueous solution, 0.55 mmol) is added. After stirring for two hours the reaction is reduced to one quarter initial volume via rotary evaporation and the remainder diluted to 50 mL with water. The aqueous solution is acidified to pH 2

by the addition of 1.0 M aqueous HCl and extracted with 1:1 tetrahydrofuran/ethyl acetate (3x), dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to provide 43 mg (79%) pure carboxylic acid as an off white solid. MS ESI (+) *m/z* 397, 398 (M<sup>+</sup>, Br pattern) detected.

*Step C: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-vinyloxy-ethoxy)-amide 29a*

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid **10c** (2.00 g, 5.0 mmol), O-(2-vinyloxy-ethyl)-hydroxylamine (0.776 g, 7.5 mmol), HOBt (0.88 g, 6.5 mmol), triethylamine (1.61 mL, 2.3 mmol) and EDCI (1.3 g, 6.5 mmol) are dissolved in dimethylformamide (52 mL) and stirred at room temperature for 48 hours. The reaction mixture is diluted with ethyl acetate, washed with water (3x), saturated potassium carbonate (2x), saturated ammonium chloride (2x), brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to an off-white solid. Trituration of the solid with diethyl ether provides 2.18 g (90%) desired product as an off-white solid. MS ESI (+) *m/z* 483, 485 (M<sup>+</sup> Br pattern) detected.

*Step D: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide 29c*















Hydrochloric acid (14 mL, 1.0 M aqueous solution, 14 mmol) is added to a suspension of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-vinyloxy-ethoxy)-amide **29a** (2.18 g, 4.50 mmol) in ethanol (50 mL) and the reaction mixture allowed to stir for 24 hours. The reaction mixture is concentrated to dryness by rotary evaporation and the solids partitioned between 3:1 ethyl

acetate/tetrahydrofuran and saturated potassium carbonate. The aqueous phase is extracted with 3:1 ethyl acetate/tetrahydrofuran (3x), the combined organics dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated to provide 2.11 g (100%) 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide as an off-white solid. MS ESI (+)  $m/z$  457, 459 ( $\text{M}^+$ , Br pattern) detected.  $^1\text{H}$  NMR (400 MHz,  $\text{MeOH-d}_4$ )  $\delta$  8.26 (s, 1H), 7.78 (s, 1H), 7.57 (d, 1H), 7.24 (dd, 1H), 6.40 (dd, 1H), 3.86 (s, 3H), 3.79 (m, 2H), 3.49 (m, 2H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{MeOH-d}_4$ ) -133.68 (s).

### Example 11

The following compounds are prepared by methods similar to those described in Example 10 by using methyl ester 8b and the appropriate alkylating agent (Step A) and the appropriate hydroxylamine (Step C):

29d		29ff	
29e		29gg	
29f		29hh	


29g		29ii	
29h		29jj	
29i		29kk	
29j		29ll	
29k		29mm	
29l		29nn	
29m		29oo	



29n		29pp	
29o		29qq	
29p		29rr	
29q		29ss	
29r		29tt	
29s		29uu	


29t		29vv	
29u		29ww	
29v		29xx	
29w		29yy	
29x		29zz	
29y		29aaa	

29z		29bbb	
29aa		29ccc	
29bb		29ddd	
29cc		29eee	
29dd		29fff	
29ee		29ggg	

**Example 12**

**6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2,3-dihydroxy-propoxy)-amide (29hhh)**

To a solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid allyloxy-amide **29tt** (20 mg, 0.04 mmol) in 0.50 mL 4:1 tetrahydrofuran/water is added  $\text{OsO}_4$  (41  $\mu\text{L}$ , 0.054 M solution in *t*-BuOH, 0.002 mmol) followed by NMO (7 mg, 0.06 mmol). The solution is stirred at room temperature for eight hours after which time HPLC analysis showed complete conversion to product. The solution is then stirred with saturated  $\text{NaHSO}_3$  and diluted with ethyl acetate. The organic phase is dried ( $\text{Na}_2\text{SO}_4$ ). Purification by FCC (DCM  $\rightarrow$  20:1 DCM/MeOH) provided 16 mg desired product as an off-white solid. MS ESI (+)  $m/z$  487, 489 ( $\text{M}^+$ , Br pattern) detected.


**Example 13**

**6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (3,4-dihydroxy-butoxy)-amide (29iii)**

6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-

carboxylic acid but-3-enyloxy-amide **29uu** is subjected to the dihydroxylation method described in Example 12. MS APCI (+)  $m/z$  501, 503 ( $M^+$  Br pattern) detected.

### Example 14





**6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzimidazole-5-carboxylic acid (2-methylamino-ethoxy)-amide TFA salt (29jjj)**

Prepared from (2-[[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzimidazole-5-carbonyl]-aminoxy]-ethyl)-methyl-carbamic acid tert-butyl ester **29ww** by trifluoroacetic acid deprotection in methylene chloride. MS APCI (+)  $m/z$  470, 472 ( $M^+$ , Br pattern) detected;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.31 (s, 1H), 7.74 (s, 1H), 7.51 (d, 1H), 7.19 (dd, 1H), 6.39 (dd, 1H), 4.11 (m, 2H), 3.97 (s, 3H), 3.12 (m, 2H), 2.72 (s, 3H);  $^{19}\text{F}$  NMR (376 MHz,  $\text{CD}_3\text{OD}$ ) -77.41 (s, 3F), -134.79 (s, 1F).

### Example 15


The following compounds are prepared by methods similar to those described in Example 10 by using methyl ester **8a** and the appropriate alkylating agent (Step A) and the appropriate hydroxylamine (Step C):







### Example 16

The following compounds are prepared by methods similar to those described in Example 10 by using methyl ester 8e and the appropriate alkylating agent (Step A) and the appropriate hydroxylamine (Step C):





### Example 17


The following compounds are prepared by methods similar to those described in Example 10 by using methyl ester 8c and the appropriate alkylating agent (Step A) and the appropriate hydroxylamine (Step C):

11m		29kkk	
-----	---	-------	--

**Example 18**

The following compounds are prepared by methods similar to those described in Example 10 by using methyl ester **8d** and the appropriate alkylating agent (Step A) and the appropriate hydroxylamine (Step C):

29lll		29mmm	
-------	---	-------	--

**Example 19**

**6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-[4-(4-methyl-piperazin-1-yl)-butyl]-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide (11o)**

**Step A:** 6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-pent-4-enyl-3H-benzimidazole-5-carboxylic acid methyl ester **9b**

7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-1H-benzimidazole-5-carboxylic acid methyl ester **8a** (0.915 g, 2.419 mmol) is suspended in DMF (18 ml) under an atmosphere of nitrogen. Bromopentene (0.430 ml, 3.629 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.502 g,

3.629 mmol) are added and the reaction mixture was warmed to 80 °C. After 1 hour, the reaction mixture is cooled to room temperature and poured into 100 ml of 1:1 ethyl acetate:diethyl ether. The organic layer is washed with water and brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The N3 and N1 alkylated products are separated by flash column chromatography, eluted with 20:1 methylene chloride:ethyl acetate. Complete separation of the isomers is obtained by performing two chromatographic separations. The higher  $R_f$  product is the N3 product **9b**, while the lower  $R_f$  product is the N1 product. The recovery of the N3 product **9b** is 0.415 g (38%): LC/MS ESI (+)  $m/z$  448, 446 (M+1 Br pattern) detected. The recovery of the N1 product was 0.486 g (45%): LC/MS ESI (+)  $m/z$  448, 446 (M+1 Br pattern) detected.

*Step B: 6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-pent-4-enyl-3H-benzoimidazole-5-carboxylic acid 10d*

6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-pent-4-enyl-3H-benzoimidazole-5-carboxylic acid methyl ester **9b** is dissolved in 1:1 THF:MeOH (10 ml) and 1 N NaOH solution (2.3 ml) is added. After 5h, the organic solvents are removed under reduced pressure and the residue diluted with water and 100 ml 1:1 THF:ethyl acetate. The layers are separated and the aqueous layer extracted with ethyl acetate. The combined organic extracts are dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure to afford 0.39 g (100%) clean desired product as a light yellow solid.

*Step C: 6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-pent-4-enyl-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide 11f*

6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-pent-4-enyl-3H-benzoimidazole-



5-carboxylic acid **10d** (0.390 g, 0.902 mmol) is dissolved in 1:1 THF:methylene chloride (6 ml) and Hunig's base (0.346 ml, 1.985 mmol) is added followed by PyBOP (0.563 g, 1.083 mmol). After 10 minutes, cyclopropyl methyl hydroxylamine hydrochloride (0.134 g, 1.083 mmol) is added. After 16 hours, the reaction mixture is diluted with ethyl acetate and washed with 0.1 N HCl, saturated NaHCO<sub>3</sub>, and brine. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The crude yellow residue is purified by FCC eluted with ethyl acetate to give 0.315 g (70%) pure desired product as a yellow solid: MS APCI (+) *m/z* 503, 501 (M+1 Br pattern) detected.

*Step D: 6-(4-Bromo-2-methyl-phenylamino)-3-(4,5-dihydroxy-pentyl)-7-fluoro-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide 11m*

6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-pent-4-enyl-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide **11f** (0.307 g, 0.612 mmol) is dissolved in 4:1 THF:water (8 ml) and 1.134 ml (0.061 mmol) of an 0.054 M OsO<sub>4</sub> solution in *t*-BuOH is added followed by NMO (0.093 g, 0.796 mmol). After 5h, the reaction mixture is quenched by the addition of 10% NaHS<sub>2</sub>O<sub>3</sub> solution. After 10 min, the reaction mixture is filtered through Celite rinsing with ethyl acetate and methylene chloride. The filtrate is diluted with ethyl acetate and washed with 0.01 N HCl, and brine. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The crude product is purified by FCC eluted with 9:1 ethyl acetate:MeOH to give 0.244 g (74%) pure desired product.

*Step E: 6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-(4-oxo-butyl)-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide 11n*

To a mixture of 6-(4-bromo-2-methyl-phenylamino)-3-(4,5-dihydroxy-pentyl)-7-fluoro-3*H*-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide **11m** (0.244 g, 0.456 mmol), THF (5 ml) and pH 7 phosphate buffer (3 ml) is added sodium periodate (0.195 g, 0.911 mmol). After 16 hours, the reaction mixture is diluted with ethyl acetate and washed with NaHCO<sub>3</sub>, and brine. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give an orange solid. Purification by FCC eluted with 4:1 methylene chloride:MeOH yields 0.189 g (82%) pure desired product as a yellow solid: MS APCI (+) *m/z* 505, 503 (M+1 Br pattern) detected; MS APCI (-) *m/z* 503, 501 (M-1 Br pattern) detected.

*Step F: 6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-[4-(4-methyl-piperazin-1-yl)-butyl]-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide 11o*

6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3-(4-oxo-butyl)-3*H* benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide **11n** (15 mg, 0.030 mmol) is dissolved in MeCN (500  $\mu$ l) and methylpiperazine (10  $\mu$ l, 0.089 mmol) is added followed by AcOH (5  $\mu$ l, 0.089 mmol). After 5 min, tetramethylammonium triacetoxyborohydride (12 mg, 0.045 mmol) is added. After 5 min, the reaction mixture is diluted with ethyl acetate and washed with NaHCO<sub>3</sub> and brine. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give 12 mg (69%) of pure desired product as a white solid. MS APCI (-) *m/z* 587, 585 (M-1 Br pattern) detected; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.99 (s, 1H), 7.98 (s, 1H), 7.30 (d, 1H), 7.08 (dd, 1H), 6.30 (d, 1H), 6.1 (broad singlet, 1H), 4.26 (t, 2H), 3.64 (d, 2H), 3.37 (s, 1H), 2.45 (broad, 8H), 2.41 (s, 3H), 2.38 (t, 2H), 2.28 (s, 3H), 1.95 (quin, 2H), 1.55 (quin, 2H), 0.98 (m, 1H), 0.50 (qt, 2H), 0.22 (qt, 2H).

**Example 20**


The following compounds are prepared by methods similar to those described in Example 19 by using the appropriate alkenyl substituted benzimidazole and the appropriate amine in the reductive amination (step F):

18a		18o	
18b		18p	
18c		18q	
18d		18r	
18e		18s	

18f		18t	
18g		18u	
18h		18v	
18i		18w	
18j		18x	
18k		18y	
18l		18z	

18m		18aa	
18n		18bb	


## Example 21



**6-(4-Bromo-2-methyl-phenylamino)-3-[4-(1,1-dioxo-1λ<sup>6</sup>-thiomorpholin-4-yl)-butyl]-7-fluoro-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide (18cc)**

To a solution of 6-(4-bromo-2-methyl-phenylamino)-7-fluoro-3-(4-thiomorpholin-4-yl-butyl)-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide **18l** (8 mg, 0.014 mmol) in 1:1:1 water/acetone/MeOH (1 mL) is added NMO (1.6 mg, 0.014 mmol) and osmium tetroxide (250  $\mu$ L, 0.054 M solution in t-BuOH, 0.014 mmol). After stirring for 24 hours, the solution is diluted with saturated sodium thiosulfate, stirred for 10 minutes and diluted with ethyl acetate. The solution is washed with brine (2x), dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure to a grey solid. FCC (10:1 dichloromethane/methanol) provides 6 mg (71%) desired product as an off-white solid. MS ESI (+)  $m/z$  622, 624 ( $\text{M}^+$ , Br pattern) detected.

## Example 22



**6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-[4-(4-methyl-piperazin-1-yl)-butyl]-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide (18dd)**

A solution of 6-(4-bromo-2-chloro-phenylamino)-3-(4-chloro-butyl)-7-fluoro-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide **18ee** (10 mg, 0.018 mmol), sodium iodide (14 mg, 0.092 mmol), and 1-methyl-piperazine (10  $\mu$ L, 0.092 mmol) are stirred at 85 °C for three hours. The reaction mixture is diluted with ethyl acetate and washed three times with water, washed twice with saturated aqueous potassium carbonate, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure to a yellow oil. Flash column chromatography (1:1 dichloromethane/methanol followed by methanol followed by 20:1 methanol/triethylamine) yields clean product (8 mg, 72%) as an off-white foam. MS ESI (+)  $m/z$  607, 609 ( $\text{M}^+$ , Br pattern) detected.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO-d}_6$ )  $\delta$  8.37 (s, 1H), 7.71 (s, 1H), 7.49 (d, 1H), 7.18 (dd, 1H), 6.40 (dd, 1H), 4.38 (t, 2H), 3.62 (d, 2H), 2.45 (broad, 8H), 2.41 (t, 2H), 2.28 (s, 3H), 1.96 (m, 2H), 1.54 (m, 2H), 1.07 (m, 1H), 0.50 (d, 2H), 0.22 (d, 2H).

**Example 23**


The following compounds are prepared by methods similar to those described in Example 22, using an appropriate amine and primary alkyl chloride.

18ff		18tt	
18gg		18uu	
18hh		18vv	
18ii		18ww	
18jj		18xx	

18kk		18yy	
18ll		18zz	
18mm		18aaa	
18nn		18bbb	
18oo		18ccc	
18pp		18ddd	



18qq		18eee	
18rr		18fff	
18ss			


**Example 24**

**6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3-oxazol-5-ylmethyl-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide (18ggg)**

6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3-(2-oxo-ethyl)-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide (0.020 g, 0.046 mmol) is dissolved in methanol (2 mL). Potassium carbonate (0.013 g, 0.093 mmol) and 1-isocyanomethanesulfonyl-4-methyl-benzene (0.010 g, 0.051 mmol) are added. The reaction mixture is stirred at reflux for 16 hours under N<sub>2</sub>, then concentrated under reduced pressure. The residue is dissolved in ethyl acetate and poured into a separatory funnel and washed with water and brine. The combined aqueous layers are reextracted

with ethyl acetate (2x). The combined ethyl acetate layers are dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The resulting solid was purified by flash column chromatography (eluting with 15:1 methylene chloride:methanol) to yield 0.011 g (50%) of the desired product. MS APCI (+)  $m/z$  470, 472 ( $\text{M}^+$ , Cl pattern) detected;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  10.51 (br s, 1H), 8.07 (s, 1H), 8.02 (s, 1H), 7.89 (s, 1H), 7.23 (s, 1H), 7.15 (d, 1H), 6.92 (dd, 1H), 6.31 (d, 1H), 6.11 (br s, 1H), 5.45 (s, 2H), 3.62 (d, 2H), 2.40 (s, 3H), 0.87 (m, 1H), 0.49 (m, 2H), 0.20 (m, 2H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ) – 134.54 (s).

### Example 25



***6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(3-oxo-3-pyrrolidin-1-yl-propyl)-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide (18hhh)***

**Step A:** 6-(4-Bromo-2-chloro-phenylamino)-3-(2-tert-butoxycarbonyl-ethyl)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester **8b** (0.50 g, 1.25 mmol) is dissolved in DMF (8 ml) under  $\text{N}_2$  and  $\text{K}_2\text{CO}_3$  (0.26 g, 1.88 mmol) is added followed by t-butyl acrylate (1.84 ml, 12.54 mmol). The reaction mixture is heated to 90 °C with stirring. After 4 h, the reaction mixture is cooled to rt and diluted with ethyl acetate. The organic layer is washed with water (3x) and brine, dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure. Purification by flash column chromatography eluted with 19:1 methylene chloride:ethyl acetate gives 0.41 g

(62%) desired product.

*Step B: 6-(4-Bromo-2-chloro-phenylamino)-3-(2-carboxy-ethyl)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester TFA salt*

6-(4-Bromo-2-chloro-phenylamino)-3-(2-tert-butoxycarbonyl-ethyl)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester (0.050 g, 0.095 mmol) is dissolved in methylene chloride (0.5 ml) and TFA (0.5 ml) is added. After 45 min, the reaction mixture is concentrated to dryness to give 0.49 g (88%) desired product: LC/MS ESI (+)  $m/z$  472, 470 ( $M^+$  Br pattern) detected;  $^1H$  NMR (400MHz, DMSO- $d_6$ )  $\delta$  8.51 (s, 1H), 8.20 (s, 1H), 8.13 (s, 1H), 7.64 (d, 1H), 7.29 (dd, 1H), 6.45 (dd, 1H), 4.55 (t, 2H), 2.89 (t, 2H).

*Step C: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(3-oxo-3-pyrrolidin-1-yl-propyl)-3H-benzoimidazole-5-carboxylic acid methyl ester*

To solution of 6-(4-bromo-2-chloro-phenylamino)-3-(2-carboxy-ethyl)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester (60 mg, 0.13 mmol) in DMF (1.8 mL) is added HOBt- $H_2O$  (24 mg, 0.16 mmol),  $Et_3N$  (0.043 mL, 0.31 mmol), pyrrolidine (0.011 mL, 0.13 mmol), and EDCI (34 mg, 0.18 mmol) at rt. The resulting yellow solution is stirred 16 h at rt. The reaction mixture is diluted with EtOAc and water, washed with sat'd aq  $NH_4Cl$ , brine, sat'd aq  $NaHCO_3$ , and brine. The organic layer is dried over  $MgSO_4$ , filtered, and concentrated *in vacuo* to give a crude material which is purified by flash chromatography (3% MeOH in  $CH_2Cl_2$ ) to afford 45 mg (67%) of the desired product: MS APCI (+)  $m/z$  523, 525 ( $M^+$ , Br pattern) detected.

*Step D: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(3-oxo-3-pyrrolidin-1-yl-propyl)-3H-benzoimidazole-5-carboxylic acid*

To a solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-(3-oxo-3-pyrrolidin-1-yl-propyl)-3H-benzoimidazole-5-carboxylic acid methyl ester (41 mg, 0.079 mmol) in THF/H<sub>2</sub>O (1.5 mL/0.75 mL) is added 0.20 mL (0.20 mmol) of 1 N aq LiOH at rt. The resulting solution is stirred 16 h. The reaction mixture is acidified with 1 N aq HCl (pH ~2 to 3) and diluted with EtOAc. The organic layer is dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo to give a crude product (42 mg) which is directly used without further purification.


*Step E: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(3-oxo-3-pyrrolidin-1-yl-propyl)-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide 18hhh*

The title compound is prepared from 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-(3-oxo-3-pyrrolidin-1-yl-propyl)-3H-benzoimidazole-5-carboxylic acid and *O*-cyclopropylmethyl-hydroxylamine hydrochloride by the standard coupling procedure described in Step A: MS APCI (+) *m/z* 578, 580 (M<sup>+</sup>, Br pattern) detected; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 11.66 (s, 1H), 8.42 (s, 1H), 8.01 (s, 1H), 7.76 (s, 1H), 7.62 (s, 1H), 7.28 (d, 1H), 6.39 (m, 1H), 4.52 (t, 2H), 3.66 (d, 2H), 3.33 (t, 2H), 3.28 (t, 2H), 2.89 (t, 2H), 1.83 (m, 2H), 1.76 (m, 2H), 1.06 (m, 1H), 0.49 (m, 2H), 0.22 (m, 2H); <sup>19</sup>F NMR (376 MHz, DMSO-*d*<sub>6</sub>) -132.94 (s, 1F).

**Example 26**

The following compounds are prepared by methods similar to those described in Example 25 using methyl ester **8b** and the appropriate amines:

18iii		18kkk	
18jjj			

**Example 27**

**6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide (11p)**

**Step A:** 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid methyl ester **11q**

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester **8b** (0.25 g, 0.63 mmol) is dissolved in N,N-dimethylformamide (5 mL). 2-Bromomethyl-tetrahydro-pyran (0.34 g, 1.88 mmol) and potassium carbonate (0.26 g, 1.88 mmol) are added and the reaction mixture is stirred at 60 °C for 12 hours under N<sub>2</sub>. The reaction mixture is poured into a separatory funnel, diluted with ethyl acetate and

water and the layers separated. The ethyl acetate layer is washed with water and brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The resulting solid residue is triturated with diethyl ether to yield a pale yellow solid (N3 regioisomer by NMR) and a yellow filtrate (mixture of N1 and N3 regioisomers by NMR). The solids are collected and washed with diethyl ether to yield 0.12 g (37%) of the pure desired N3 regioisomeric product as a pale yellow solid. MS ESI (+)  $m/z$  496, 498 ( $\text{M}^+$ , Br pattern) detected.

*Step B: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid 11r*

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid methyl ester **11q** is suspended in 4:1 tetrahydrofuran/water (2.5 mL) and aqueous 1 M LiOH is added (2.5 mL). After stirring at room temperature for 16 hours, the reaction mixture is homogeneous and the reaction is complete. The reaction mixture is cooled to 0 °C, diluted with water and aqueous 2 M HCl is added dropwise until the pH of the solution is 1-2, at which time it turns to a suspension. The reaction mixture is poured into a separatory funnel and diluted with ethyl acetate/tetrahydrofuran and water and the layers separated. The aqueous layer is extracted with ethyl acetate. The combined organic layers are washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure to yield 0.11 g (100%) of the pure desired product as a white solid. MS ESI (+)  $m/z$  482, 484 ( $\text{M}^+$ , Br pattern) detected.

*Step C: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid (2-vinyloxy-ethoxy)-amide 11s*

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-

3H-benzoimidazole-5-carboxylic acid **11r** (0.11 g, 0.23 mmol) is dissolved in N,N-dimethylformamide (2 mL). HOBT (0.037 g, 0.27 mmol) and triethylamine (0.094 mL, 0.68 mmol) are added. Then O-(2-vinyloxy-ethyl)-hydroxylamine (0.028 g, 0.27 mmol) and EDCI (0.056 g, 0.29 mmol) are added and the reaction mixture is stirred at room temperature under N<sub>2</sub> until HPLC shows the reaction is complete (2-3 days). The reaction mixture is poured into a separatory funnel, diluted with ethyl acetate and water and the layers separated. The ethyl acetate layer is washed successively with aqueous saturated NH<sub>4</sub>Cl (2x), brine (1x), aqueous saturated sodium bicarbonate (2x), water (1x), and brine (1x), dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The resulting solid is purified by FCC (eluting with 15:1 methylene chloride:methanol) to yield 0.039 g (79%) of the pure desired product as an off-white solid. MS ESI (+) *m/z* 567, 569 (M<sup>+</sup>, Br pattern) detected.

*Step D: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide 11p*

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid (2-vinyloxy-ethoxy)-amide **11s** (0.039 g, 0.068 mmol) is dissolved in ethanol (2 mL) and aqueous 2 M HCl (200 uL) is added. The reaction mixture is stirred at room temperature for 30 minutes. The reaction mixture is diluted with water and then neutralized with aqueous 2 M NaOH (~200 uL) until pH 7 and concentrated under reduced pressure. The residue is partitioned between ethyl acetate and brine in a separatory funnel and the layers separated. The ethyl acetate layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to yield 0.034 g (91%) of the pure desired product as an off-white solid. MS ESI (+) *m/z* 541, 543 (M<sup>+</sup>, Br pattern)


detected;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.29 (s, 1H), 7.75 (s, 1H), 7.49 (d, 1H), 7.18 (dd, 1H), 6.40 (dd, 1H), 4.40 (dd, A of ABX pattern, 1H), 4.28 (dd, B of ABX pattern, 1H), 3.92 (m, X of ABX pattern, 1H), 3.66 (t, 2H), 3.35 (m, 1H), 1.89 (m, 1H), 1.76 (m, 1H), 2.28 (s, 3H), 1.54 (m, 3H), 1.30 (m, 1H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CD}_3\text{OD}$ )  $-134.87$  (s).

### Example 28

The following compounds are prepared by methods similar to that described in Example 27 by using the appropriate methyl ester and alkylating agent (Step A) and the appropriate hydroxylamine in (Step C).

11t		11x	
11u		11y	
11v		11z	
11w		11aa	



**Example 29**

**6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide (11bb)**

**Step A:** 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzimidazole-5-carboxylic acid methyl ester **11cc**

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid methyl ester **8b** (1.55 g, 3.89 mmol) is dissolved in 15 ml DMF under N<sub>2</sub>. K<sub>2</sub>CO<sub>3</sub> (0.70 g, 5.06 mmol) is added followed by methyl vinyl sulfone (0.41 ml, 4.67 mmol). After stirring 16 h at room temperature, the reaction mixture is diluted with ethyl acetate and water. The layers are separated and the organic layer is washed with water (3x) and brine. The combined aqueous washes are extracted with ethyl acetate. The combined organic extracts are dried (MgSO<sub>4</sub>) and concentrated under reduced. Purification by dissolving the residue in methylene chloride and precipitating with diethyl ether, repeated several times, gives 1.16 g (59%) pure desired product as a yellow solid: MS APCI (+) *m/z* 506, 504 (M+ Br pattern) and 400, 398 (M – methyl ethyl sulfone Br pattern).

**Step B:** 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide **11bb**




6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzimidazole-5-carboxylic acid methyl ester **11cc** is subjected to methods previously described to give 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-

ethyl)-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide: MS APCI (+)  $m/z$  561, 559 (M+ Br pattern) and MS APCI (-)  $m/z$  559, 557 (M- Br pattern) detected;  $^1\text{H}$  NMR (400MHz, DMSO- $d_6$ )  $\delta$  11.75 (s, 1H), 8.47 (s, 1H), 8.04 (s, 1H), 7.77 (s, 1H), 7.62 (d, 1H), 7.28 (dd, 1H), 6.40 (dd, 1H), 4.78 (t, 2H), 3.82 (t, 2H), 3.62 (d, 2H), 3.07 (s, 3H), 1.02 (m, 1H), 0.49 (m, 2H), 0.21 (m, 2H);  $^{19}\text{F}$  NMR (376MHz, DMSO- $d_6$ ) -132.66 (s).


### Example 30

The following compounds were prepared similarly using the appropriate methyl ester and Michael acceptor and methods described previously.

11dd		11jj	
11ee		11kk	
11ff		11ll	
11gg		11mm	

11hh		11nn	
11ii			

### Example 31



***[6-(5-Amino-[1,3,4]oxadiazol-2-yl)-4-fluoro-1H-benzoimidazol-5-yl]-(4-bromo-2-methyl-phenyl)-amine (24a)***


**Step A:** 6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid hydrazide **20a**

6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester **8a** (0.051 g, 0.135 mmol) is suspended in EtOH (5 ml) and hydrazine hydrate (0.118 g, 2.023 mmol) is added. The reaction mixture is heated at reflux for 16 hours. The reaction mixture is concentrated under reduced pressure and purified by FCC eluted with 97:3 ethyl acetate:MeOH to give 0.041 g (81%) of clean desired product: LC/MS ESI (+) *m/z* 378, 380 (M+ Br pattern) detected.

**Step B:** *[6-(5-Amino-[1,3,4]oxadiazol-2-yl)-4-fluoro-1H-benzoimidazol-5-yl]-(4-bromo-2-methyl-phenyl)-amine 24a*


6-(4-Bromo-2-methyl-phenylamino)-7-fluoro-3*H*-benzoimidazole-5-carboxylic acid hydrazide **20a** (0.041 g, 0.109 mmol) is suspended in 1,4-dioxane (1.5 ml) and 36  $\mu$ l of a 3 M solution of cyanogen bromide in methylene chloride is added.  $\text{NaHCO}_3$  (9 mg, 0.109 mmol) in water (1.5 ml) is then added. After 16 hours, the reaction mixture is diluted with water and brine and extracted with THF. The combined organic extracts are dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. Purification by FCC eluted with 98:2 ethyl acetate:MeOH gives 24 mg (55%) of pure desired product as a yellow solid: LC/MS ESI (+)  $m/z$  403, 405 ( $\text{M}^+$  Br pattern) detected;  $^1\text{H}$ -NMR (400 MHz,  $\text{DMSO-d}_6$ )  $\delta$  12.97 (s, 1H), 8.42 (s, 1H), 7.94 (s, 1H), 7.74 (s, 1H), 7.36 (s, 2H), 7.33 (d, 1H), 7.15 (d, 1H), 6.40 (bs, 1H), 2.34 (s, 3H).

### Example 32




***[6-(5-Amino-[1,3,4]oxadiazol-2-yl)-4-fluoro-1*H*-benzoimidazol-5-yl]-(4-chloro-2-methyl-phenyl)-amine (24b)***

[6-(5-Amino-[1,3,4]oxadiazol-2-yl)-4-fluoro-1*H*-benzoimidazol-5-yl]-(4-chloro-2-methyl-phenyl)-amine **24b** is prepared as described in example 31 starting with 6-(4-chloro-2-methyl-phenylamino)-7-fluoro-3*H*-benzoimidazole-5-carboxylic acid methyl ester **8e**. LC/MS ESI (+)  $m/z$  359, 361 ( $\text{M}^+$  Cl pattern) detected;  $^1\text{H}$  NMR (400MHz,  $\text{DMSO-d}_6$ )  $\delta$  8.42 (s, 1H), 8.00 (bs, 1H), 7.78 (bs, 1H) 7.48 (s, 2H), 7.22 (s, 1H), 7.04 (d, 1H), 6.48 (bs, 1H), 2.37 (s, 3H).

**Example 33**

***[6-(5-Amino-[1,3,4]oxadiazol-2-yl)-4-fluoro-1H-benzoimidazol-5-yl]-(4-bromo-2-chloro-phenyl)-amine (24c)***

[6-(5-Amino-[1,3,4]oxadiazol-2-yl)-4-fluoro-1H-benzoimidazol-5-yl]-(4-bromo-2-chloro-phenyl)-amine **24c** is prepared as described in example 31 starting with 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester **8b**. MS APCI (+)  $m/z$  425, 423 (M+ Br pattern) and MS APCI (-)  $m/z$  423, 421 (M- Br pattern) detected.


**Example 34**

***6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid hydrazide (20b)***

6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid hydrazide **20b** is prepared as described in example 31, step A from 6-(4-chloro-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methyl ester **8e**. LC/MS ESI (+)  $m/z$  334, 336 (M+ Cl pattern) detected;  $^1\text{H}$  NMR (400MHz, DMSO- $d_6$ )


$\delta$  13.09 (bs, 1H), 9.98 (s, 1H), 8.40 (s, 1H), 8.17 (bs, 1H), 7.64 (bs, 1H), 7.20 (s, 1H), 7.03 (d, 1H), 6.41 (bs, 1H), 4.49 (s, 2H), 2.23 (s, 3H).

### Example 35




**5-[6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-  
[1,3,4]oxadiazol-2-ol (22a)**

6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid hydrazide **20b** (0.050 g, 0.150 mmol) is suspended in PhMe (2 ml) and a 20% phosgene solution in PhMe (0.24 ml, 0.45 mmol) is added. The reaction mixture is stirred at reflux under N<sub>2</sub> for 1h and then cooled to rt. The reaction mixture is quenched by the addition of a 1:1 mixture of THF and 10% HCl (20 ml). The layers are separated and the aqueous layer is extracted with THF (3x). The combined organic layer is washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give 54 mg (99%) of desired product as a yellow solid: LC/MS ESI (+) *m/z* 360, 362 (M+ Cl pattern) detected; <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>)  $\delta$  12.64 (s, 1H), 8.83 (s, 1H), 7.88 (s, 1H), 7.30 (s, 1H), 7.20 (d, 1H), 7.00 (dd, 1H), 6.38 (dd, 1H), 2.30 (s, 3H).

**Example 36**

**(4-Chloro-2-methyl-phenyl)-(4-fluoro-6-[1,3,4]oxadiazol-2-yl-1H-benzoimidazol-5-yl)-  
amine (21a)**


6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid hydrazide **20b** (0.048 g, 0.144 mmol) is suspended in 3 ml absolute EtOH and HC(OEt)<sub>3</sub> (0.60 ml, 3.54 mmol) is added followed by catalytic pTsOH·H<sub>2</sub>O. The reaction mixture is heated to reflux under N<sub>2</sub>. After 2h, the reaction mixture is cooled to room temperature and concentrated under reduced pressure. Purification by flash column chromatography (97:3 ethyl acetate:MeOH) gives 36 mg (73%) desired product as a light yellow solid. LC/MS ESI (+) *m/z* 344, 346 (M+ Cl pattern) detected; <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) δ 13.10 (bs, 1H), 9.39 (s, 1H), 8.49 (s, 1H), 8.10 (bs, 1H), 7.78 (bs, 1H), 7.20 (d, 1H), 7.00 (dd, 1H), 6.41 (bs, 1H), 2.18 (s, 3H).

**Example 37**

**5-[6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-  
[1,3,4]oxadiazole-2-thiol (23a)**

6-(4-Chloro-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid hydrazide **20b** (0.050 g, 0.150 mmol) is suspended in 3 ml absolute EtOH and cooled to 0 °C under N<sub>2</sub>. CS<sub>2</sub> is added (26 mg, 0.346 mmol) followed by powdered KOH (8 mg, 0.150 mmol). After stirring at 0 °C for 30 min, the reaction mixture is heated to reflux. After 3.5 h, the reaction mixture is quenched by the addition of water, followed by the addition of ethyl acetate and 1N HCl. The layers are separated and the aqueous layer is extracted with ethyl acetate. The combined organic extracts are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give the desired product as a yellow solid: LC/MS ESI (+) *m/z* 376, 378 (M+ Cl pattern) detected; <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) δ 8.51 (s, 1H), 7.92 (s, 1H), 7.19 (s, 1H), 7.12 (s, 1H), 6.98 (d, 1H), 6.29 (d, 1H), 2.28 (s, 3H).

### Example 38



**6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid methylamide (1100)**

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid **10c** (0.029 g, 0.076 mmol) is dissolved in N,N-dimethylformamide (1.1mL). HOBT (0.016 g, 0.10 mmol), triethylamine (0.028 mL, 0.20 mmol), methylamine (0.059 mL, 0.12 mmol, 2 M solution in tetrahydrofuran), and EDCI (0.019 g, 0.10 mmol) are added consecutively to the reaction mixture at room temperature. The solution is stirred at room temperature for 16 hours under N<sub>2</sub>. The reaction mixture is poured into a separatory funnel and diluted with ethyl acetate and water and the layers separated. The ethyl acetate layer is washed successively with aqueous saturated NH<sub>4</sub>Cl (2x), brine (1x), aqueous




saturated sodium bicarbonate (2x), water (1x), and brine (1x), dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure. The resulting solid is purified by FCC (eluting with 19:1 methylene chloride:methanol) to yield 0.013 g (42%) of the pure desired product. MS APCI (+)  $m/z$  397, 399 ( $\text{M}^+$ , Br pattern) detected;  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO-d}_6$ )  $\delta$  8.76 (broad s, 1H), 8.69 (m, 1H), 8.41 (s, 1H), 7.76 (s, 1H), 7.63 (d, 1H), 7.30 (dd, 1H), 6.50 (dd, 1H), 2.76 and 2.75 (s and s, 3H total, amide rotamers).  $^{19}\text{F}$  NMR (376 MHz,  $\text{DMSO-d}_6$ ) -132.69 (s).

### Example 39


The following compounds are prepared using methods similar to that described above in Example 38 by using the appropriate carboxylic acid and amine. In those cases that contain two amine functionalities, the appropriate mono Boc protected amine is used in the coupling reaction and the Boc group is later removed in a final step under standard TFA deprotection conditions.

11pp		11uu	
11qq		11vv	
11rr		11ww	

11ss		11xx	
11tt			


**Example 40*****[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzimidazol-5-yl]-methanol(10e)***

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid methyl ester **8b** (1.06 g, 2.65 mmol) is suspended in tetrahydrofuran (25 mL) and cooled to -78 °C. Lithium aluminum hydride (8.03 mL, 8.03 mmol, 1M solution in tetrahydrofuran) is added dropwise to the reaction mixture. After stirring for 10 minutes at -78 °C, the reaction mixture is warmed to 0 °C and becomes a homogeneous solution. The reaction mixture is stirred for 5 minutes at 0 °C and then cooled again to -78 °C. The reaction mixture is quenched with MeOH, diluted with Rochelle's salt, warmed to room temperature and stirred for 1 hour. The reaction mixture is then poured into a separatory funnel, diluted with ethyl acetate, and the layers separated. The aqueous phase is extracted with ethyl acetate. The combined ethyl acetate layers are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to yield 0.98 g (100%) of the pure desired product as a pale yellow solid. MS ESI (+) *m/z* 370, 372 (M<sup>+</sup>, Br pattern) detected.

**Example 41****6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzimidazole-5-carbaldehyde (10f)**

[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzimidazol-5-yl]-methanol

10e (0.96 g, 2.58 mmol) is dissolved in tetrahydrofuran/acetone (1:1, 15 mL), and  $\text{MnO}_2$  (2.24 g, 25.8 mmol) is added. The reaction mixture is stirred at 50 °C for 10 hours under  $\text{N}_2$ . The reaction mixture is filtered through silica gel and eluted with methylene chloride/methanol (10:1, 1 L). The filtrate is concentrated under reduced pressure to a small volume and then filtered through an Acrodisc syringe filter to remove small amounts of  $\text{MnO}_2$  that passed through the silica gel. The filtrate is concentrated under reduced pressure and the residue is purified by flash column chromatography (eluting with 20:1 methylene chloride:methanol) to yield 0.81 g (85%) of the pure desired product as a bright yellow solid. MS ESI (+)  $m/z$  368, 370 ( $\text{M}^+$ , Br pattern) detected.

**Example 42**

**1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzimidazol-5-yl]-2-hydroxy-ethanone (10g)**

*Step A: 1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzimidazol-5-yl]-2-methoxymethoxy-ethanol 10i*

To a solution of tributyl-methoxymethoxymethyl-stannane (864 mg, 2.37 mmol, prepared by the procedure reported in *J. Org. Chem.* **1988**, *53*, 4131) in THF (8 mL) at -78 °C is added n-BuLi (0.94 mL, 2.35 mmol, 2.5 M solution in hexane). After stirring for 3 min, a solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carbaldehyde **10h** (59 mg, 0.15 mmol) in THF (2 mL) is added at -78 °C. After stirring for 40 min at -78 °C, the reaction is quenched with saturated aqueous NH<sub>4</sub>Cl at -78 °C, warmed to room temperature, and diluted with EtOAc. The organic layer is washed with brine, dried over MgSO<sub>4</sub>, filtered, concentrated, and purified by flash chromatography (1.5 % MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to give the desired product (45 mg, 64%): MS APCI (+) *m/z* 458, 460 (M<sup>+</sup>, Br pattern) detected.


*Step B: 1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazol-5-yl]-2-methoxymethoxy-ethanone 10j*

A solution of 1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazol-5-yl]-2-methoxymethoxy-ethanol **10i** (44 mg, 0.096 mmol) and the Dess-Martin periodinane (49 mg, 0.12 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1.5 mL) is stirred for 1.5 h at rt. The reaction mixture is diluted with ether (3 mL). Saturated aqueous NaHCO<sub>3</sub> (1 mL) containing sodium thiosulfate pentahydrate (74 mg) is added. The resulting mixture is stirred for 10 min and diluted with EtOAc. The organic layer is washed with saturated aqueous NaHCO<sub>3</sub> and brine, dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo* to give a crude material which is purified by flash chromatography (1.5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to afford the desired product (31 mg, 71%): MS APCI (+) *m/z* 456, 458 (M<sup>+</sup>, Br pattern) detected.

*Step C: 1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazol-5-yl]-2-hydroxy-ethanone 10g*

A mixture of 1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazol-5-yl]-2-methoxymethoxy-ethanone **10j** (15 mg, 0.033 mmol), 10% aqueous HCl (0.3 mL), methanol (0.01 mL), and water (0.05 mL) is stirred for 3 days at rt. The reaction mixture is neutralized with saturated aqueous NaHCO<sub>3</sub>, and diluted with EtOAc. The organic layer is washed with brine, dried over MgSO<sub>4</sub>, filtered, concentrated *in vacuo*, and purified by flash chromatography (1.5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to afford the desired product (7.3 mg, 54%): MS APCI (+) *m/z* 412, 414 (M<sup>+</sup>, Br pattern) detected; <sup>1</sup>H NMR (400 MHz, acetone-*d*<sub>6</sub>) δ 8.64 (s, 1H), 8.34 (s, 1H), 8.16 (s, 1H), 7.58 (d, 1H), 7.31 (dd, 1H), 6.59 (dd, 1H), 4.94 (s, 2H), 4.06 (s, 3H); <sup>19</sup>F NMR (376 MHz, acetone-*d*<sub>6</sub>) -132.45 (s, 1F).

#### Example 43



*1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-hydroxy-ethanone (10k)*

*Step A: 1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methoxymethoxy-ethanol 10l*

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzimidazole-5-carbaldehyde **10f** is treated with tributyl-methoxymethoxymethyl-stannane according to the procedure


described in Example 42, Step A to yield compound **10l**. MS APCI (+)  $m/z$  444, 446 ( $M^+$ , Br pattern) detected.

*Step B: 1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methoxymethoxy-ethanone 10m*

To a solution of oxalyl chloride (0.11 mL, 0.22 mmol) in  $CH_2Cl_2$  (1 mL) at  $-78^\circ C$  is added DMSO (0.016 mL, 0.22 mmol). After stirring for 3 min, a solution of 1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methoxymethoxy-ethanol **10l** (25 mg, 0.056 mmol) in methylene chloride (1 mL) is added. The resulting solution is stirred for 30 min at  $-78^\circ C$ . TEA (0.1 mL, 0.71 mmol) is added. The reaction mixture is slowly warmed to room temperature, stirred for 5 min at room temperature, and diluted with water and  $CH_2Cl_2$ . The organic layer is separated, dried over  $MgSO_4$ , filtered, and concentrated to give the crude product which is directly used without further purification.

*Step C: 1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-hydroxy-ethanone 10k*

1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methoxymethoxy-ethanone **10m** is deprotected according to the procedure described in Example 42, Step C to yield compound **10k**. MS APCI (+)  $m/z$  398, 400 ( $M^+$ , Br pattern) detected;  $^1H$  NMR (400 MHz,  $CD_3OD$ )  $\delta$  8.38 (s, 1H), 8.04 (s, 1H), 7.52 (d, 1H), 7.22 (dd, 1H), 6.53 (dd, 1H), 4.90 (m, 2H);  $^{19}F$  NMR (376 MHz,  $CD_3OD$ )  $-133.96$  (s, 1F).

**Example 44**

***1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-ethoxy-ethanone (10n)***

**Step A:** *1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-ethoxy-ethanol 10o*


To a solution of lithiomethyl ethyl ether in THF (6 mL) (prepared from 4,4'-di-tert-butylbiphenyl (585 mg, 2.20 mmol), Li (18 mg, 2.59 mmol), and EtOCH<sub>2</sub>Cl (0.20 mL, 2.05 mmol) by the procedure reported in *Tetrahedron* 1996, 52, 1643) is added a solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carbaldehyde **10f** (29 mg, 0.080 mmol) in THF (1 mL) at -78 °C. The resulting solution is stirred for 1 h and then quenched with saturated aqueous NH<sub>4</sub>Cl at -78 °C, warmed to room temperature, and extracted with EtOAc. The organic layer is washed with brine, dried over MgSO<sub>4</sub>, filtered, concentrated in vacuo, and purified by flash chromatography (100% CH<sub>2</sub>Cl<sub>2</sub> to 3% to 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to give the desired product (15 mg, 44%): MS APCI (+) *m/z* 428, 430 (M<sup>+</sup>, Br pattern) detected.

**Step B:** *1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-ethoxy-ethanone 10n*

The title compound is prepared from 1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-ethoxy-ethanol **10o** according to the procedure described in Example 42, Step B except that the reaction mixture is not treated with saturated aqueous NaHCO<sub>3</sub> containing sodium thiosulfate pentahydrate. MS APCI (+)

$m/z$  426, 428 ( $M^+$ , Br pattern) detected;  $^1H$  NMR (400 MHz,  $CD_3OD$ )  $\delta$  8.36 (s, 1H), 8.04 (s, 1H), 7.51 (d, 1H), 7.21 (dd, 1H), 6.51 (dd, 1H), 4.76 (s, 2H), 3.57 (q, 2H), 1.19 (t, 3H);  $^{19}F$  NMR (376 MHz,  $CD_3OD$ )  $-133.96$  (s).


#### Example 45



***1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methoxy-ethanone (10p)***

1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methoxy-ethanone **10p** is prepared from 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carbaldehyde **10f** and lithiummethyl methyl ether by the procedures described in Example 44. MS APCI (+)  $m/z$  412, 414 ( $M^+$ , Br pattern) detected.

#### Example 46



***2-Benzyloxy-1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-ethanone (10q)***

**Step A:** 2-Benzyloxy-1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-ethanol **10r**




To a solution of benzyloxymethyl lithium in THF (2 mL, prepared from n-Bu<sub>3</sub>SnCH<sub>2</sub>OBn (505 mg, 1.23 mmol) and n-BuLi (0.49 mL, 1.22 mmol, 2.5 M solution in hexane) by the procedure reported in *J. Am. Chem. Soc.* 1978, 100, 1481) is added a solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3*H*-benzoimidazole-5-carbaldehyde 10f (51 mg, 0.14 mmol) in THF (3 mL) at -78 °C. The resulting solution is stirred for 1 h at -78 °C. The reaction is quenched with saturated aqueous NH<sub>4</sub>Cl, and extracted with EtOAc. The organic layer is dried over MgSO<sub>4</sub>, filtered, concentrated *in vacuo*, and purified by flash chromatography (100% CH<sub>2</sub>Cl<sub>2</sub> to 3% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to afford the desired product (46 mg, 68%): MS APCI (+) *m/z* 490, 492 (M<sup>+</sup>, Br pattern) detected.

*Step B: 2-Benzyloxy-1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-ethanone 10q*

The title compound is prepared from 2-benzyloxy-1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3*H*-benzoimidazol-5-yl]-ethanol 10r by the procedure described in Example 42, Step B except that the reaction mixture is not treated with saturated aqueous NaHCO<sub>3</sub> containing sodium thiosulfate pentahydrate: MS APCI (+) *m/z* 488, 490 (M<sup>+</sup>, Br pattern) detected; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 8.37 (s, 1H), 8.02 (s, 1H), 7.51 (d, 1H), 7.26 (m, 5H), 7.19 (dd, 1H), 6.46 (dd, 1H), 4.77 (s, 2H), 4.58 (s, 2H); <sup>19</sup>F NMR (376 MHz, CD<sub>3</sub>OD) -134.52 (s).

## Example 47



***1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methanesulfonyl-ethanone (10s)***

**Step A:** *1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methanesulfonyl-ethanol 10t*


To a solution of methyl sulfone (65 mg, 0.68 mmol) in THF (1.5 mL) is added a solution of n-BuLi (0.27 mL, 0.68 mmol, 2.5 M solution in hexane) at -78 °C. After stirring for 5 min, HMPA (0.1 mL) is added. After stirring for additional 10 min, a solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carbaldehyde **10f** (26 mg, 0.069 mmol) in THF (1 mL) is added. The resulting solution is stirred for 1.5 h at -78 °C. The reaction is quenched with saturated aqueous NH<sub>4</sub>Cl, warmed to room temperature, and diluted with EtOAc. The organic layer is washed with water, dried over MgSO<sub>4</sub>, filtered, concentrated *in vacuo*, and purified by flash chromatography (3% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to give the crude desired product (31 mg, 96%) which is used directly without further purification: MS APCI (+) *m/z* 462, 464 (M<sup>+</sup>, Br pattern) detected.

**Step B:** *1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methanesulfonyl-ethanone 10s*

The title compound is prepared from 1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methanesulfonyl-ethanol **10t** by the procedure described in Example 42, Step B except that the reaction mixture is not treated with

saturated aqueous NaHCO<sub>3</sub> containing sodium thiosulfate pentahydrate: MS APCI (+) *m/z* 460, 462 (M<sup>+</sup>, Br pattern) detected; <sup>1</sup>H NMR (400 MHz, acetone-*d*<sub>6</sub>) δ 8.44 (s, 1H), 8.33 (s, 1H), 7.59 (s, 1H), 7.32 (d, 1H), 6.68 (dd, 1H), 5.00 (s, 1H), 3.15 (s, 3H); <sup>19</sup>F NMR (376 MHz, acetone-*d*<sub>6</sub>) -132.97 (s).

### Example 48



#### *1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-ethane-1,2-diol (10u)*


*Step A: 1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-(isopropoxy-dimethyl-silanyl)-ethanol 10v*

To a solution of the Grignard reagent prepared from Mg and chloromethyl dimethylisopropoxy silane (*Org. Synth.* **1992**, 69, 96 ) [4.4 mL, 3.26 mmol, 0.74 M solution (based on 90% purity)] in THF, is added a solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carbaldehyde **10f** (200 mg, 0.54 mmol) in THF (1 mL) at -78 °C. After stirring for 1 h at -78 °C, the reaction is quenched with saturated aqueous NH<sub>4</sub>Cl, and extracted with EtOAc. The organic layer is dried over MgSO<sub>4</sub>, filtered, concentrated *in vacuo* to afford the crude desired product which is directly used without further purification.

*Step B: 1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-ethane-1,2-diol 10u*

To the crude 1-[6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3H-benzimidazol-5-yl]-2-(isopropoxy-dimethyl-silanyl)-ethanol **10v** in MeOH-THF (5 mL-5 mL) is added  $\text{KHCO}_3$  (54 mg, 0.54 mmol), and KF (74 mg, 1.27 mmol), and 30% aqueous  $\text{H}_2\text{O}_2$  (0.20 mL) at rt. After stirring for 3.5 h at room temperature, the reaction mixture is diluted with water, and extracted with EtOAc. The organic layer is dried over  $\text{MgSO}_4$ , filtered, concentrated *in vacuo*, and purified by flash chromatography (8% to 10 % MeOH in  $\text{CH}_2\text{Cl}_2$ ) to give the desired product (74 mg, 34%): MS APCI (+)  $m/z$  400, 402 ( $\text{M}^+$ , Br pattern) detected;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.20 (s, 1H), 7.62 (broad s, 1H), 7.47 (d, 1H), 7.14 (dd, 1H), 6.30 (d, 1H), 4.96 (t, 1H), 3.64 (m, 2H);  $^{19}\text{F}$  NMR (376 MHz,  $\text{CD}_3\text{OD}$ ) -136.87 (s).

#### Example 49




#### ***[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzimidazol-5-yl]-pyridin-2-yl-methanol (10w)***

To a solution of 2-bromopyridine (0.10 mL, 1.04 mmol) in THF (3 mL) at -78 °C is added *n*-BuLi (0.39 mL, 0.98 mmol, 2.5 M solution in hexane). After stirring for 10 min at -78 °C, a solution of 6-(4-bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzimidazole-5-carbaldehyde **10h** (25 mg, 0.064 mmol) in THF (1 mL) is added. The resulting reaction mixture is stirred for 1.5 h at -78 °C, quenched with saturated aqueous  $\text{NH}_4\text{Cl}$ , and extracted with EtOAc. The organic layer is dried over  $\text{MgSO}_4$ , filtered, concentrated *in vacuo*, and purified by flash chromatography (2.5% MeOH in  $\text{CH}_2\text{Cl}_2$ ) to

afford the desired product (18 mg, 62%): MS APCI (+)  $m/z$  461, 463 ( $M^+$ , Br pattern) detected;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.31 (d, 1H), 8.16 (s, 1H), 7.65 (m, 3H), 7.38 (d, 1H), 7.10 (m, 1H), 7.00 (dd, 1H), 6.11 (dd, 1H), 6.05 (s, 1H), 3.94 (s, 3H);  $^{19}\text{F}$  NMR (376 MHz,  $\text{CD}_3\text{OD}$ )  $-135.79$  (s).

### Example 50



**(4-Bromo-2-chloro-phenyl)-(4-fluoro-6-oxazol-5-yl-1H-benzimidazol-5-yl)-amine**  
(10x)

*Step A:* [6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzimidazol-5-yl]-methanol 10y

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzimidazole-5-carboxylic acid methyl ester 11cc (0.300 g, 0.594 mmol) is suspended in a mixture of EtOH (6 ml) and THF (4 ml) under  $\text{N}_2$ .  $\text{NaBH}_4$  (0.112 g, 2.97 mmol) is added. After approximately 4 days stirring, reaction mixture is quenched by the addition of AcOH until the reaction mixture reaches pH 7. The reaction mixture is concentrated to dryness under reduced pressure and the residue partitioned between ethyl acetate and water. The layers are separated and the organic layer is washed with water (3x), brine, and dried ( $\text{Na}_2\text{SO}_4$ ). The organic layer is concentrated under reduced pressure until a white precipitate forms which is collected by filtration to give 0.225 g (79%) clean desired product: LC/MS ESI (+)  $m/z$  478, 476 ( $M^+$  Br pattern) detected.

*Step B: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzoimidazole-5-carbaldehyde 10z*

[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzoimidazol-5-yl]-methanol **10y** (0.050, 0.105 mmol) is dissolved in 1:1 THF:acetone (2 ml) under N<sub>2</sub> and MnO<sub>2</sub> (0.046 g, 0.524 mmol) is added. The reaction mixture is stirred at room temperature for 16 h, and then at 55 °C for 5 h. Additional MnO<sub>2</sub> (0.046 g, 0.524 mmol) is added and the reaction mixture stirred at 55 °C for 2 h. The reaction mixture is concentrated to dryness and the residue dissolved in 10:1 methylene chloride:MeOH. The solution is filtered through a silica gel plug eluted with 10:1 methylene chloride:MeOH. The resulting filtrate is concentrated under reduced pressure to give 41 mg (82%) desired product as a bright yellow solid.

*Step C: (4-Bromo-2-chloro-phenyl)-(4-fluoro-6-oxazol-5-yl-1H-benzoimidazol-5-yl)-amine 10x*


6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzoimidazole-5-carbaldehyde **10z** (0.025 g, 0.053 mmol) is suspended in MeOH (2 ml) and K<sub>2</sub>CO<sub>3</sub> (0.015 g, 0.105 mmol) is added followed by tosylmethyl isocyanide (0.011 g, 0.058 mmol). The reaction mixture is heated to reflux under N<sub>2</sub> for 16 h. After cooling, additional tosylmethyl isocyanide (0.011 g, 0.058 mmol) is added and the reaction mixture heated to reflux under N<sub>2</sub> for 16 h. The reaction mixture is cooled to room temperature, concentrated under reduced pressure and dissolved in ethyl acetate. The organic solution is washed with water and brine. The combined aqueous washes are extracted with ethyl acetate. The combined organic extracts are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. Purification by flash column chromatography eluted with 20:1 methylene chloride:MeOH gives 4 mg (18%) desired product **10x** and 1

mg (4%) (4-bromo-2-chloro-phenyl)-[4-fluoro-1-(2-methanesulfonyl-ethyl)-6-oxazol-5-yl]-1H-benzoimidazol-5-yl]-amine.

(4-Bromo-2-chloro-phenyl)-(4-fluoro-6-oxazol-5-yl-1H-benzoimidazol-5-yl)-amine **10x**. LC/MS ESI (+)  $m/z$  409, 407 ( $M+Br$  pattern) detected;  $^1H$  NMR (400MHz, MeOH- $d_4$ )  $\delta$  8.33 (s, 1H), 8.24 (s, 1H), 7.94 (bs, 1H), 7.51 (d, 1H), 7.33 (s, 1H), 7.07 (dd, 1H), 6.14 (dd, 1H).

(4-Bromo-2-chloro-phenyl)-[4-fluoro-1-(2-methanesulfonyl-ethyl)-6-oxazol-5-yl]-1H-benzoimidazol-5-yl]-amine. LC/MS ESI (+)  $m/z$  515, 513 ( $M+Br$  pattern) detected;  $^1H$  NMR (400MHz, MeOH- $d_4$ )  $\delta$  8.39 (s, 1H), 8.28 (s, 1H), 8.03 (s, 1H), 7.52 (d, 1H), 7.37 (s, 1H), 7.07 (m, 1H), 6.14 (dd, 1H), 3.83 (t, 2H), 2.99 (s, 3H), 1.18 (t, 2H).

### Example 51



#### *(4-Bromo-2-chloro-phenyl)-[4-fluoro-6-(3H-imidazol-4-yl)-1H-benzoimidazol-5-yl]-amine (10aa)*

*Step A: (4-Bromo-2-chloro-phenyl)-{4-fluoro-1-(2-methanesulfonyl-ethyl)-6-[4-(toluene-4-sulfonyl)-4,5-dihydro-oxazol-5-yl]-1H-benzoimidazol-5-yl}-amine 10bb*


6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(2-methanesulfonyl-ethyl)-3H-benzoimidazole-5-carbaldehyde **10z** (0.050 g, 0.107 mmol) is suspended in EtOH (0.5 ml) under  $N_2$  and tosylmethyl isocyanide (0.020 g, 0.105 mmol) is added followed by catalytic NaCN (~1 mg). After 2 h, 2 ml THF is added to assist with solubility. After stirring for 16 h at room temperature, a second equivalent of tosylmethyl isocyanide

(0.020 g, 0.105 mmol) is added. After 8 h, the reaction mixture is concentrated under reduced pressure and used as is in the next reaction: LC/MS ESI (+)  $m/z$  671, 669 ( $M+ Br$  pattern) detected.

*Step B: (4-Bromo-2-chloro-phenyl)-[4-fluoro-6-(3H-imidazol-4-yl)-1H-benzoimidazol-5-yl]-amine 10aa*

(4-Bromo-2-chloro-phenyl)-{4-fluoro-1-(2-methanesulfonyl-ethyl)-6-[4-(toluene-4-sulfonyl)-4,5-dihydro-oxazol-5-yl]-1H-benzoimidazol-5-yl}-amine **10bb** (0.072 g, 0.107 mmol) is treated with 2.4 ml of a 2.0 M  $NH_3$  in MeOH solution in a sealed pressure tube. The reaction mixture is then heated to 90 °C with stirring for 20 h and further stirred at room temperature for 3 days. The reaction mixture is transferred to a round bottom flask and concentrated under reduced pressure. Purification by flash column chromatography, twice, eluted with 10:1 methylene chloride:MeOH, followed by successive tritutions with methylene chloride and then diethyl ether gives 3 mg (7%) desired product: LC/MS ESI (+)  $m/z$  408, 406 ( $M+ Br$  pattern) detected;  $^1H$  NMR (400MHz, MeOH- $d_4$ )  $\delta$  8.23 (s, 1H), 7.87 (s, 1H), 7.74 (s, 1H), 7.46 (m, 1H), 7.32 (d, 1H), 7.05 (m, 1H), 6.20 (dd, 1H).

### Example 52



**6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3-methyl-3H-benzimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide (10cc)**



*Step A: 3-Chloro-2,4-difluoro-5-nitro-benzoic acid 2a*

3-Chloro-2,4-difluoro-benzoic acid **1a** (3.00 g, 15.6 mmol) is added to a stirred solution of concentrated H<sub>2</sub>SO<sub>4</sub> (16 mL) and fuming nitric acid (0.85 mL, 20.3 mmol). After 3 hours a precipitate forms. The yellow slurry is poured onto ice water (100 mL). The aqueous mixture is extracted with diethyl ether (3x). The organic extracts are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give 3.50 g (95%) of clean desired product as a pale yellow solid.

*Step B: 4-Amino-3-chloro-2-fluoro-5-nitro-benzoic acid 3a*

Ammonium hydroxide solution (6.88 g, ~30% in water, 58.9 mmol) is added to a solution of 3-chloro-2,4-difluoro-5-nitro-benzoic acid **2a** (3.5 g, 14.7 mmol) in water (16 mL) at 0 °C with stirring. Upon completion of the ammonium hydroxide addition the reaction mixture is warmed to room temperature. After 5 hours the reaction mixture is cooled to 0 °C and concentrated HCl is carefully added until the pH of the reaction mixture is near zero. The solid is collected by filtration and washed with water and diethyl ether. The solids are transferred to a round bottom flask as a solution in MeOH and EtOAc and concentrated under reduced pressure to give 2.96 g of a yellow solid. The filtrate is partitioned between diethyl ether and water and the organic layer is washed with brine. The combined organic extracts are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give 0.65 g of product. Recovered a total of 3.61 g (104%) of pure desired product, that is carried forward without further purification.

*Step C: 4-Amino-3-chloro-2-fluoro-5-nitro-benzoic acid methyl ester 4a*

To a stirred solution of 4-amino-3-chloro-2-fluoro-5-nitro-benzoic acid **3a** (3.61 g, 15.4 mmol) in THF (30 mL) and MeOH (10 mL), TMS diazomethane (9.23 mL, 2.0 M

solution in hexanes, 18.5 mmol) is added. After completion of reaction, the reaction mixture is concentrated via rotary evaporation with acetic acid in the trap. The recovered oily solid is triturated with diethyl ether to provide 1.51 g of a yellow solid. The filtrate is concentrated and triturated with diethyl ether to give an additional 0.69 g of yellow solid. A total of 2.20 g (57%) of pure desired product is recovered.

*Step D: 4-Amino-3-chloro-5-nitro-2-phenylamino-benzoic acid methyl ester 5c*

4-Amino-3-chloro-2-fluoro-5-nitro-benzoic acid methyl ester **4a** (2.20 g, 8.84 mmol) is suspended in MeOH (9.4 mL) and aniline (3.22 mL, 35.4 mmol) is added. The reaction mixture is heated to reflux with stirring under a nitrogen atmosphere. After 19 hours, the reaction is complete. Distilled water (3.22 mL) is added to the reaction mixture and refluxing is continued for one hour. The reaction mixture is cooled to 0 °C in an ice bath for 20 minutes. The reaction mixture is filtered and washed with 3:10 distilled water/MeOH (65 mL total) and then with MeOH. The solid is dissolved with CH<sub>2</sub>Cl<sub>2</sub> and concentrated under reduced pressure to give 2.40 g (84%) of pure desired product. MS APCI (-) *m/z* 320.3 (M-1) detected.

*Step E: 4,5-Diamino-3-chloro-2-phenylamino-benzoic acid methyl ester 6b*

4-Amino-3-chloro-5-nitro-2-phenylamino-benzoic acid methyl ester **5c** (0.50 g, 1.55 mmol) is dissolved into 2:1 EtOH/MeOH (15.5 mL). Saturated aqueous NH<sub>4</sub>Cl (15 mL), Zn powder (1.02 g, 15.6 mmol), and THF (10 mL) are added. After stirring for 20 hours, the reaction mixture is diluted with CH<sub>2</sub>Cl<sub>2</sub>/THF and water. The organic layer is washed with water (3x). The combined organic extracts are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The solids are triturated with ether to give 0.32 g (70%) clean desired product.

*Step F: 7-Chloro-6-phenylamino-3H-benzoimidazole-5-carboxylic acid methyl ester 7c*

4,5-Diamino-3-chloro-2-phenylamino-benzoic acid methyl ester **6b** (0.32 g, 1.09 mmol) and formamidine acetate (72 mg, 1.64 mmol) in EtOH (36 mL) are heated, with stirring, to 80 °C. After 44 hours, the reaction mixture is cooled to room temperature and diluted with EtOAc and washed with water (3x), saturated NaHCO<sub>3</sub>, and brine. The combined organic extracts are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give 0.33 g (99%) clean desired product as a solid. MS APCI (+) *m/z* 302.3 (M+1) detected.

*Step G: 6-(4-Bromo-phenylamino)-7-chloro-3H-benzoimidazole-5-carboxylic acid methyl ester 8g*

7-Chloro-6-phenylamino-3H-benzoimidazole-5-carboxylic acid methyl ester **7c** (0.327 g, 1.08 mmol) is dissolved into DMF (16 mL) and NBS (0.193 g, 1.08 mmol) is added. After one hour, the reaction mixture is quenched by the addition of saturated aqueous NaHSO<sub>3</sub>. The reaction mixture is then partitioned between EtOAc/THF and water. The organic layer is washed with water and brine. The combined organic extracts are dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The recovered solid is triturated with ether to give 0.225 g (54%) pure desired product. MS ESI (+) *m/z* 382, 384 (M<sup>+</sup>, Br pattern) detected.

*Step H: 6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3H-benzoimidazole-5-carboxylic acid methyl ester 10dd*

6-(4-Bromo-phenylamino)-7-chloro-3H-benzimidazole-5-carboxylic acid methyl ester **8g** (0.225 g, 0.591 mmol) is dissolved in DMF (2 mL) and NCS (79 mg, 0.591 mmol) is added. After the NCS is in solution concentrated HCl (0.005 mL, 0.059 mmol) is added. After 2 hours, sodium bicarbonate, water and NaHSO<sub>3</sub> are added to the reaction mixture. Solids are filtered and washed with water and ether to give 0.141 g (57%) of clean desired product as a tan solid. MS APCI (-) *m/z* 414, 416 (M-, Br pattern) detected.

*Step I: 6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3-methyl-3H-benzimidazole-5-carboxylic acid methyl ester 10ee*

6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3H-benzimidazole-5-carboxylic acid methyl ester **10dd** (0.141 g, 0.34 mmol), potassium carbonate (0.141 g, 1.02 mmol), and iodomethane (0.063 mL, 1.02 mmol) are dissolved in dimethylformamide (3 mL). After 20 hours, the reaction mixture is diluted with EtOAc and washed with water (3x), potassium carbonate, and brine. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to a brown oil. The N3 and N1 alkylated regioisomers are separated by flash chromatography (EtOAc). The recovery of the N3 alkylated regioisomer is 20.4 mg (28%). MS ESI (+) *m/z* 428, 430 (M+, Br pattern) detected.

*Step J: 6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3-methyl-3H-benzimidazole-5-carboxylic acid 10ff*

6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3-methyl-3H-benzimidazole-5-carboxylic acid methyl ester **10ee** (21 mg, 0.048 mmol) is dissolved into 2:1 THF/water (1.2 mL) and NaOH (0.190 mL, 1.0 M aqueous solution, 0.190 mmol) is added. After stirring for 4 hours the reaction is diluted with water and acidified to pH 2 by addition of 1.0 M HCl. The mixture is then extracted with 3:1 EtOAc/THF (3x), dried (Na<sub>2</sub>SO<sub>4</sub>) and

concentrated to give quantitative yield of desired product as a white solid. MS APCI (+)  $m/z$  414, 416 (M<sup>+</sup>, Br pattern) detected.

*Step K: 6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3-methyl-3H-benzimidazole-5-carboxylic acid (2-vinyloxy-ethoxy)-amide 10gg*


6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3-methyl-3H-benzimidazole-5-carboxylic acid **10ff** (32 mg, 0.077 mmol), O-(2-vinyloxy-ethyl)-hydroxylamine (0.010 mL, 0.092 mmol), HOBt (13 mg, 0.093 mmol), triethylamine (0.011 mL, 0.077 mmol), and EDCI (19 mg, 0.10 mmol) are dissolved into dimethylformamide (1.0 mL) and allowed to stir under a nitrogen atmosphere at room temperature for 24 hours. The reaction mixture is diluted with EtOAc, washed with water (3x), 10% potassium carbonate (2x), saturated ammonium chloride, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated under reduced pressure to give 39 mg of 85% pure material. MS APCI (-)  $m/z$  497, 501 (M<sup>-</sup>, Br pattern) detected.

*Step L: 6-(4-Bromo-2-chloro-phenylamino)-7-chloro-3-methyl-3H-benzimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide 10cc*

Hydrochloric acid (0.78 mL, 1.0 M aqueous solution, 0.78 mmol) is added to a suspension of 6-(4-bromo-2-chloro-phenylamino)-7-chloro-3-methyl-3H-benzimidazole-5-carboxylic acid **10gg** (2-vinyloxy-ethoxy)-amide (39 mg, 0.078 mmol) in MeOH (1 mL). After one hour, the reaction mixture is neutralized to pH 7 and concentrated under reduced pressure. The solids are dissolved in EtOAc, washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated under reduced pressure. Flash chromatography (20:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH) provides 9 mg (23%) of pure product: MS APCI (+)  $m/z$  473, 475

(M<sup>+</sup>, Br pattern) detected; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.30 (s, 1H), 8.08 (s, 1H), 7.57 (d, 1H), 7.15 (dd, 1H), 6.21 (d, 1H), 3.97 (s, 3H) 3.86 (m, 2H), 3.57 (m, 2H).


### Example 53



***6-(4-Bromo-2-chloro-phenylamino)-3H-benzoimidazole-5-carboxylic acid (2-hydroxyethoxy)-amide (10hh)***

The above compound is prepared in an analogous fashion to Example 52 except that Step I is eliminated. MS APCI (-) *m/z* 457, 461 (M<sup>-</sup>, Br pattern) detected; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 8.40 (s, 1H), 7.85 (s, 1H), 7.50 (d, 1H), 7.14 (dd, 1H), 6.21 (d, 1H), 3.84 (m, 2H), 3.61 (m, 2H).

### Example 54



***6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-2-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxyethoxy)-amide (10ii)***

*Step A: 4,5-Diamino-3-fluoro-2-phenylamino-benzoic acid methyl ester 6c*

4-Amino-3-fluoro-5-nitro-2-phenylamino-benzoic acid methyl ester 26a (11.44 g, 37.48 mmol) is suspended in ethanol (400 mL) and ammonium formate (11.80 g, 187.0

mmol) and 20 % Pd(OH)<sub>2</sub>/C (10.00 g, 18.79 mmol) are added. The reaction mixture is stirred at 95 °C under N<sub>2</sub> for 30 minutes. The reaction mixture is cooled to room temperature and then filtered through celite, rinsing with ethanol. The filtrate is concentrated under reduced pressure to give 9.63 g (93 %) of the pure desired product as a purple/red solid. MS ESI (+) *m/z* 276 (M+1) detected.

*Step B: 7-Fluoro-2-methyl-6-phenylamino-3H-benzoimidazole-5-carboxylic acid methyl ester 31a*


4,5-Diamino-3-fluoro-2-phenylamino-benzoic acid methyl ester 6c (0.20 g, 0.73 mmol) is suspended in ethanol (3 mL) and 5 M aqueous HCl (1 mL, 5.00 mmol) is added. The reaction mixture is brought to reflux under N<sub>2</sub> and then 2,4-pentanedione (0.150 mL, 1.45 mmol) is added. The reaction mixture is stirred at reflux for 60 minutes. The reaction mixture is cooled to room temperature and treated with saturated aqueous NaHCO<sub>3</sub> until the pH of the reaction mixture is pH 7 and is then concentrated under reduced pressure to dryness. The residue is diluted with ethyl acetate and water, poured into a separatory funnel and the layers separated. The ethyl acetate layer is washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The red solid residue is triturated with diethyl ether to yield a light brown solid and a red filtrate. The solid is collected and washed with diethyl ether to yield 0.20 g (91 %) of the pure desired product as a light brown solid. MS ESI (+) *m/z* 300 (M+1) detected.

*Step C: 6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-2-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide 10ii*

7-Fluoro-2-methyl-6-phenylamino-3H-benzoimidazole-5-carboxylic acid methyl ester 31a is converted by the bromination, chlorination, hydrolysis, coupling, and

hydrolysis procedures already described to yield the pure desired product as an off-white solid. MS ESI (+)  $m/z$  457, 459 ( $M^+$ , Br pattern) detected;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.58 (s, 1H), 7.49 (d, 1H), 7.18 (dd, 1H), 6.41 (m, 1H), 3.91 (t, 2H), 3.65 (t, 2H), 2.61 (s, 3H);  $^{19}\text{F}$  NMR (376 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  -135.84 (s).

### Example 55



**6-(4-Cyano-2-methyl-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid  
cyclopropylmethoxy-amide (11yy)**

*Step A: 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzimidazole-5-carboxylic acid methyl ester 10jj*

7-Fluoro-6-*o*-tolylamino-1H-benzimidazole-5-carboxylic acid methyl ester **7a** (1.47 g, 4.92 mmol) is suspended in 1:1 THF:MeOH mixture (44 ml) and cooled to  $-78^\circ\text{C}$  under a nitrogen atmosphere. A solution of NIS (1.66 g, 7.39 mmol) in THF (2 ml) is added followed by a MeOH (2 ml) solution of TsOH $\cdot$ H $_2$ O (1.87 g, 9.84 mmol). After 30 minutes, the reaction mixture is warmed to  $0^\circ\text{C}$  and 1 ml methylene chloride is added. The reaction is slowly allowed to warm to room temperature with stirring over 16 hours. The reaction mixture is quenched by the addition of 10%  $\text{Na}_2\text{S}_2\text{O}_4$  solution. The reaction mixture is then diluted with water and ethyl acetate and the layers separated. The aqueous layer is extracted with ethyl acetate. The combined organic extracts are dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The recovered solid is triturated with



MeOH to give 1.45 g (69%) pure desired product: MS ESI (+)  $m/z$  426 (M+1) detected; MS ESI (-)  $m/z$  424 (M-1) detected.

*Step B: 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-(2-trimethylsilyl-ethoxymethyl)-benzoimidazole-5-carboxylic acid methyl ester 10kk*

7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid methyl ester **10jj** (0.200 g, 0.470 mmol) is suspended in DMF (2 ml) under N<sub>2</sub> and cooled to 0 °C in an ice-water bath. NaH (60% dispersion in oil, 0.018 g, 0.470 mmol) is added. After 10 min, the reaction mixture is warmed to room temperature and stirred for 30 min. After cooling to 0 °C, SEMCl (0.087 ml, 0.494 mmol) is added and the reaction is allowed to warm to room temperature with stirring overnight. The reaction mixture is quenched by the addition of water and brine. The reaction mixture is extracted with ethyl acetate. The combined organic extracts are washed with water and brine, and dried (MgSO<sub>4</sub>) and concentrated under reduced pressure. Purification by flash column chromatography eluted with 1:1 hexanes:ethyl acetate gives 0.182 g (70%) of desired product as a 1:1 mixture of N1 and N3 isomers as a white foam.

*Step C: 6-(4-Cyano-2-methyl-phenylamino)-7-fluoro-(2-trimethylsilyl-ethoxymethyl)-benzoimidazole-5-carboxylic acid methyl ester 10ll*

To a stirred solution of a 1:1 mixture of N1:N3 isomers of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-(2-trimethylsilyl-ethoxymethyl)-benzoimidazole-5-carboxylic acid methyl ester **10jj** (0.060 g, 0.108 mmol) in 1 ml DMF at room temperature under N<sub>2</sub> is added dppf (2 mg, 0.004 mmol) followed by Pd<sub>2</sub>dba<sub>3</sub> (2 mg, 0.002 mmol) and Zn(CN)<sub>2</sub> (8 mg, 0.065 mmol) (Tetrahedron Lett. 1999, 40, 8193-8195). The reaction mixture is heated to 120 °C for 45 min. The reaction mixture is cooled to room temperature and

quenched by the addition of 5 ml of a 4:1:5 mixture of sat  $\text{NH}_4\text{Cl}$ :conc  $\text{NH}_4\text{OH}$ :water. The aqueous layer is extracted with ethyl acetate. The combined organic extracts are washed with water (3x), brine, and dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure. Purification by flash column chromatography eluted with 1:1 hexanes:ethyl acetate gives 38 mg (77%) of desired product as a 1:1 mixture of N1 and N3 isomers: APCI MS (+)  $m/z$  455 (M+1) detected.

*Step D: 6-(4-Cyano-2-methyl-phenylamino)-7-fluoro-(2-trimethylsilyl-ethoxymethyl)-benzimidazole-5-carboxylic acid 10mm*

A 1:1 mixture of N1:N3 isomers of 6-(4-cyano-2-methyl-phenylamino)-7-fluoro-(2-trimethylsilyl-ethoxymethyl)-benzimidazole-5-carboxylic acid methyl ester **10II** (31 mg, 0.068 mmol) is hydrolyzed with aqueous sodium hydroxide as described previously to give 26 mg (87%) of desired product.


*Step E: 6-(4-Cyano-2-methyl-phenylamino)-7-fluoro-(2-trimethylsilyl-ethoxymethyl)-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide 11zz*

A 1:1 mixture of N1:N3 isomers of 6-(4-cyano-2-methyl-phenylamino)-7-fluoro-(2-trimethylsilyl-ethoxymethyl)-benzimidazole-5-carboxylic acid **10mm** (26 mg, 0.059 mmol) is coupled with EDCI and cyclopropyl methyl hydroxylamine hydrochloride as described previously to give 28 mg (93%) of desired product: APCI MS (+)  $m/z$  510 (M+1) detected.

*Step F: 6-(4-Cyano-2-methyl-phenylamino)-7-fluoro-3H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide 11yy*

To a slurry of a 1:1 mixture of N1:N3 isomers 6-(4-cyano-2-methyl-phenylamino)-7-fluoro-(2-trimethylsilyl-ethoxymethyl)-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide **11zz** (28 mg, 0.055 mmol) in 0.5 ml EtOH is added 0.5 ml 10% HCl. The reaction mixture is heated to 50 °C with stirring overnight (Whitten *et al.*, *JOC* 1986, *51*, 1891-1894). An additional 0.5 ml 10% HCl is added and the reaction mixture stirred at 70 °C overnight. The reaction mixture is cooled to room temperature and neutralized to pH ~8 with 1.5 ml 1N NaOH. The reaction mixture is extracted with ethyl acetate, dried (MgSO<sub>4</sub>) and concentrated under reduced pressure to give 14 mg (60%) of 90% pure product as a mixture of rotamers: MS APCI (+) *m/z* 380 (M+1) detected; MS APCI (-) *m/z* 378 (M-1) detected; <sup>1</sup>H NMR (400MHz, MeOH-d<sub>4</sub>) δ 8.41 (bs, 1H), 7.75 (m, 1H), 7.50 (s, 1H), 7.38 (d, 1H), 6.51 (m, 1H), 3.72 (d, 0.5 H), 3.65 (d, 1.5 H), 2.41 (s, 3H), 0.98 (1H, m), 0.58 (d, 1.5 H), 0.40 (d, 0.5 H), 0.25 (d, 1.5 H), 0.19 (d, 0.5 H).

### Example 56



**6-(4-Ethynyl-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid  
cyclopropylmethoxy-amide **11aaa****

**Step A.** 7-Fluoro-6-(2-methyl-4-trimethylsilyl-ethynyl-phenylamino)-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide **11bbb**

7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide **11ccc** (0.025 g, 0.052 mmol) is dissolved in 1:1 acetonitrile/triethylamine (0.50 mL). Ethynyl-trimethylsilane (0.013 mL, 0.092 mmol),

$\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$  (0.004 g, 0.006 mmol), and  $\text{CuI}$  (0.002 g, 0.011 mmol) are added consecutively and the reaction mixture is stirred at 60 °C for 1 hour under  $\text{N}_2$ . The reaction mixture is cooled to room temperature and concentrated under reduced pressure. The residue is purified by FCC (eluting with 20:1 methylene chloride:methanol) to yield 0.020 g (87%) of the desired product.

*Step B: 6-(4-Ethynyl-2-methyl-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide 11aaa*


7-Fluoro-6-(2-methyl-4-trimethylsilanylethynyl-phenylamino)-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide **11bbb** (0.020 g, 0.044 mmol) is dissolved in tetrahydrofuran (0.50 mL) and the reaction solution is cooled to 0 °C. TBAF (50  $\mu\text{L}$ , 0.050 mmol, 1 M solution in tetrahydrofuran) is added. The reaction mixture is warmed to room temperature and additional TBAF (25  $\mu\text{L}$ , 0.025 mmol, 1 M solution in tetrahydrofuran) is added. The reaction mixture is stirred at 50 °C for 2 hours under  $\text{N}_2$ . The reaction mixture is cooled to room temperature, a few drops of  $\text{H}_2\text{O}$  are added and then it is concentrated under reduced pressure. The residue is purified by FCC (eluting with 20:1 methylene chloride:methanol) to yield 0.011 g (65%) of the pure desired product. MS APCI (-)  $m/z$  377 (M-1) detected;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  10.56 (broad s, 1H), 8.12 (s, 1H), 7.99 (s, 1H), 7.28 (s, 1H), 7.11 (d, 1H), 6.42 (broad, 1H), 3.70 (br s, 2H), 2.96 (d, 1H), 2.37 (s, 3H), 0.85 (m, 1H), 0.50 (m, 2H), 0.22 (m, 2H).

The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the

foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

## WHAT IS CLAIMED IS:

1. A compound of the formula



and pharmaceutically accepted salts, prodrugs and solvates thereof, wherein:

— is an optional bond, provided that one and only one nitrogen of the ring is double-bonded;

$R^1$ ,  $R^2$ ,  $R^9$  and  $R^{10}$  are independently selected from hydrogen, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-OR^3$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $NR^4C(O)OR^6$ ,  $-OC(O)R^3$ ,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-NR^4C(O)R^3$ ,  $-C(O)NR^3R^4$ ,  $-NR^5C(O)NR^3R^4$ ,  $-NR^5C(NCN)NR^3R^4$ ,  $-NR^3R^4$ , and

$C_1$ - $C_{10}$  alkyl,  $C_2$ - $C_{10}$  alkenyl,  $C_2$ - $C_{10}$  alkynyl,  $C_3$ - $C_{10}$  cycloalkyl,  $C_3$ - $C_{10}$  cycloalkylalkyl,  $-S(O)_j(C_1-C_6 \text{ alkyl})$ ,  $-S(O)_j(CR^4R^5)_m\text{-aryl}$ , aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl,  $-O(CR^4R^5)_m\text{-aryl}$ ,  $-NR^4(CR^4R^5)_m\text{-aryl}$ ,  $-O(CR^4R^5)_m\text{-heteroaryl}$ ,  $-NR^4(CR^4R^5)_m\text{-heteroaryl}$ ,  $-O(CR^4R^5)_m\text{-heterocyclyl}$  and  $-NR^4(CR^4R^5)_m\text{-heterocyclyl}$ , where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $-OC(O)R^3$ ,  $-NR^4C(O)OR^6$ ,  $-NR^4C(O)R^3$ ,

$-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{NCN})\text{NR}^3\text{R}^4$ ,  $-\text{OR}^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

$\text{R}^3$  is selected from hydrogen, trifluoromethyl, and

$\text{C}_1\text{-C}_{10}$  alkyl,  $\text{C}_2\text{-C}_{10}$  alkenyl,  $\text{C}_2\text{-C}_{10}$  alkynyl,  $\text{C}_3\text{-C}_{10}$  cycloalkyl,  $\text{C}_3\text{-C}_{10}$  cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-\text{NR}'\text{SO}_2\text{R}''''$ ,  $-\text{SO}_2\text{NR}'\text{R}''$ ,  $-\text{C}(\text{O})\text{R}'$ ,  $-\text{C}(\text{O})\text{OR}'$ ,  $-\text{OC}(\text{O})\text{R}'$ ,  $-\text{NR}'\text{C}(\text{O})\text{OR}''''$ ,  $-\text{NR}'\text{C}(\text{O})\text{R}''$ ,  $-\text{C}(\text{O})\text{NR}'\text{R}''$ ,  $-\text{SR}'$ ,  $-\text{S}(\text{O})\text{R}''''$ ,  $-\text{SO}_2\text{R}''''$ ,  $-\text{NR}'\text{R}''$ ,  $-\text{NR}'\text{C}(\text{O})\text{NR}''\text{R}''''$ ,  $-\text{NR}'\text{C}(\text{NCN})\text{NR}''\text{R}''''$ ,  $-\text{OR}'$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

$\text{R}'$ ,  $\text{R}''$  and  $\text{R}'''$  independently are selected from hydrogen, lower alkyl, lower alkenyl, aryl and arylalkyl;

$\text{R}''''$  is selected from lower alkyl, lower alkenyl, aryl and arylalkyl; or

Any two of  $\text{R}'$ ,  $\text{R}''$ ,  $\text{R}'''$  or  $\text{R}''''$  can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl; or

$\text{R}^3$  and  $\text{R}^4$  can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from

halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl; or

R<sup>4</sup> and R<sup>5</sup> independently represent hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; or

R<sup>4</sup> and R<sup>5</sup> together with the atom to which they are attached form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R''', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl;

R<sup>6</sup> is selected from trifluoromethyl, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclalkyl, where each alkyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl;

R<sup>7</sup> is selected from hydrogen, and



C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclalkyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -SO<sub>2</sub>R<sup>6</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl;

W is selected from heteroaryl, heterocyclyl, -C(O)OR<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -C(O)NR<sup>4</sup>OR<sup>3</sup>, -C(O)R<sup>4</sup>OR<sup>3</sup>, -C(O)(C<sub>3</sub>-C<sub>10</sub> cycloalkyl), -C(O)(C<sub>1</sub>-C<sub>10</sub> alkyl), -C(O)(aryl), -C(O)(heteroaryl) and -C(O)(heterocyclyl), each of which is optionally substituted with 1-5 groups independently selected from

-NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, -R<sup>2</sup>, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, and C<sub>2</sub>-C<sub>10</sub> alkynyl, each of which is optionally substituted with 1 or 2 groups independently selected from -NR<sup>3</sup>R<sup>4</sup> and -OR<sup>3</sup>;

R<sup>8</sup> is selected from hydrogen, -SCF<sub>3</sub>, -Cl, -Br, -F, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -OR<sup>3</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, and


C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, -S(O)<sub>j</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), -S(O)<sub>j</sub>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclalkyl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-

aryl,  $-\text{NR}^4(\text{CR}^4\text{R}^5)_m\text{-aryl}$ ,  $-\text{O}(\text{CR}^4\text{R}^5)_m\text{-heteroaryl}$ ,  $-\text{NR}^4(\text{CR}^4\text{R}^5)_m\text{-heteroaryl}$ ,  $-\text{O}(\text{CR}^4\text{R}^5)_m\text{-heterocyclyl}$  and  $-\text{NR}^4(\text{CR}^4\text{R}^5)_m\text{-heterocyclyl}$ , where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-\text{NR}^4\text{SO}_2\text{R}^6$ ,  $-\text{SO}_2\text{NR}^3\text{R}^4$ ,  $-\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{OR}^3$ ,  $-\text{OC}(\text{O})\text{R}^3$ ,  $-\text{NR}^4\text{C}(\text{O})\text{OR}^6$ ,  $-\text{NR}^4\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{NCN})\text{NR}^3\text{R}^4$ ,  $-\text{OR}^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

$m$  is 0, 1, 2, 3, 4 or 5; and

$j$  is 1 or 2.

2. A compound of the formula



and pharmaceutically accepted salts, prodrugs and solvates thereof, wherein:

— is an optional bond, provided that one and only one nitrogen of the ring is double-bonded;

$\text{R}^1$ ,  $\text{R}^9$  and  $\text{R}^{10}$  are independently selected from hydrogen, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-\text{OR}^3$ ,  $-\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{OR}^3$ ,  $\text{NR}^4\text{C}(\text{O})\text{OR}^6$ ,  $-\text{OC}(\text{O})\text{R}^3$ ,  $-\text{NR}^4\text{SO}_2\text{R}^6$ ,  $-\text{SO}_2\text{NR}^3\text{R}^4$ ,  $-\text{NR}^4\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{NCN})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^3\text{R}^4$ , and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, -S(O)<sub>j</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), -S(O)<sub>j</sub>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heteroaryl, -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heteroaryl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heterocyclyl and -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heterocyclyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

R<sup>3</sup> is selected from hydrogen, trifluoromethyl, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -SR<sup>3</sup>, -S(O)R<sup>3</sup>, -SO<sub>2</sub>R<sup>6</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> independently are selected from hydrogen, lower alkyl, lower alkenyl, aryl and arylalkyl;

R<sup>''''</sup> is selected from lower alkyl, lower alkenyl, aryl and arylalkyl; or

Any two of R', R'', R''' or R<sup>''''</sup> can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl; or

R<sup>3</sup> and R<sup>4</sup> can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl and heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R<sup>''''</sup>, -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR<sup>''''</sup>, -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R<sup>''''</sup>, -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl; or

R<sup>4</sup> and R<sup>5</sup> independently represent hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; or

R<sup>4</sup> and R<sup>5</sup> together with the atom to which they are attached form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R<sup>''''</sup>, -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR<sup>''''</sup>, -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R<sup>''''</sup>, -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl;

R<sup>6</sup> is selected from trifluoromethyl, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl, arylalkyl, heteraryl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, where each alkyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

R<sup>7</sup> is selected from hydrogen, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -SO<sub>2</sub>R<sup>6</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

W is selected from heteroaryl, heterocyclyl, -C(O)OR<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -C(O)NR<sup>4</sup>OR<sup>3</sup>, -C(O)R<sup>4</sup>OR<sup>3</sup>, -C(O)(C<sub>3</sub>-C<sub>10</sub> cycloalkyl), -C(O)(C<sub>1</sub>-C<sub>10</sub> alkyl), -C(O)(aryl), -C(O)(heteroaryl) and -C(O)(heterocyclyl), each of which is optionally substituted with 1-5 groups independently selected from

-NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, -R<sup>2</sup>, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, and C<sub>2</sub>-C<sub>10</sub> alkynyl, each of which is optionally substituted with 1 or 2 groups independently selected from -NR<sup>3</sup>R<sup>4</sup> and -OR<sup>3</sup>;


R<sup>8</sup> is selected from hydrogen, -SCF<sub>3</sub>, -Cl, -Br, -F, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -OR<sup>3</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, -S(O)<sub>j</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), -S(O)<sub>j</sub>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, aryl, arylalkyl, heteraryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-aryl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heteroaryl, -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heteroaryl, -O(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heterocyclyl and -NR<sup>4</sup>(CR<sup>4</sup>R<sup>5</sup>)<sub>m</sub>-heterocyclyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

m is 0, 1, 2, 3, 4 or 5; and

j is 1 or 2.

3. A compound of the formula



and pharmaceutically accepted salts, prodrugs and solvates thereof, wherein:

$R^1$ ,  $R^2$ , and  $R^9$  are independently selected from hydrogen, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-OR^3$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $NR^4C(O)OR^6$ ,  $-OC(O)R^3$ ,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-NR^4C(O)R^3$ ,  $-C(O)NR^3R^4$ ,  $-NR^5C(O)NR^3R^4$ ,  $-NR^5C(NCN)NR^3R^4$ ,  $-NR^3R^4$ , and  $C_1$ - $C_{10}$  alkyl,  $C_2$ - $C_{10}$  alkenyl,  $C_2$ - $C_{10}$  alkynyl,  $C_3$ - $C_{10}$  cycloalkyl,  $C_3$ - $C_{10}$  cycloalkylalkyl,  $-S(O)_j(C_1$ - $C_6$  alkyl),  $-S(O)_j(CR^4R^5)_m$ -aryl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl,  $-O(CR^4R^5)_m$ -aryl,  $-NR^4(CR^4R^5)_m$ -aryl,  $-O(CR^4R^5)_m$ -heteroaryl,  $-NR^4(CR^4R^5)_m$ -heteroaryl,  $-O(CR^4R^5)_m$ -heterocyclyl and  $-NR^4(CR^4R^5)_m$ -heterocyclyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $-OC(O)R^3$ ,  $-NR^4C(O)OR^6$ ,  $-NR^4C(O)R^3$ ,  $-C(O)NR^3R^4$ ,  $-NR^3R^4$ ,  $-NR^5C(O)NR^3R^4$ ,  $-NR^5C(NCN)NR^3R^4$ ,  $-OR^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

$R^3$  is selected from hydrogen, trifluoromethyl, and

$C_1$ - $C_{10}$  alkyl,  $C_2$ - $C_{10}$  alkenyl,  $C_2$ - $C_{10}$  alkynyl,  $C_3$ - $C_{10}$  cycloalkyl,  $C_3$ - $C_{10}$  cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl,

and heterocyclalkyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocycl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-NR'SO_2R''''$ ,  $-SO_2NR'R''$ ,  $-C(O)R'$ ,  $-C(O)OR'$ ,  $-OC(O)R'$ ,  $-NR'C(O)OR''''$ ,  $-NR'C(O)R''$ ,  $-C(O)NR'R''$ ,  $-SR''''$ ,  $-S(O)R''''$ ,  $-SO_2R'$ ,  $-NR'R''$ ,  $-NR'C(O)NR''R'''$ ,  $-NR'C(NCN)NR''R'''$ ,  $-OR'$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocycl, and heterocyclalkyl;

$R'$ ,  $R''$  and  $R'''$  independently are selected from hydrogen, lower alkyl, lower alkenyl, aryl and arylalkyl;

$R''''$  is selected from lower alkyl, lower alkenyl, aryl and arylalkyl; or

Any two of  $R'$ ,  $R''$ ,  $R'''$  or  $R''''$  can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocycl, and heterocyclalkyl; or

$R^3$  and  $R^4$  can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-NR'SO_2R''''$ ,  $-SO_2NR'R''$ ,  $-C(O)R'$ ,  $-C(O)OR'$ ,  $-OC(O)R'$ ,  $-NR'C(O)OR''''$ ,  $-NR'C(O)R''$ ,  $-C(O)NR'R''$ ,  $-SO_2R''''$ ,  $-NR'R''$ ,  $-NR'C(O)NR''R'''$ ,  $-NR'C(NCN)NR''R'''$ ,  $-OR'$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocycl, and heterocyclalkyl; or



R<sup>4</sup> and R<sup>5</sup> independently represent hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; or

R<sup>4</sup> and R<sup>5</sup> can be taken together with the atom to which they are attached to form a 4 to 10 membered carbocyclic, heteroaryl or heterocyclic ring, each of which is optionally substituted with one to three groups independently selected from halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R'', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

R<sup>6</sup> is selected from trifluoromethyl, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, where each alkyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR'SO<sub>2</sub>R''', -SO<sub>2</sub>NR'R'', -C(O)R', -C(O)OR', -OC(O)R', -NR'C(O)OR''', -NR'C(O)R'', -C(O)NR'R'', -SO<sub>2</sub>R''', -NR'R', -NR'C(O)NR''R''', -NR'C(NCN)NR''R''', -OR', aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

R<sup>7</sup> is selected from hydrogen, and

C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>2</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>3</sub>-C<sub>10</sub> cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro,

trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-\text{NR}^4\text{SO}_2\text{R}^6$ ,  $-\text{SO}_2\text{NR}^3\text{R}^4$ ,  $-\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{OR}^3$ ,  $-\text{OC}(\text{O})\text{R}^3$ ,  $-\text{NR}^4\text{C}(\text{O})\text{OR}^6$ ,  $-\text{NR}^4\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{SO}_2\text{R}^6$ ,  $-\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{NCN})\text{NR}^3\text{R}^4$ ,  $-\text{OR}^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

A is selected from  $-\text{C}(\text{O})\text{OR}^3$  or  $-\text{C}(\text{O})\text{NR}^4\text{OR}^3$ ;


$\text{R}^8$  is selected from hydrogen,  $-\text{SCF}_3$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{F}$ , cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-\text{OR}^3$ ,  $-\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{OR}^3$ ,  $-\text{NR}^4\text{C}(\text{O})\text{OR}^6$ ,  $-\text{OC}(\text{O})\text{R}^3$ ,  $-\text{NR}^4\text{SO}_2\text{R}^6$ ,  $-\text{SO}_2\text{NR}^3\text{R}^4$ ,  $-\text{NR}^4\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^3\text{R}^4$ , and

$\text{C}_1\text{-C}_{10}$  alkyl,  $\text{C}_2\text{-C}_{10}$  alkenyl,  $\text{C}_2\text{-C}_{10}$  alkynyl,  $\text{C}_3\text{-C}_{10}$  cycloalkyl,  $\text{C}_3\text{-C}_{10}$  cycloalkylalkyl,  $-\text{S}(\text{O})_j(\text{C}_1\text{-C}_6 \text{ alkyl})$ ,  $-\text{S}(\text{O})_j(\text{CR}^4\text{R}^5)_m\text{-aryl}$ , aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl,  $-\text{O}(\text{CR}^4\text{R}^5)_m\text{-aryl}$ ,  $-\text{NR}^4(\text{CR}^4\text{R}^5)_m\text{-aryl}$ ,  $-\text{O}(\text{CR}^4\text{R}^5)_m\text{-heteroaryl}$ ,  $-\text{NR}^4(\text{CR}^4\text{R}^5)_m\text{-heteroaryl}$ ,  $-\text{O}(\text{CR}^4\text{R}^5)_m\text{-heterocyclyl}$  and  $-\text{NR}^4(\text{CR}^4\text{R}^5)_m\text{-heterocyclyl}$ , where each alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl portion is optionally substituted with one to five groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-\text{NR}^4\text{SO}_2\text{R}^6$ ,  $-\text{SO}_2\text{NR}^3\text{R}^4$ ,  $-\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{OR}^3$ ,  $-\text{OC}(\text{O})\text{R}^3$ ,  $-\text{NR}^4\text{C}(\text{O})\text{OR}^6$ ,  $-\text{NR}^4\text{C}(\text{O})\text{R}^3$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{C}(\text{NCN})\text{NR}^3\text{R}^4$ ,  $-\text{OR}^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl;

m is 0, 1, 2, 3, 4 or 5; and

j is 1 or 2.

4. A compound according to claim 3 which is



5. A compound according to claim 4 wherein

R<sup>7</sup> is C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkylalkyl, C<sub>3</sub>-C<sub>7</sub> heterocycloalkyl or C<sub>3</sub>-C<sub>7</sub> heterocycloalkylalkyl, each of which can be optionally substituted with 1 - 3 groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR<sup>4</sup>SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, -C(O)R<sup>3</sup>, -C(O)OR<sup>3</sup>, -OC(O)R<sup>3</sup>, -SO<sub>2</sub>R<sup>6</sup>, -NR<sup>4</sup>C(O)OR<sup>6</sup>, -NR<sup>4</sup>C(O)R<sup>3</sup>, -C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -NR<sup>5</sup>C(NCN)NR<sup>3</sup>R<sup>4</sup>, -OR<sup>3</sup>, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl;

R<sup>9</sup> is hydrogen or halogen;

R<sup>1</sup> is lower alkyl or halogen; and

R<sup>8</sup> is -OCF<sub>3</sub> or halogen.

6. A compound according to claim 5 wherein R<sup>9</sup> is fluoro.
7. A compound according to claim 6 wherein R<sup>1</sup> is methyl or chloro.
8. A compound according to claim 7 wherein R<sup>8</sup> is chloro or bromo.

9. A compound according to claim 5 wherein A is  $-C(O)NR^4OR^3$ .

10. A compound according to claim 1 wherein

$R^7$  is  $C_1$ - $C_{10}$  alkyl,  $C_3$ - $C_7$  cycloalkyl,  $C_3$ - $C_7$  cycloalkylalkyl,  $C_3$ - $C_7$  heterocycloalkyl or  $C_3$ - $C_7$  heterocycloalkylalkyl, each of which can be optionally substituted with 1 - 3 groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $-OC(O)R^3$ ,  $-SO_2R^6$ ,  $-NR^4C(O)OR^6$ ,  $-NR^4C(O)R^3$ ,  $-C(O)NR^3R^4$ ,  $-NR^3R^4$ ,  $-NR^5C(O)NR^3R^4$ ,  $-NR^5C(NCN)NR^3R^4$ ,  $-OR^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl;

$R^8$  is  $-OCF_3$ ,  $-Br$  or  $-Cl$ ,  $R^2$  is hydrogen, and  $R^1$  is lower alkyl or halogen;

$R^9$  is hydrogen or halogen;

$R^{10}$  is hydrogen; and

W is  $-C(O)OR^3$  or  $-C(O)NR^4OR^3$ .

11. A compound according to claim 10 wherein W is  $-C(O)NR^4OR^3$ .

12. A compound according to claim 2 wherein

$R^7$  is  $C_1$ - $C_{10}$  alkyl,  $C_3$ - $C_7$  cycloalkyl or  $C_3$ - $C_7$  cycloalkylalkyl, each of which can be optionally substituted with 1 - 3 groups independently selected from oxo, halogen, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido,  $-NR^4SO_2R^6$ ,  $-SO_2NR^3R^4$ ,  $-C(O)R^3$ ,  $-C(O)OR^3$ ,  $-OC(O)R^3$ ,  $-SO_2R^3$ ,  $-NR^4C(O)OR^6$ ,  $-NR^4C(O)R^3$ ,  $-C(O)NR^3R^4$ ,  $-NR^3R^4$ ,  $-NR^5C(O)NR^3R^4$ ,  $-NR^5C(NCN)NR^3R^4$ ,  $-OR^3$ , aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclalkyl;

$R^8$  is  $-OCF_3$ ,  $-Br$  or  $-Cl$ , and  $R^1$  is lower alkyl or halogen;

R<sup>9</sup> is hydrogen or halogen;

R<sup>10</sup> is hydrogen; and

W is -C(O)OR<sup>3</sup> or -C(O)NR<sup>4</sup>OR<sup>3</sup>.

13. A compound according to claim 12 wherein W is -C(O)NR<sup>4</sup>OR<sup>3</sup>.

14. A compound according to claim 1 which is selected from

7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-3H-benzoimidazole-5-carboxylic acid  
cyclopropylmethoxy-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid  
cyclopropylmethoxy-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic  
acid (2-hydroxy-ethoxy)-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic  
acid (2,3-dihydroxy-propoxy)-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-  
benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide;

[6-(5-Amino-[1,3,4]oxadiazol-2-yl)-4-fluoro-1H-benzoimidazol-5-yl]-(4-bromo-2-  
methyl-phenyl)-amine;

1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazol-5-yl]-2-  
hydroxy-ethanone;

1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methoxy-  
ethanone;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic  
acid (2-hydroxy-1,1-dimethyl-ethoxy)-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-furan-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide;  
6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide;  
6-(4-Bromo-2-fluoro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide; and  
6-(2,4-Dichloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide.

15. A composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

16. A method of inhibiting MEK activity in a mammal comprising administering an effective amount of a compound of claim 1 to the mammal.

17. A method for treatment of a hyperproliferative disorder in a mammal comprising administering an effective amount of a compound of claim 1 to the mammal.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/07864


<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : A61K 31/4184; C07D 235/04, 235/06 US CL : 514/394; 548/304.4, 304.7, 310.4 According to International Patent Classification (IPC) or to both national classification and IPC														
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/394; 548/304.4, 304.7, 310.4  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) STN CAS ONLINE														
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>														
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
X	WO 00/42022 A1 (WARNER-LAMBERT COMPANY) 20 July 2000 (20/07/00), see entire document, especially the compound listed on page 38, lines 25-26 (CA Registry No. 284030-61-1).	1-7, 15-17												
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"><tr><td>* Special categories of cited documents:</td><td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td></tr><tr><td>"A" document defining the general state of the art which is not considered to be of particular relevance</td><td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td></tr><tr><td>"B" earlier application or patent published on or after the international filing date</td><td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td></tr><tr><td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td><td>"&amp;" document member of the same patent family</td></tr><tr><td>"O" document referring to an oral disclosure, use, exhibition or other means</td><td></td></tr><tr><td>"P" document published prior to the international filing date but later than the priority date claimed</td><td></td></tr></table>			* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"B" earlier application or patent published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family	"O" document referring to an oral disclosure, use, exhibition or other means		"P" document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
"B" earlier application or patent published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family													
"O" document referring to an oral disclosure, use, exhibition or other means														
"P" document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 10 June 2003 (10.06.2003)		Date of mailing of the international search report <b>30 JUN 2003</b>												
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450 Facsimile No. (703)305-3230		Authorized officer Laura L. Stockton, Ph.D. Telephone No. 703/308-1235 												

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
957		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole		437.02 [M+1]
958		1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.37 (d, 1H), 8.03 (dd, 1H), 7.65 (d, 1H), 7.38-7.50 (m, 5H), 6.24 (d, 1H), 4.49 (s, 2H), 4.03 (q, 2H), 2.68 (s, 3H), 1.24 (t, 3H)	
959		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-[5-(trifluoromethyl)-1H-tetrazol-1-yl]-1H-benzimidazole		474.99 [M+1]
960		5-(3,5-difluorophenyl)-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.91 (d, 1H), 7.64 (d, 1H), 7.27-7.50 (m, 5H), 7.10-7.18 (m, 3H), 6.77 (m, 1H), 6.20 (d, 1H), 4.36 (s, 2H), 4.02 (q, 2H), 1.25 (t, 3H)	
961		(1Z)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-methyloxime	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.94 (d, 1H), 7.73 (dd, 1H), 7.73 (dd, 1H), 7.63 (d, 1H), 7.52 (m, 1H), 7.38-7.40 (m, 3H), 7.27 (m, 1H), 6.19 (d, 1H), 4.32 (s, 2H), 3.87-4.10 (m, 5H), 2.30 (s, 3H), 1.22 (t, 3H.)	
962		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole		425.10 [M+1]
963		(1E)-1-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)ethanone oxime	<sup>1</sup> H NMR (CD <sub>3</sub> OD): 7.81 (d, 1H), 7.64-7.68 (m, 2H), 7.42-7.53 (m, 2H), 7.16-7.32 (m, 3H), 6.28 (d, 1H), 4.44 (s, 2H), 4.12 (q, 2H), 2.30 (s, 3H), 1.23 (t, 3H)	
964		(1E)-1-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)ethanone O-ethyloxime	Syn and anti <sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.95 (d, 1H+H1), 7.73 (dd, 1H+H1), 7.63 (d, 1H+H1), 7.44 (m, 1H+H1), 7.24-7.33 (m, 3H+H1), [6.18 (d, 1H+H1), 7.13 (d, 1H+H1)], 4.33 (s, 2H), [4.26 (q, 2H), 4.13 (q, 2H1)], 3.98 (q, 2H+2H1), [2.31 (s, 3H), 2.27 (s, 3H1)], [1.35 (t, 3H), 1.28 (t, 3H1)], 1.22 (t, 3H+H1)]	



TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
965		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(5-pyridin-3-yl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	<sup>1</sup> H NMR (d <sub>6</sub> DMSO): 9.40(1H, s), 8.87(1H, s), 8.63(1H, d), 8.41(1H, s), 8.20(1H, d), 8.01(1H, d), 7.76(2H, m), 7.59(1H, m), 7.48-7.41(2H, m), 6.50(1H, s), 4.61(2H, s), 4.31(2H, q), 1.19(3H, t)	484.5 [M+1]
966		2-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)-4-methoxybenzaldehyde	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 9.86 (s, 1H), 8.03 (d, 1H), 7.26-7.78 (m, 7H), 6.94-7.18 (m, 3H), 6.22 (dd, 1H), 4.38 (s, 2H), 4.04 (q, 2H), 3.91 (s, 3H), 1.27 (t, 3H)	
967		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-N,N-dimethyl-1H-benzimidazole-5-carboxamide	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.78 (s, 1H), 7.63 (s, 1H), 7.10-7.48 (m, 6H), 6.18 (s, 1H), 4.35 (s, 2H), 4.00 (q, 2H), 3.13 (s, 3H), 3.05 (s, 3H), 1.22 (t, 3H)	
968		5-(3,5-dimethylisoxazol-4-yl)-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.62-7.64 (m, 2H), 7.25-7.50 (m, 4H), 7.11-7.18 (m, 2H), 6.21 (d, 1H), 4.36 (s, 2H), 4.02 (q, 2H), 2.42 (s, 3H), 2.29 (s, 3H), 1.26 (t, 3H)	
969		5-bromo-2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazole	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.87 (d, 1H), 7.63 (d, 1H), 7.50 (m, 1H), 7.35-7.42 (m, 4H), 7.18 (d, 1H), 6.18 (d, 1H), 4.31 (s, 2H), 3.96 (q, 2H), 1.21 (t, 3H)	
970		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(5-pyridin-4-yl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	Dihydrochloride <sup>1</sup> H NMR (d <sub>6</sub> DMSO): 8.98(2H, m), 8.43(1H, s), 8.35(2H, m), 8.19(1H, d), 7.99(1H, d), 7.75(1H, d), 7.59-7.569(1H, m), 7.50-7.41(2H, m), 6.49(1H, s), 4.58(2H, s), 4.30(2H, q), 1.19(3H, t)	484.5 [M+1]
971		(1E)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone oxime	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 10.92 (br s, 1H), 8.58 (s, 1H), 7.52-7.64 (m, 3H), 7.29 (d, 1H), 6.20 (d, 1H), 4.36 (s, 2H), 3.97 (q, 2H), 2.38 (s, 3H), 1.21 (t, 3H)	
972		(1E)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-ethylloxime	Syn and anti <sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.95 (d, 1H+H1), 7.73 (dd, 1H+H1), 7.63 (d, 1H+H1), 7.44 (m, 1H+H1), 7.24-7.33 (m, 3H+H1), [6.18 (d, 1H+H1), 7.13 (d, 1H+H1)], 4.33 (s, 2H), [4.26 (q, 2H), 4.13 (q, 2H1)], 3.98 (q, 2H+2H1), [2.31 (s, 3H), 2.27 (s, 3H1)], [1.35 (t, 3H), 1.28 (t, 3H1)], 1.22 (t, 3H+H1)]	









TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
973		(1E)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-(tert-butyl)oxime	Syn and anti <sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.00 (d, 1H+H1), 7.75 (dd, 1H+H1), 7.63 (d, 1H+H1), 7.27 (d, 1H+H1), 7.38-7.40 (m, 3H+H1), 6.20 (d, 1H)[6.18(d, 1H)], 4.32 (s, 2H+H1), 3.99 (q, 2H+H1), 2.28 (s, 3H), [2.27(s, 3H)], 1.37 (s, 9H), [1.32 (s, 9H)], 1.22 (t, 3H, t3H)	
974		(1E)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-tetrahydro-2H-pyran-2-yl oxime	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.95 (d, 1H), 7.79 (dd, 1H), 7.63 (d, 1H), 7.52 (m, 1H), 7.37-7.40 (m, 3H), 7.26 (dd, 1H), 6.19 (d, 1H), 5.43 (m, 1H), 4.32 (s, 2H), 3.85-4.02 (m, 4H), 3.66 (m, 2H), 1.62-1.92 (m, 4H), 1.22 (t, 3H)	
975		(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)acetonitrile	7.66 (m, 1H), 7.63 (d, 1H), 7.24-7.49 (m, 5H), 7.13 (m, 1H), 6.17 (d, 1H), 4.34 (s, 2H), 3.99 (q, 2H), 3.87 (s, 2H), 1.22 (t, 3H)	
976		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole	<sup>1</sup> H NMR (d <sub>6</sub> DMSO): 8.17(1H, s), 8.04-7.98(2H, m), 7.71(1H, d), 7.55-7.50(2H, m), 7.45-7.43(1H, m), 7.29-7.24(1H, m), 6.40(1H, d), 4.77(2H, s), 4.36(2H, q), 2.58(3H, s), 1.24(3H, t)	403.6 [M+1]
977		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	<sup>1</sup> H NMR (CD <sub>3</sub> OD): 8.79 (s, 1H), 8.31 (d, 1H), 7.69 (s, 1H), 7.60 (d, H), 7.46 (dd, 1H), 7.30 (d, 2H), 7.16-7.20 (m, 1H), 6.37 (s, 1H), 4.55 (s, 2H), 4.20 (q, 2H), 1.24 (t, 3H)	
978		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-imidazo[4,5-c]pyridine	<sup>1</sup> H NMR (CD <sub>3</sub> OD): 8.79 (s, 1H), 8.31 (d, 1H), 7.69 (s, 1H), 7.60 (d, H), 7.46 (dd, 1H), 7.30 (d, 2H), 7.16-7.20 (m, 1H), 6.37 (s, 1H), 4.55 (s, 2H), 4.20 (q, 2H), 1.24 (t, 3H)	
979		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-6-carbonitrile		
980		1-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)propan-1-one		

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
981		2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1-(2-methoxyethyl)-1H-benzimidazole-5-carbonitrile		
982		2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1-(3-methoxypropyl)-1H-benzimidazole-5-carbonitrile		390.1 [M+1]; 388.3 [M-1]
983		1-(2-cyanoethyl)-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-5-carbonitrile		371.3 [M+1]
984		1-ethyl-5-(5-ethyl-1,3,4-oxadiazol-2-yl)-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	1 H NMR (d6DMSO): 8.17(1H, s), 8.01-7.94(2H, m), 7.71(1H, s), 7.53-7.51(2H, m), 7.44-7.42(1H, m), 7.28-7.24(1H, m), 6.39(1H, s), 4.73(2H, s), 4.33(2H, q), 2.94(2H, q), 1.32(3H, t), 1.23(3H, t)	417.5 [M+1]
985		5-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)-1,3,4-oxadiazol-2-amine	1 H NMR (d6DMSO): 8.17(1H, s), 8.01-7.94(2H, m), 7.71(1H, s), 7.53-7.51(2H, m), 7.44-7.42(1H, m), 7.28-7.24(1H, m), 6.39(1H, s), 4.73(2H, s), 4.33(2H, q), 2.94(2H, q), 1.32(3H, t), 1.23(3H, t)	417.5 [M+1]
986		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl](hydroxy)methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile	1H NMR (CDCl3): 7.93 (d, 1H), 7.67 (d, 1H), 7.55 (dd, 1H), 7.40 (dd, 1H); 7.24 (m, 1H), 7.10-7.15 (m, 2H), 6.20 (d, 1H), 6.01 (s, 1H), 4.98 (br, s, 1H); 4.04 (q, 2H), 1.21 (t, 3H)	
987		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-imidazo[4,5-b]pyridine		322.3 [M+1]; 320.3 [M-1]
988		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl](fluoro)methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile	1H NMR (CDCl3): 8.07 (d, 1H), 7.77 (d, 1H), 7.60 (dd, 1H), 7.47 (dd, 1H), 7.11-7.23 (m, 3H), 6.85 [d, 1H, J(H,F)=47.4], 6.50 (t, 1H), 4.27 (m, 2H), 1.36 (t, 3H)	

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
989		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(5-pyridin-3-yl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	<sup>1</sup> H NMR (d <sub>6</sub> DMSO): 9.41(1H, s), 8.88(1H, d), 8.67(1H, d), 8.46(1H, s), 8.25(1H, d), 8.10(1H, d), 7.81-7.72(2H, m), 7.55-7.46(3H, m), 7.30-7.26(1H, m), 6.43(1H, s), 4.84(2H, s), 4.40(2H, q), 1.27(3H, t)	
990		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-isoxazol-5-yl-1H-benzimidazole	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.30 (d, 1H), 8.16 (dd, 1H), 7.76 (dd, 1H), 7.64 (d, 1H), 7.38-7.50 (m, 2H), 7.25-7.32 (m, 2H), 7.14 (m, 1H), 6.52 (d, 1H), 6.21 (d, 1H), 4.37 (s, 2H), 4.01 (q, 2H), 1.25 (t, 3H)	
991		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl](hydroxymethyl)-1H-benzimidazole-5-carbonitrile	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.02 (dd, 1H), 7.47-7.62 (m 5H), 7.39 (d, 1H), 7.18 (m, 1H), 6.12 (d, 1H), 6.00 (s, 1H), 5.05 (brs, 1H), 3.83 (q, 2H), 1.08 (t, 3H)	
992		1-ethyl-2-(fluoro[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-5-carbonitrile	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.12 (dd, 1H), 7.73 (d, 1H), 7.62 (dd, 1H), 7.44-7.53 (m, 2H), 7.31-7.39 (m, 2H), 7.17 (m, 1H), 6.83 (d, 1H, J <sub>H,F</sub> =46.8 MZ), 4.39 (m, 2H), 1.43 (t, 3H)	
993		methyl 5-[(5-cyano-1-ethyl-1H-benzimidazol-2-yl)oxy]-1-(3-fluorophenyl)-1H-pyrazole-4-carboxylate		406.3 [M+1]
994		2-[[1-(2,5-difluorophenyl)-1H-pyrazol-5-yl](difluoromethyl)-1-ethyl-1H-benzimidazole-5-carbonitrile	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.09 (m, 1H), 7.84 (M, 1H), 7.64 (dd, 1H), 7.48 (dd, 1H), 6.96-7.18 (m, 3H), 6.85 (d, 1H), 4.33 (q, 2H), 1.38 (t, 3H)	
995		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-6-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole		389.2 [M+1]
996		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole		403.3 [M+1]









TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
997		1-(2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazol-1-yl)acetone		349.3 [M+1]; 347.2 [M-1]
998		2-(difluoro[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.078 (m, 1H), 7.80 (m, 1H), 7.63 (dd, 1H), 7.48 (dd, 1H), 7.13-7.28 (m, 3H), 6.99 (m, 1H), 6.81 (d, 1H), 4.35 (q, 2H), 1.39 (t, 3H)	
999		2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1-(3-fluoropropyl)-1H-benzimidazole		353.2 [M+1]; 351.2 [M-1]
1000		1-(2-fluoroethyl)-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazole		339.2 [M+1]
1001		4-(2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazol-1-yl)butanenitrile		360.2 [M+1]; 358.3 [M-1]
1002		1-(1-ethyl-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazol-5-yl)-2,2,2-trifluoroethanone		417.2 [M+1]
1003		1-(2-((1-(3-chlorophenyl)-1H-pyrazol-5-yl)methyl)-1-ethyl-1H-benzimidazol-5-yl)-2,2,2-trifluoroethanone		
1004		5-chloro-2-((1-(2,5-difluorophenyl)-1H-pyrazol-5-yl)(difluoro)methyl)-1-ethyl-1H-benzimidazole		










TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1005		(1E)-1-(1-ethyl-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazol-5-yl)ethanone O-(2-ethoxyethyl)oxime	<sup>1</sup> H NMR, 8ppm (CDCl <sub>3</sub> ): 7.95(m, 1H), 7.72(dd, 1H), 7.63(d, 1H), 7.45(m, 1H), 7.11-7.31(m, 4H), 6.18(d, 1H), 4.34-4.37(m, 4H), 3.98(q, 2H), 3.77(t, 2H), 3.58(q, 2H), 2.33(s, 3H), 1.18-1.26(m, 6H)	
1006		1-ethyl-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-N,N-dimethyl-1H-benzimidazole-6-carboxamide	<sup>1</sup> H NMR, 8ppm (CDCl <sub>3</sub> ): 7.72(dd, 1H), 7.63(d, 1H), 7.40-7.49 (m, 2H), 7.24-7.34(m, 3H), 7.13(d, 1H), 6.18(d, 1H), 4.35(s, 2H), 3.99(q, 2H), 3.13(s, 2H), 3.07(s, 3H), 1.23(t, 3H)	
1007		3-ethyl-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-3H-imidazo[4,5-b]pyridine		
1008		2-((1-(2,5-difluorophenyl)-1H-pyrazol-5-yl)methyl)-3-ethyl-3H-imidazo[4,5-b]pyridine	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.19 (1H, d), 7.61 (1H, d), 7.48-7.44(1H, m), 7.28-7.13(4H, m), 6.14(1H, m), 4.44(2H, s), 3.93(2H, q), 1.17(3H, t),	356.2 [M+1]
1009		4-chloro-1-ethyl-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-imidazo[4,5-c]pyridine	<sup>1</sup> H nmr (CDCl <sub>3</sub> ): 8.19(1H, d), 7.61(1H, d), 7.48-7.44(1H, m), 7.28-7.13(4H, m), 6.14(1H, m), 4.44(2H, s), 3.93(2H, q), 1.17(3H, t)	356.2 [M+1]
1010		3-ethyl-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-5-methoxy-3H-imidazo[4,5-b]pyridine		392.3 [M+1]
1011		2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-3-isopropyl-3H-imidazo[4,5-b]pyridine		
1012		1-(cyclopropylmethyl)-2-((1-(2,5-difluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-imidazo[4,5-b]pyridine	<sup>1</sup> H-1 NMR (CDCl <sub>3</sub> ): 0.22 (m, 2H), 0.54(m, 2H), 0.95(m, 1H), 3.84(d, 2H), 4.32(s, 2H), 6.18(s, 2H), 7.13-7.25 (m, 4H), 7.65 (m, 2H), 8.52 (d, 1H)	366 [M+1]
1013		1-(cyclopropylmethyl)-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-imidazo[4,5-b]pyridine	<sup>1</sup> H-1 NMR (CDCl <sub>3</sub> ): 0.22 (m, 2H), 0.54(m, 2H), 0.95(m, 1H), 3.84(d, 2H), 4.32(s, 2H), 6.18(s, 2H), 7.13-7.25 (m, 4H), 7.65 (m, 2H), 8.52 (d, 1H)	366 [M+1]

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1014		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-propyl-1H-imidazo[4,5-b]pyridine	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 0.82(t, 3H), 1.62(h, 2H), 3.85(t, 2H), 4.33 (s, 2H), 6.20 (s, 1H), 7.05-7.25(m, 4H), 7.82(m, 2H), 8.53(d, 1H)	354 [M+1]
1015		2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-3-propyl-1H-imidazo[4,5-b]pyridine	(CDCl <sub>3</sub> ) 8.43(dd, J = 4.8, 1.5 Hz, 1H), 8.10 (dd, J = 8.0, 1.4 Hz, 1H), 7.64 (d, J = 2.1 Hz, 1H), 7.47 (m, 1H), 7.23-7.34 (m, 3H), 7.12 (m, 1H), 6.31 (d, J = 1.8 Hz, 1H), 4.60 (s, 2H), 4.03 (t, J = 7.7 Hz, 2H), 1.65 (m, 2H), 0.84 (t, J = 7.5 Hz, 3H)	m/z 366.2 [M+1]
1016		2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-methyl-3-propyl-3H-imidazo[4,5-b]pyridine	(CDCl <sub>3</sub> ) 7.85 (d, J = 7.8 Hz, 1H), 7.64 (d, J = 1.5 Hz, 1H), 7.43 (m, 1H), 7.24-7.29 (m, 2H), 7.12 (m, 1H), 7.06 (d, J = 8.1 Hz, 1H), 6.22 (d, J = 1.5 Hz, 1H), 4.33 (s, 2H), 4.01 (t, J = 7.5 Hz, 2H), 2.64 (s, 3H), 1.67 (m, 2H), 0.84 (t, J = 7.3 Hz, 3H)	m/z 350.3 [M+1]
1017		3-ethyl-2-([1-(3-(trifluoromethyl)phenyl)-1H-pyrazol-5-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl <sub>3</sub> ) 8.36 (dd, J = 4.8, 1.3 Hz, 1H), 7.98 (dd, J = 8.0, 1.3 Hz, 1H), 7.78 (m, 1H), 7.56-7.72 (m, 4H), 7.22 (dd, J = 8.1, 4.8 Hz, 1H), 6.29 (d, J = 1.5 Hz, 1H), 4.36 (s, 2H), 4.17 (q, J = 7.2 Hz, 2H), 1.27 (t, J = 7.2 Hz, 3H)	m/z 372.2 [M+1]
1018		3-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-methyl-3H-imidazo[4,5-b]pyridine	(CDCl <sub>3</sub> ) 7.85 (d, J = 8.1 Hz, 1H), 7.64 (d, J = 1.8 Hz, 1H), 7.44 (m, 1H), 7.24-7.29 (m, 2H), 7.13 (m, 1H), 7.06 (d, J = 8.1 Hz, 1H), 6.21 (d, J = 1.5 Hz, 1H), 4.33 (s, 2H), 4.12 (q, J = 7.3 Hz, 2H), 2.65 (s, 3H), 1.25 (t, J = 7.3 Hz, 3H)	m/z 336.2 [M+1]
1019		3-ethyl-2-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-5-methyl-3H-imidazo[4,5-b]pyridine		m/z 337.2 [M+1]
1020		8-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-2,9-dimethyl-9H-purine	(CDCl <sub>3</sub> ) 8.81 (s, 1H), 7.87 (m, 2H), 7.69 (d, 1H), 6.67 (dd, 1H), 6.33 (s, 1H), 4.86 (s, 2H), 3.90 (s, 3H), 2.79 (s, 3H)	
1021		2-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-1,6-dimethyl-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 8.82 (s, 1H), 7.88 (m, 1H), 7.84 (m, 1H), 7.67 (s, 1H), 7.11 (s, 1H), 6.69 (m, 1H), 6.27 (s, 1H), 4.85 (s, 2H), 3.81 (s, 3H), 2.66 (s, 3H)	m/z 323 [M+1]

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1022		8-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-2,9-dimethyl-9H-purine	(CDCl <sub>3</sub> ) 8.90 (s, 1H), 7.63(d, 1H), 7.42 (m, 1H), 7.26 (m, 2H), 7.12 (m, 1H), 6.16 (d, 1H), 4.32 (s, 2H), 3.60 (s, 3H), 2.78 (s, 3H)	
1023		2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1-methyl-6-(trifluoromethyl)-1H-benzimidazole	(CDCl <sub>3</sub> ) 7.79 (d, 1H), 7.59 (d, 2H), 7.54 (m, 2H), 7.30 (m, 2H), 7.16 (m, 1H), 6.12 (d, 1H), 4.38 (s, 2H), 3.61 (s, 3H),	
1024		9-ethyl-8-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-2-methyl-9H-purine	(CDCl <sub>3</sub> ) 8.82 (s, 1H), 7.88 (m, 1H), 7.83 (q, 1H), 7.69 (d, 1H), 6.69 (m, 1H), 6.35 (s, 1H), 4.95 (s, 2H), 4.44 (q, 2H), 2.82 (s, 3H), 1.52 (t, 3H)	
1025		3-ethyl-2-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-3H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 8.78 (s, 1H), 8.38 (d, J = 3.9 Hz, 1H), 7.83-7.89 (m, 2H), 7.66 (s, 1H), 7.56 (d, J = 3.9 Hz, 1H), 7.25 (d, J = 9.9 Hz, 1H), 6.70 (d, J = 9.9 Hz, 1H), 4.81 (s, 2H), 4.39 (q, J = 5.4 Hz, 2H), 1.49 (t, J = 5.4 Hz, 3H)	m/z 323 [M+1]
1026		1-ethyl-2-([1-(3-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-6-methyl-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 8.82 (s, 1H), 8.28 (br d, J = 4.0 Hz, 1H), 7.70 (d, J = 1.6 Hz, 1H), 7.61 (m, 1H), 7.35 (m, 1H), 7.07 (s, 1H), 6.23 (s, 1H), 4.56 (s, 2H), 4.09 (q, J = 7.2 Hz, 2H), 2.65 (s, 3H), 1.25 (t, J = 7.2 Hz, 3H)	m/z 337 [M+1]
1027		2-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-3-methyl-3H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 8.78 (br s, 1H), 8.38 (br s, 1H), 7.87 (m, 2H), 7.68 (s, 1H), 7.55 (br s, 1H), 7.25 (s, 1H), 6.69 (br d, J = 7.5 Hz, 1H), 6.30 (s, 2H), 4.88 (s, 3H)	m/z 309.2 [M+1]
1028		6-ethyl-2-([1-(3-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-1-methyl-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 8.86 (s, 1H), 8.30 (d, J = 4.4 Hz, 1H), 7.71 (s, 1H), 7.62 (m, 1H), 7.36 (m, 1H), 7.07 (s, 1H), 6.18 (s, 1H), 4.58 (s, 2H), 3.64 (s, 3H), 2.94 (q, J = 7.6 Hz, 2H), 1.36 (t, J = 7.6 Hz, 3H)	m/z 337 [M+1]
1029		2-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-3,5-dimethyl-3H-imidazo[4,5-b]pyridine	(CDCl <sub>3</sub> ) 8.86 (s, 1H), 8.30 (d, J = 4.4 Hz, 1H), 7.71 (s, 1H), 7.62 (m, 1H), 7.36 (m, 1H), 7.07 (s, 1H), 6.18 (s, 1H), 4.58 (s, 2H), 3.64 (s, 3H), 2.94 (q, J = 7.6 Hz, 2H), 1.36 (t, J = 7.6 Hz, 3H)	m/z 337 [M+1]



TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1030		1-ethyl-2-([1-(1,3-thiazol-2-yl)-1H-pyrazol-5-yl]methyl)-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 9.00 (s, 1H), 8.43 (d, 1H), 7.63 (d, 1H), 7.43 (d, 1H), 7.15 (d, 1H), 7.06 (d, 1H), 6.25 (d, 1H), 4.99 (s, 2H), 4.25 (q, 2H), 1.36 (t, 3H)	
1031		1-ethyl-2-[1-(1,3-thiazol-2-yl)-1H-pyrazol-5-yl]ethyl-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 9.05 (s, 1H), 8.39 (d, 1H), 7.59 (d, 1H), 7.49 (d, 1H), 7.25 (d, 1H), 7.09 (d, 1H), 6.28 (d, 1H), 5.85 (q, 1H), 4.19 (q, 2H), 1.85 (d, 3H), 1.29 (t, 3H)	
1032		1-ethyl-2-([1-(1,3-thiazol-2-yl)-1H-pyrazol-5-yl]methyl)-6-(trifluoromethyl)-1H-imidazo[4,5-c]pyridine	(CD <sub>3</sub> OD) 8.7 (s, 1H), 8.17 (s, 1H), 7.74 (d, J = 3.6 Hz, 1H), 7.33 (d, J = 3.6 Hz, 1H), 7.21 (d, J = 3.6 Hz, 1H), 7.21 (d, J = 3.6 Hz, 1H), 6.53 (s, 1H), 5.01 (s, 2H), 4.54 (q, J = 7.2 Hz, 2H), 1.52 (t, J = 7.2 Hz, 3H)	
1033		3-ethyl-2-([1-(1,3-thiazol-2-yl)-1H-pyrazol-5-yl]methyl)-3H-imidazo[4,5-c]pyridine	(CD <sub>3</sub> OD) 9.42 (s, 1H), 8.44 (d, J = 6.6 Hz, 1H), 7.9 (d, J = 6.3 Hz, 1H), 7.7 (s, 1H), 7.26 (d, J = 3.6 Hz, 1H), 7.20 (d, J = 3.6 Hz, 1H), 6.63 (s, 1H), 5.09 (s, 2H), 4.68 (q, J = 7.5 Hz, 2H), 1.62 (t, J = 7.5 Hz, 3H)	m/z 311 [M+1]

TABLE 8				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1035		6-chloro-2-[(4-methyl-2-phenyl-1H-imidazol-1-yl)methyl]-1-propyl-1H-benzimidazole	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.62-7.71 (m, 3H), 7.45-7.53 (m, 3H), 7.23-7.27 (m, 2H), 6.74 (q, 1H), 5.42 (s, 2H), 3.59 (t, 2H), 2.21 (d, 3H), 1.37 (m, 2H), 0.70 (t, 3H)	
1036		2-[(4-methyl-2-phenyl-1H-imidazol-1-yl)methyl]-1-propyl-1H-benzimidazole	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.80 (m, 1H), 7.25-7.67 (m, 8H), 6.74 (q, 1H), 5.43 (s, 2H), 3.62 (t, 2H), 2.20 (d, 3H), 1.38 (m, 2H), 0.69 (t, 3H)	
1037		6-chloro-2-[(4-chloro-2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl]-1-propyl-1H-benzimidazole	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 7.70 (d, 1H), 7.37-7.50 (m, 3H), 7.17-7.32 (m, 3H), 6.96 (s, 1H), 5.43 (s, 2H), 3.70 (t, 2H), 1.49 (m, 2H), 0.77 (t, 3H)	
1038		2-[(4-methyl-2-phenyl-1H-imidazol-1-yl)methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.39 (dd, 1H), 8.06 (dd, 1H), 7.63-7.65 (m, 2H), 7.45-7.55 (m, 3H), 7.26 (m, 1H), 6.77 (q, 1H), 5.46 (s, 2H), 3.82 (t, 2H), 2.23 (d, 3H), 1.46 (m, 2H), 0.73 (t, 3H)	
1039		3-ethyl-2-[(2-(3-fluorophenyl)-4-methyl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-b]pyridine	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.39 (dd, 1H), 8.06 (dd, 1H), 7.38-7.50 (m, 3H), 7.12-7.29 (m, 2H), 6.77 (q, 1H), 5.46 (s, 2H), 3.98 (q, 2H), 2.22 (d, 3H), 1.08 (t, 3H)	
1040		2-[(2-(3-fluorophenyl)-4-methyl-1H-imidazol-1-yl)methyl]-3-methyl-3H-imidazo[4,5-b]pyridine		
1041		2-[(2-(2,5-difluorophenyl)-4-methyl-1H-imidazol-1-yl)methyl]-3-methyl-3H-imidazo[4,5-b]pyridine		









TABLE 8				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1042		3-(cyclopropylmethyl)-2-((2-(2,5-difluorophenyl)-4-methyl-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): 8.38 (dd, 1H), 8.04 (dd, 1H), 7.40 (m, 1H), 7.15-7.27 (m, 3H), 6.79 (d, 1H), 5.35 (s, 2H), 3.82 (d, 2H), 2.22 d, 3H), 0.75 (m, 1H), 0.39-0.44 (m, 2H), 0.23-0.28 (m, 2H)	
1043		1-ethyl-2-((2-(3-fluorophenyl)-4-methyl-1H-imidazol-1-yl)methyl)-1H-benzimidazole-5-carbonitrile		m/z 360 [M+1]
1044		1-ethyl-2-((2-(3-fluorophenyl)-5-methyl-1H-imidazol-1-yl)methyl)-1H-benzimidazole-5-carbonitrile		m/z 360 [M+1]
1045		2-((2-methyl-4-phenyl-1,3-thiazol-5-yl)methyl)-1-propyl-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 9.06 (s, 1H), 8.39 (d, J = 5.6 Hz, 1H), 7.61 (m, 2H), 7.47 (m, 2H), 7.41 (m, 1H), 7.20 (d, J = 5.6 Hz, 1H), 4.54 (s, 2H), 3.77 (t, J = 7.6 Hz, 2H), 2.68 (s, 3H),	m/z 349 [M+1]
1046		5-bromo-1-ethyl-2-((3-(3-fluorophenyl)-1H-pyrazol-4-yl)methyl)-1H-benzimidazole		401.0 [M+1]
1047		5-bromo-1-ethyl-2-((5-(3-fluorophenyl)isoxazol-4-yl)methyl)-1H-benzimidazole	(CDCl <sub>3</sub> ) 8.40 (s, 1H), 7.88 (s, 1H), 7.31-7.52 (m, 4H), 7.18-7.22 (m, 2H), 4.14 (s, 2H), 4.04 (q, 2H), 1.18 (t, 3H)	
1048		5-bromo-1-ethyl-2-((5-(3-fluorophenyl)-1-methyl-1H-pyrazol-4-yl)methyl)-1H-benzimidazole	(CDCl <sub>3</sub> ) 7.81 (s, 1H), 7.38-7.45 (m, 2H), 7.32 (dd, 1H), 7.00-7.18 (m, 4H), 4.00 (s, 2H), 3.90 (q, 2H), 3.78 (s, 3H), 1.15 (t, 3H)	
1049		5-bromo-1-ethyl-2-((3-(3-fluorophenyl)-1-methyl-1H-pyrazol-4-yl)methyl)-1H-benzimidazole	(CDCl <sub>3</sub> ) 7.88 (s, 1H), 7.27-7.42 (m, 4H), 7.00-7.20 (m, 3H), 4.21 (s, 2H), 3.95 (t, 2H), 3.83 (s, 3H), 1.15 (t, 3H)	

TABLE 8				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1050		5-bromo-1-ethyl-2-([4-(3-fluorophenyl)-1,3-thiazol-5-yl]methyl)-1H-benzimidazole	(CDCl <sub>3</sub> ) 8.78 (s, 1H), 7.91 (s, 1H), 7.38-7.48 (m, 4H), 7.11-7.19 (m, 2H), 4.59 (s, 2H), 3.90 (q, 2H), 1.12 (t, 3H)	
1051		1-ethyl-2-([4-(6-fluoropyridin-2-yl)-1,3-thiazol-5-yl]methyl)-6-methyl-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 8.92 (s, 1H), 8.75 (s, 1H), 8.16 (d, 1H), 7.90 (dd, 1H), 7.08 (s, 1H), 6.87 (d, 1H), 5.28 (s, 2H), 4.32 (q, 2H), 2.63 (s, 3H), 1.32 (t, 3H)	
1052		2-([3,5-bis(3-fluorophenyl)isoxazol-4-yl]methyl)-1-ethyl-6-(trifluoromethyl)-1H-benzimidazole		m/z 484 [M+1]
1053		2-(2,4'-bi-1,3-thiazol-5'-ylmethyl)-1-ethyl-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 9.05 (s, 1H), 8.76 (s, 1H), 8.41 (d, 1H), 7.88 (d, 1H), 7.41 (d, 1H), 7.25 (d, 1H), 5.35 (s, 2H), 4.26 (q, 2H), 1.25 (t, 3H)	
1054		1-ethyl-2-([2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl)-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 9.10 (s, 1H), 8.47 (d, J = 6.0 Hz, 1H), 7.88 (d, J = 3.2 Hz, 1H), 7.58 (s, 1H), 7.48 (d, J = 3.2 Hz, 1H), 7.31 (d, J = 5.6 Hz, 1H), 6.37 (s, 2H), 4.31 (q, J = 7.2 Hz, 2H), 1.24 (t, J = 7.2 Hz, 3H)	m/z 379 [M+1]
1055		3-ethyl-6-isopropyl-2-([2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 8.75 (s, 1H), 7.87 (d, J = 3.3 Hz, 1H), 7.59 (d, J = 0.9 Hz, 1H), 7.54 (s, 1H), 7.47 (d, J = 3.6 Hz, 1H), 6.32 (s, 2H), 4.38 (q, J = 7.2 Hz, 2H), 3.21 (sept, J = 6.9 Hz, 1H), 1.35 (d, J = 6.9 Hz, 6H), 1.30 (t, J = 7.2 Hz, 3H)	m/z 421 [M+1]
1056		1-ethyl-6-isopropyl-2-([2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl)-1H-imidazo[4,5-c]pyridine	(CDCl <sub>3</sub> ) 9.02 (s, 1H), 7.88 (d, J = 2.7 Hz, 1H), 7.55 (d, J = 0.9 Hz, 1H), 7.48 (d, J = 3.3 Hz, 1H), 7.13 (s, 1H), 6.34 (s, 2H), 4.28 (q, J = 7.2 Hz, 2H), 3.20 (sept, J = 6.9 Hz, 1H), 1.36 (d, J = 6.9 Hz, 6H), 1.22 (t, J = 7.2 Hz, 3H)	m/z 421 [M+1]

Example 515 Preparation of radiolabeled probe compounds of the invention

The compounds of the invention are prepared as radiolabeled probes by carrying out their synthesis using precursors comprising at least one atom that is a radioisotope. The radioisotope is preferably selected from of at least one of  
10 carbon (preferably  $^{14}\text{C}$ ), hydrogen (preferably  $^3\text{H}$ ), sulfur (preferably  $^{35}\text{S}$ ), or iodine (preferably  $^{125}\text{I}$ ). Such radiolabeled probes are conveniently synthesized by a radioisotope supplier specializing in custom synthesis of radiolabeled probe compounds. Such suppliers include Amersham Corporation, Arlington Heights,  
15 IL; Cambridge Isotope Laboratories, Inc. Andover, MA; SRI International, Menlo Park, CA; Wizard Laboratories, West Sacramento, CA; ChemSyn Laboratories, Lexena, KS; American Radiolabeled Chemicals, Inc., St. Louis, MO; and Moravsek Biochemicals Inc., Brea, CA.

20 Tritium labeled probe compounds are also conveniently prepared catalytically via platinum-catalyzed exchange in tritiated acetic acid, acid-catalyzed exchange in tritiated trifluoroacetic acid, or heterogeneous-catalyzed exchange with tritium gas. Such preparations are also conveniently carried out  
25 as a custom radiolabeling by any of the suppliers listed in the preceding paragraph using the compound of the invention as substrate. In addition, certain precursors may be subjected to tritium-halogen exchange with tritium gas, tritium gas reduction of unsaturated bonds, or reduction using sodium borotritide, as  
30 appropriate.

Example 52Receptor autoradiography

Receptor autoradiography (receptor mapping) is carried out in vitro as described by Kuhar in sections 8.1.1 to 8.1.9 of  
5 Current Protocols in Pharmacology (1998) John Wiley & Sons, New York, using radiolabeled compounds of the invention prepared as described in the preceding Example.

Example 5310 Binding Assay

The high affinity and high selectivity of preferred compounds of the invention for the benzodiazepine site of the GABA<sub>A</sub> receptor can be confirmed using the binding assay described by Thomas and Tallman (*J. Bio. Chem.* 1981; 156:9838-9842, and *J.*  
15 *Neurosci.* 1983; 3:433-440).

Rat cortical tissue is dissected and homogenized in 25 volumes (w/v) of Buffer A (0.05 M Tris HCl buffer, pH 7.4 at 4 °C). The tissue homogenate is centrifuged in the cold (4 °C) at 20,000 x g for 20 minutes. The supernatant is decanted, the  
20 pellet rehomogenized in the same volume of buffer, and centrifuged again at 20,000 x g. The supernatant of this centrifugation step is decanted. The resulting pellet may be stored at -20 °C overnight. The pellet is then thawed and resuspended in 25 volumes of Buffer A (original wt/vol),  
25 centrifuged at 20,000 x g and the supernatant decanted. This wash step is repeated once. The pellet is finally resuspended in 50 volumes of Buffer A.

Incubations contain 100 µl of tissue homogenate, 100 µl of radioligand, (0.5 nM <sup>3</sup>H-Ro15-1788 [<sup>3</sup>H-Flumazenil], specific  
30 activity 80 Ci/mmol), and test compound or control (see below), and are brought to a total volume of 500 µl with Buffer A. Incubations are carried for 30 minutes at 4°C and then rapidly filtered through Whatman GFB filters to separate free and bound

ligand. Filters are washed twice with fresh Buffer A and counted in a liquid scintillation counter. Nonspecific binding (control) is determined by displacement of  $^3\text{H}$  Ro15-1788 with 10  $\mu\text{M}$  Diazepam (Research Biochemicals International, Natick, MA): Data are collected in triplicate, averaged, and percent inhibition of total specific binding (Total Specific Binding = Total - Nonspecific) is calculated for each compound.

A competition binding curve may obtained with up to 11 points spanning the compound concentration range from  $10^{-12}\text{M}$  to  $10^{-5}\text{M}$  obtained per curve by the method described above for determining percent inhibition.  $K_i$  values are calculated according the Cheng-Prusoff equation. Each of the compounds disclosed in Tables 6-8 was tested in this fashion and each was found to have a  $K_i$  of  $< 4\mu\text{M}$ . Preferred compounds of the invention exhibit  $K_i$  values of less than 100 nM and more preferred compounds of the invention exhibit  $K_i$  values of less than 10 nM.

#### Example 54

##### 20 Electrophysiology

The following assay can be used to determine if a compound of the invention acts as an agonist, an antagonist, or an inverse agonist at the benzodiazepine site of the  $\text{GABA}_A$  receptor.

Assays are carried out as described in White and Gurley (NeuroReport 6: 1313-1316, 1995) and White, Gurley, Hartnett, Stirling, and Gregory (Receptors and Channels 3: 1-5, 1995) with modifications. Electrophysiological recordings are carried out using the two electrode voltage-clamp technique at a membrane holding potential of -70 mV. *Xenopus Laevis* oocytes are enzymatically isolated and injected with non-polyadenylated cRNA mixed in a ratio of 4:1:4 for  $\alpha$ ,  $\beta$  and  $\gamma$  subunits, respectively. Of the nine combinations of  $\alpha$ ,  $\beta$  and  $\gamma$  subunits described in the White et al. publications, preferred combinations are  $\alpha_1\beta_2\gamma_2$ ,

$\alpha_2\beta_3\gamma_2$ ,  $\alpha_3\beta_3\gamma_2$ , and  $\alpha_5\beta_3\gamma_2$ . Preferably all of the subunit cRNAs in each combination are human clones or all are rat clones. The sequence of each of these cloned subunits is available from GENBANK, e.g., human  $\alpha_1$ , GENBANK accession no. X14766, human  $\alpha_2$ , GENBANK accession no. A28100; human  $\alpha_3$ , GENBANK accession no. A28102; human  $\alpha_5$ , GENBANK accession no. A28104; human  $\beta_2$ , GENBANK accession no. NM 021911; human  $\beta_3$ , GENBANK accession no. M82919 and accession no. Z20136; human  $\gamma_2$ , GENBANK accession no. X15376; rat  $\alpha_1$ , GENBANK accession no. L08490, rat  $\alpha_2$ , GENBANK accession no. L08491; rat  $\alpha_3$ , GENBANK accession no. L08492; rat  $\alpha_5$ , GENBANK accession no. L08494; rat  $\beta_2$ , GENBANK accession no. X15467; rat  $\beta_3$ , GENBANK accession no. X15468; and rat  $\gamma_2$ , GENBANK accession no. L08497. For each subunit combination, sufficient message for each constituent subunit is injected to provide current amplitudes of >10 nA when 1  $\mu$ M GABA is applied.

Compounds are evaluated against a GABA concentration that evokes <10% of the maximal evokable GABA current (e.g. 1 $\mu$ M-9 $\mu$ M). Each oocyte is exposed to increasing concentrations of a compound being evaluated (test compound) in order to evaluate a concentration/effect relationship. Test compound efficacy is calculated as a percent-change in current amplitude:  $100 * ((I_c/I) - 1)$ , where  $I_c$  is the GABA evoked current amplitude observed in the presence of test compound and  $I$  is the GABA evoked current amplitude observed in the absence of the test compound.

Specificity of a test compound for the benzodiazepine site is determined following completion of a concentration/effect curve. After washing the oocyte sufficiently to remove previously applied test compound, the oocyte is exposed to GABA + 1  $\mu$ M RO15-1788, followed by exposure to GABA + 1  $\mu$ M RO15-1788 + test compound. Percent change due to addition of compound is calculated as described above. Any percent change observed in the presence of RO15-1788 is subtracted from the percent changes in current amplitude observed in the absence of 1  $\mu$ M RO15-1788.



These net values are used for the calculation of average efficacy and  $EC_{50}$  values by standard methods. To evaluate average efficacy and  $EC_{50}$  values, the concentration/effect data are averaged across cells and fit to the logistic equation.

5


It is to be understood that the foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the scope of the present invention as set forth in the following

10

claims.

What is claimed is:

1. A compound of the formula:



5 or a pharmaceutically acceptable salt thereof, wherein:

Z<sub>1</sub> is nitrogen or CR<sub>1</sub>;

Z<sub>2</sub> is nitrogen or CR<sub>2</sub>;

Z<sub>3</sub> is nitrogen or CR<sub>3</sub>;

Z<sub>4</sub> is nitrogen or CR<sub>4</sub>;

10 provided that no more than two of Z<sub>1</sub>, Z<sub>2</sub>, Z<sub>3</sub>, and Z<sub>4</sub> are nitrogen;

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from


i) hydrogen, halogen, hydroxy, nitro, cyano, amino, haloalkyl, and haloalkoxy,

ii) alkyl, alkoxy, cycloalkyl, alkenyl, alkynyl,

15 (cycloalkyl)alkyl, -NH(R<sub>10</sub>), -N(R<sub>10</sub>)(R<sub>11</sub>), hydroxyalkyl, aminoalkyl, (R<sub>10</sub>)NHalkyl-, (R<sub>10</sub>)(R<sub>11</sub>)Nalkyl-, alkanoyl, alkoxycarbonyl, (heterocycloalkyl)alkyl, alkylsulfonyl, alkylthio, mono- or dialkylaminocarbonyl, heterocycloalkyl, aryl, and heteroaryl, each of which is optionally  
20 substituted with 1, 2, 3, or 4 of R<sub>20</sub>,

wherein R<sub>10</sub> and R<sub>11</sub> are independently selected at each occurrence from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, (cycloalkyl)alkyl, aryl, arylalkyl, alkanoyl, and mono and dialkylaminoalkyl;  
25 and


iii) a group of the formula:



where G is a bond, alkyl, -O-, -C(=O)-, or -CH<sub>2</sub>C(=O)-, and

$R_A$  is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ ,

iv) a group of the formula



where J is N, CH, or C-alkyl, and

$R_B$  and  $R_C$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, (cycloalkyl)alkyl, heterocycloalkyl, aryl, arylalkyl, alkanoyl, heteroaryl, and mono and dialkylaminoalkyl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, alkoxy, and alkyl;

$R_B$  and  $R_C$  and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain:

a) one or more double bonds,  
 b) one or more of oxo, O, S, SO, SO<sub>2</sub>, or N- $R_D$  wherein  $R_D$  is hydrogen,  $Ar_1$ , alkyl, cycloalkyl, heterocycloalkyl, or  $Ar_1$ alkyl; wherein  $Ar_1$  is aryl or heteroaryl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, alkoxy, and alkyl, and/or

c) one or more substituents  $R_{20}$ ;

v)  $-OC(=O)R_E$ ,  $-C(=O)OR_E$ ,  $-C(=O)NH_2$ ,  $-C(=O)NHR_E$ ,  $-C(=O)NR_ER_F$ ,  $-S(O)_nR_E$ ,  $-S(O)_nNH_2$ ,  $-S(O)_nNHR_E$ ,  $-S(O)_nNR_ER_F$ ,  $-NHC(=O)R_E$ ,  $-C(=NR_E)R_F$ ,  $-HC=N-OH$ ,  $-HC=N(alkoxy)$ ,  $-HC=N(alkyl)$ ,  $-NR_EC(=O)R_F$ ,  $-NHS(O)_mR_E$ , and  $-NR_ES(O)_mR_F$ , where m is 0, 1 or 2, and

R<sub>E</sub> and R<sub>F</sub> are independently selected at each occurrence from alkyl, cycloalkyl, heterocycloalkyl, alkoxy, mono- or dialkylamino, aryl, or heteroaryl each of which is optionally substituted by 1, 2, or 3 of R<sub>30</sub>;

5 R<sub>20</sub> is independently selected at each occurrence from the group consisting of: halogen; hydroxy; nitro; cyano; amino; alkyl; alkoxy optionally substituted with amino or mono- or dialkylamino; cycloalkyl; cycloalkylalkyl; cycloalkylalkoxy; alkenyl; alkynyl; haloalkyl; oxo;  
10 haloalkoxy; mono- and dialkylamino; aminoalkyl; and mono- and dialkylaminoalkyl;

R<sub>30</sub> is independently selected at each occurrence from halogen, hydroxy, nitro, cyano, amino, alkyl, alkoxy optionally substituted with amino or mono- or  
15 dialkylamino, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, heterocycloalkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, oxo, mono- and dialkylamino, aminoalkyl, and mono- and dialkylaminoalkyl;


R<sub>5</sub> represents hydrogen or haloalkyl; or

20 R<sub>5</sub> represents alkyl, cycloalkyl, or (cycloalkyl)alkyl, each of which may contain one or more double or triple bonds, and each of which is optionally substituted with 1, 2, or 3 of R<sub>30</sub>, or

R<sub>5</sub> represents aryl, arylalkyl, heteroaryl, or heteroarylalkyl  
25 each of which is optionally substituted with 1, 2, or 3 substituents selected from the group consisting of haloalkyl, amino, -NH(R<sub>10</sub>), -N(R<sub>10</sub>)(R<sub>11</sub>), carboxamido, (R<sub>10</sub>)NHcarbonyl, (R<sub>10</sub>)(R<sub>11</sub>)Ncarbonyl, halogen, hydroxy, nitro, cyano, amino, alkyl, alkoxy optionally substituted with  
30 amino or mono- or dialkylamino, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, heterocycloalkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, aminoalkyl, and mono- and dialkylaminoalkyl;

Q represents  $-C(R_6)(R_7)$  or oxygen,

with the proviso that Q is not oxygen when  $X_2$  is nitrogen;  
 $R_6$  and  $R_7$  independently represent hydrogen, fluorine, or alkyl;  
 the group:



5

represents a 5 to 7 membered heteroaryl or heterocycloalkyl ring  
 containing up to 4 heteroatoms independently selected from  
 nitrogen, sulfur, and oxygen, said 5 to 7 membered  
 heteroaryl or heterocycloalkyl ring is substituted at each  
 10 carbon atom by R, and substituted at each nitrogen atom  
 available for substitution by R', wherein

R is independently chosen at each occurrence from hydrogen,  
 halogen, amino, alkyl, alkenyl, alkynyl, alkoxy,  
 cycloalkyl, (cycloalkyl)alkyl, haloalkyl, haloalkoxy,  
 15 carboxamido, and 3- to 7-membered carbocyclic or  
 heterocyclic groups which are saturated, unsaturated,  
 or aromatic, which may be further substituted with one  
 or more substituents independently selected from  
 halogen, oxo, hydroxy, alkyl, and alkoxy;

20 R' is independently chosen at each occurrence from alkyl,  
 hydrogen, cycloalkyl, cycloalkyl(alkyl), and 3- to 7-  
 membered carbocyclic or heterocyclic groups which are  
 saturated, unsaturated, or aromatic, which 3- to 7-  
 membered carbocyclic or heterocyclic groups are  
 25 optionally substituted with one or more substituents  
 independently selected from halogen, oxo, hydroxy,  
 alkyl, and alkoxy;

$X_1$  and  $X_2$  independently represent nitrogen, carbon or CH;

Y is nitrogen, oxygen, carbon,  $-CH-$ ,  $-CH_2-$ , or absent; and

30 W represents aryl or heteroaryl, wherein the aryl or heteroaryl  
 group is optionally substituted with up to 4 groups

independently selected from  $R_{30}$ ,  $-\text{CO}_2\text{H}$ ,  $-\text{C}(=\text{O})\text{OR}_E$ ,  $-\text{C}(=\text{O})\text{NHR}_E$ ,  $-\text{C}(=\text{O})\text{NR}_E\text{R}_F$ ,  $-\text{C}(\text{O})\text{R}_E$ , and  $-\text{S}(\text{O})_m\text{R}_E$ ,  $-\text{OR}_E$ , where  $R_{30}$  and  $R_E$  are as defined above and  $m$  is 0, 1, or 2.


5 2. A compound or salt according to Claim 1, wherein

$R_1$ ,  $R_2$ ,  $R_3$ , and  $R_4$  are independently selected from

i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo( $\text{C}_1\text{-C}_6$ ) alkyl, and halo( $\text{C}_1\text{-C}_6$ )alkoxy,

10 ii) ( $\text{C}_1\text{-C}_6$ )alkyl, ( $\text{C}_1\text{-C}_6$ )alkoxy, ( $\text{C}_3\text{-C}_8$ )cycloalkyl, ( $\text{C}_2\text{-C}_6$ )alkenyl, alkynyl, (( $\text{C}_3\text{-C}_8$ )cycloalkyl)( $\text{C}_1\text{-C}_4$ )alkyl,  $-\text{NH}(\text{R}_{10})$ ,  $-\text{N}(\text{R}_{10})(\text{R}_{11})$ , hydroxy( $\text{C}_1\text{-C}_6$ )alkyl, amino( $\text{C}_1\text{-C}_6$ )alkyl, ( $\text{R}_{10})\text{NH}(\text{C}_1\text{-C}_6)$ alkyl, ( $\text{R}_{10})(\text{R}_{11})\text{N}(\text{C}_1\text{-C}_6)$ alkyl, ( $\text{C}_1\text{-C}_6$ )alkanoyl, ( $\text{C}_1\text{-C}_6$ )alkoxycarbonyl, ( $\text{C}_1\text{-C}_6$ )alkylsulfonyl, ( $\text{C}_1\text{-C}_6$ )alkylthio, 15 (heterocycloalkyl) $\text{C}_1\text{-C}_4$ alkyl, aryl, and heteroaryl, each of which is optionally substituted with 1, 2, 3, or 4 of  $\text{R}_{20}$ , wherein  $\text{R}_{10}$  and  $\text{R}_{11}$  are independently selected from the group consisting of ( $\text{C}_1\text{-C}_6$ )alkyl, ( $\text{C}_2\text{-C}_6$ )alkenyl, ( $\text{C}_1\text{-C}_6$ )alkoxy, ( $\text{C}_3\text{-C}_8$ )cycloalkyl, ( $\text{C}_3\text{-C}_8$ )cycloalkylalkyl, 20 aryl, aryl( $\text{C}_1\text{-C}_6$ )alkyl, ( $\text{C}_1\text{-C}_6$ )alkanoyl, and mono and di( $\text{C}_1\text{-C}_6$ )alkylaminoalkyl;


iii) a group of the formula:



where  $G$  is ( $\text{C}_1\text{-C}_6$ )alkyl,  $-\text{O}-$ ,  $-\text{C}(=\text{O})-$ , or  $-\text{CH}_2\text{C}(=\text{O})-$ , and

25  $\text{R}_A$  is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring consisting of from 3 to 8 ring atoms, and each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O; said saturated, partially unsaturated, or 30 aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of  $\text{R}_{20}$ , and

iv) a group of the formula



where J is N, CH, or C-(C<sub>1</sub>-C<sub>6</sub>)alkyl and

R<sub>B</sub> and R<sub>C</sub> are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl) (C<sub>1</sub>-C<sub>4</sub>)alkyl, heterocycloalkyl, aryl, aryl(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkanoyl, heteroaryl, and mono and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, C<sub>1</sub>-C<sub>6</sub>alkoxy, and C<sub>1</sub>-C<sub>6</sub>alkyl; or

R<sub>B</sub> and R<sub>C</sub> and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

a) one or more double bonds;

b) one or more of oxo, O, S, SO, SO<sub>2</sub>, and N-R<sub>D</sub> wherein R<sub>D</sub> is hydrogen, Ar<sub>1</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, heterocycloalkyl, or Ar<sub>1</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl; wherein Ar<sub>1</sub> is aryl or heteroaryl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, C<sub>1</sub>-C<sub>6</sub>alkoxy, and C<sub>1</sub>-C<sub>6</sub>alkyl; and/or

c) one or more substituents R<sub>20</sub>;

v) -OC(=O)R<sub>E</sub>, -C(=O)OR<sub>E</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NHR<sub>E</sub>, -C(=O)NR<sub>E</sub>R<sub>F</sub>, -S(O)<sub>n</sub>R<sub>E</sub>, -S(O)<sub>n</sub>NH<sub>2</sub>, -S(O)<sub>n</sub>NHR<sub>E</sub>, -S(O)<sub>n</sub>NR<sub>E</sub>R<sub>F</sub>, -NHC(=O)R<sub>E</sub>, -C(=NR<sub>E</sub>)R<sub>F</sub>, -HC=N-OH, -HC=N(C<sub>1</sub>-C<sub>6</sub>alkoxy), -HC=N(C<sub>1</sub>-C<sub>6</sub>alkyl), -NR<sub>E</sub>C(=O)R<sub>F</sub>, -NHS(O)<sub>m</sub>R<sub>E</sub>, and -NR<sub>E</sub>S(O)<sub>m</sub>R<sub>F</sub>, where m is 0, 1 or 2, and

R<sub>E</sub> and R<sub>F</sub> are independently selected at each occurrence from (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, heterocycloalkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aryl, and

heteroaryl each of which is optionally substituted by  
1, 2, or 3 of R<sub>30</sub>;

R<sub>20</sub> is independently selected at each occurrence from the  
group consisting of halogen; hydroxy; nitro; cyano;  
5 amino; (C<sub>1</sub>-C<sub>6</sub>)alkyl; (C<sub>1</sub>-C<sub>6</sub>)alkoxy optionally  
substituted with amino or mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino;  
(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl; (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl;  
(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkoxy; (C<sub>2</sub>-C<sub>6</sub>)alkenyl; (C<sub>2</sub>-  
C<sub>6</sub>)alkynyl; halo(C<sub>1</sub>-C<sub>6</sub>)alkyl; halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy; oxo;  
10 mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino; amino(C<sub>1</sub>-C<sub>6</sub>)alkyl; and  
mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>30</sub> is independently selected at each occurrence from  
halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl,  
(C<sub>1</sub>-C<sub>6</sub>)alkoxy optionally substituted with amino or  
15 mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl,  
(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl,  
(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkoxy, heterocycloalkyl,  
(C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, oxo, mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino,  
20 amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and  
mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> represents hydrogen or halo(C<sub>1</sub>-C<sub>6</sub>)alkyl; or

R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, or  
(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl, each of which may contain  
25 one or more double or triple bonds, and each of which is  
optionally substituted with 1, 2, or 3 of R<sub>30</sub> or

R<sub>5</sub> represents aryl, aryl(C<sub>1</sub>-C<sub>4</sub>)alkyl, heteroaryl, or  
heteroaryl(C<sub>1</sub>-C<sub>4</sub>)alkyl each of which is optionally  
substituted with 1, 2, or 3 substituents selected from the  
30 group consisting of:

halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, NH(R<sub>10</sub>), N(R<sub>10</sub>)(R<sub>11</sub>), carboxamido,  
NH(R<sub>10</sub>)carbonyl, N(R<sub>10</sub>)(R<sub>11</sub>)carbonyl, halogen, hydroxy,  
nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy optionally




substituted with amino or mono- or  
 di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl,  
 (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkoxy, heterocyclo(C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 5 (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
 halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and  
 mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

Q represents -C(R<sub>6</sub>)(R<sub>7</sub>) or oxygen,

with the proviso that Q is not oxygen when X<sub>2</sub> is nitrogen;

10 R<sub>6</sub> and R<sub>7</sub> independently represent hydrogen, fluorine, or  
 C<sub>1</sub>-C<sub>6</sub>alkyl;

the group:



represents a 5 to 7 membered heteroaryl or heterocycloalkyl ring


15 containing up to 4 heteroatoms selected from nitrogen,  
 sulfur, and oxygen, said 5 to 7 membered heteroaryl or  
 heterocycloalkyl ring is substituted at each carbon atom by  
 R, and is substituted at each nitrogen atom available for  
 substitution by R', wherein

20 R is independently chosen at each occurrence from hydrogen,  
 halogen, amino, C<sub>1</sub>-C<sub>6</sub>alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-  
 C<sub>6</sub>)alkynyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-  
 C<sub>8</sub>cycloalkyl)(C<sub>1</sub>-C<sub>4</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, haloalkoxy,  
 carboxamido, and 3- to 7-membered carbocyclic or  
 25 heterocyclic groups which are saturated, unsaturated,  
 or aromatic, which may be further substituted with one  
 or more substituents independently selected from  
 halogen, oxo, hydroxy, C<sub>1-4</sub>alkyl, and -O(C<sub>1-4</sub>alkyl);

R' is independently chosen at each occurrence from hydrogen,  
 30 C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl(C<sub>1</sub>-C<sub>4</sub>alkyl),  
 and 3- to 7-membered carbocyclic or heterocyclic groups

which are saturated, unsaturated, or aromatic, which 3-  
to 7-membered carbocyclic or heterocyclic groups are  
optionally substituted with one or more substituents  
independently selected from halogen, oxo, hydroxy,  
5 C<sub>1-4</sub>alkyl, and -O(C<sub>1-4</sub>alkyl); and  
X<sub>1</sub>, X<sub>2</sub>, W, and Y are as defined in Claim 1.

3. A compound or salt according to Claim 2 of the formula:



10 wherein Z<sub>1</sub>, Z<sub>2</sub>, Z<sub>3</sub>, Z<sub>4</sub>, R<sub>5</sub>, Q, X<sub>1</sub>, X<sub>2</sub>, and W are as defined in  
Claim 2;

X<sub>3</sub> and X<sub>4</sub> are independently selected from the group consisting of  
carbon, CR, N, O, S, NH, and N(C<sub>1</sub>-C<sub>6</sub>)alkyl;  
provided that at least one of X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub>, and X<sub>4</sub> is carbon or  
15 CR, wherein  
R is independently chosen at each occurrence from hydrogen,  
halogen, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy,  
(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-  
C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
20 halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, carboxamido, and 3- to 7-membered  
carbocyclic or heterocyclic groups which are saturated,  
unsaturated, or aromatic, which may be further  
substituted with one or more substituents independently  
selected from halogen, oxo, hydroxy, C<sub>1-4</sub>alkyl, and -  
25 O(C<sub>1-4</sub>alkyl).

4. A compound or salt according to Claim 1 wherein  
Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; and Z<sub>4</sub> is CR<sub>4</sub>.

5. A compound or salt according to Claim 2 wherein  
Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; and Z<sub>4</sub> is CR<sub>4</sub>.

6. A compound or salt according to Claim 3 wherein  
5 Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; and Z<sub>4</sub> is CR<sub>4</sub>.

7. A compound or salt according to Claim 6, wherein  
X<sub>2</sub> is carbon; and Q is oxygen.

10 8. A compound or salt according to Claim 6, wherein  
X<sub>2</sub> is N; and Q is C(R<sub>6</sub>) (R<sub>7</sub>).

9. A compound or salt according to Claim 6, wherein  
X<sub>2</sub> is carbon; and Q is C(R<sub>6</sub>) (R<sub>7</sub>).


15

10. A compound or salt according to Claim 6, wherein X<sub>1</sub> is  
carbon; X<sub>2</sub> is N; and Q is C(R<sub>6</sub>) (R<sub>7</sub>).

11. A compound or salt according to Claim 6, wherein X<sub>1</sub> is  
20 nitrogen; X<sub>2</sub> is carbon; and Q is C(R<sub>6</sub>) (R<sub>7</sub>).

12. A compound or salt according to Claim 6, wherein Q is  
C(R<sub>6</sub>) (R<sub>7</sub>).

25 13. A compound or salt according to Claim 6 of the formula



wherein R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, Q, and W are as defined in Claim 6.

14. A compound or salt according to Claim 13 wherein Q is C(R<sub>6</sub>)(R<sub>7</sub>).

15. A compound or salt according to Claim 14, wherein:
- 5 R is independently selected at each occurrence from the group consisting of
- i) hydrogen, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, and
- 10 ii) phenyl and pyridyl each of which is optionally substituted with up to 3 substituents independently chosen from halogen, hydroxy, C<sub>1</sub>-C<sub>4</sub>alkyl, and -O(C<sub>1</sub>-C<sub>4</sub>alkyl);
- R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, heterocycloalkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;
- 15 R<sub>5</sub> represents hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, phenyl, benzyl, thiophenyl, thiazoyl, pyridyl, imidazolyl, pyrazolyl, or pyrimidinyl;
- R<sub>6</sub> and R<sub>7</sub> independently represent hydrogen, fluorine, or C<sub>1</sub>-C<sub>6</sub>alkyl; and
- 20 W represents phenyl, thienyl, thiazoyl, pyridyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, or pyrimidinyl, each of which is optionally substituted with up to 4 independently selected R<sub>30</sub> groups.

- 30 16. A compound or salt according to Claim 14, wherein:
- R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;

R<sub>1</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or  
5 di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;

Q is CH<sub>2</sub>; and

W represents phenyl, furanyl, thienyl, thiazoyl, pyridyl,  
10 imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, pyrimidinyl, benzimidazolyl, quinolinyl, isoquinolinyl each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

15 17. A compound or salt according to Claim 16 wherein R<sub>1</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and W is phenyl, pyridyl, or thiazolyl, each which is optionally substituted by one or more substituents independently chosen  
20 from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

25 18. A compound or salt according to Claim 17, wherein W is 2-thiazolyl, 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

19. A compound or salt according to Claim 17, wherein R, R<sub>1</sub>, and R<sub>4</sub> are hydrogen.


30 20. A compound or salt according to Claim 17, wherein R<sub>5</sub> is ethyl or n-propyl.

21. A compound or salt according to Claim 17 wherein  $R_2$  is chosen from

- i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo( $C_1-C_6$ )alkyl, and halo( $C_1-C_6$ )alkoxy,
- 5 ii)  $C_1-C_6$ alkyl,  $C_1-C_6$ alkoxy,  $C_3-C_8$ cycloalkyl,  $C_2-C_6$ alkenyl,  $C_2-C_6$ alkynyl, ( $C_3-C_8$ cycloalkyl) $C_1-C_4$ alkyl,  $-NH(R_{10})$ ,  $-N(R_{10})(R_{11})$ ,  $(R_{10})NH(C_1-C_6)alkyl$ ,  $(R_{10})(R_{11})N(C_1-C_6)alkyl$ , (heterocycloalkyl)alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

10

22. A compound or salt according to Claim 17 wherein  $R_2$  is a group of the formula



where J is N, CH, or C- $(C_1-C_6)alkyl$  and


- 15  $R_B$  and  $R_C$  are independently selected from the group consisting of hydrogen,  $(C_1-C_6)alkyl$ ,  $(C_2-C_6)alkenyl$ ,  $(C_2-C_6)alkynyl$ ,  $C_3-C_8$ cycloalkyl, and  $(C_3-C_8cycloalkyl)(C_1-C_4)alkyl$ ; or  $R_B$  and  $R_C$  and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

20

- a) one or more double bonds,
- b) one or more of oxo, O, S, SO, SO<sub>2</sub>, and N- $R_D$  wherein  $R_D$  is hydrogen or  $(C_1-C_6)alkyl$ ;
- c) one or more substituents  $R_{20}$ .

25

23. A compound or salt according to Claim 17 wherein  $R_2$  is a group of the formula:



where G is a bond or  $C_1-C_2alkyl$ ; and

$R_A$  is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially  
5 unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

24. A compound or salt according to Claim 23 wherein  $R_A$  is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl,  
10 triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

25. A compound or salt according to Claim 14, wherein:  
 $R$  is independently selected at each occurrence from the group  
15 consisting of hydrogen, halogen, and  $(C_1-C_2)$ alkyl;  
 $R_1$ ,  $R_2$ , and  $R_4$  are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_3-C_8)$ cycloalkyl,  $(C_3-C_8)$ cycloalkyl  $(C_1-C_6)$ alkyl, halo  $(C_1-C_6)$ alkyl, halo  $(C_1-C_6)$ alkoxy, mono or  
20 di  $(C_1-C_6)$ alkylamino, amino  $(C_1-C_6)$ alkyl, and mono- and di  $(C_1-C_6)$ alkylamino  $(C_1-C_6)$ alkyl;

$R_5$  represents  $(C_1-C_6)$ alkyl;

$Q$  is  $CH_2$ ; and

$W$  represents phenyl, furanyl, thienyl, thiazoyl, pyridyl,  
25 imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, pyrimidinyl, benzimidazolyl, quinolinyl, isoquinolinyl each of which is optionally substituted with up to 4  $R_{30}$  groups.

30 26. A compound or salt according to Claim 25 wherein  $R_1$ ,  $R_2$ , and  $R_4$  are independently selected from hydrogen, halogen, trifluoromethyl,  $C_1-C_2$  alkyl, and cyano; and

W is phenyl, pyridyl, or thiazolyl, each of which is optionally substituted by one or more substituents independently chosen from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

5

27. A compound or salt according to Claim 26, wherein W is 2-thiazolyl, 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

10

28. A compound or salt according to Claim 26, wherein R, R<sub>1</sub>, and R<sub>4</sub> are hydrogen.

29. A compound or salt according to Claim 26, wherein R<sub>5</sub> is ethyl or n-propyl.


15

30. A compound or salt according to Claim 26 wherein R<sub>3</sub> is chosen from

- i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy,
- 20 ii) C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, -NH(R<sub>10</sub>), -N(R<sub>10</sub>)(R<sub>11</sub>), (R<sub>10</sub>)NH(C<sub>1</sub>-C<sub>6</sub>)alkyl, (R<sub>10</sub>)(R<sub>11</sub>)N(C<sub>1</sub>-C<sub>6</sub>)alkyl, (heterocycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

25

31. A compound or salt according to Claim 26 wherein R<sub>3</sub> is a group of the formula



where J is N, CH, or C-(C<sub>1</sub>-C<sub>6</sub>)alkyl and



$R_B$  and  $R_C$  are independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl,  $C_3-C_8$ cycloalkyl, and  $(C_3-C_8$ cycloalkyl)  $(C_1-C_4)$ alkyl; or


5  $R_B$  and  $R_C$  and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

a) one or more double bonds,

b) one or more of oxo, O, S, SO,  $SO_2$ , and  $N-R_D$  wherein  $R_D$  is hydrogen or  $(C_1-C_6)$ alkyl;

10 c) one or more substituents  $R_{20}$ .

32. A compound or salt according to Claim 26 wherein  $R_3$  is a group of the formula:



15 where G is a bond or  $C_1-C_2$ alkyl; and


$R_A$  is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially  
20 unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

33. A compound or salt according to Claim 32 wherein  $R_A$  is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl,  
25 triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

34. A compound or salt according to Claim 26 wherein  $R_3$  is  $-HC=N-OH$  or  $-HC=N(C_1-C_6$  alkoxy).

30

35. A compound or salt according to Claim 6 of the formula



wherein R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and W are as defined in Claim 6.

- 5           36. A compound or salt according to Claim 35, wherein:  
 R is independently selected at each occurrence from the group  
 consisting of
- i) hydrogen, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-  
 C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
 10       halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, and
  - ii) phenyl and pyridyl each of which is optionally substituted  
 with up to 3 substituents independently chosen from halogen,  
 hydroxy, C<sub>1-4</sub>alkyl, and -O(C<sub>1-4</sub>alkyl);
- R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen,  
 15       halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-  
 C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
 (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, heterocycloalkyl, halo(C<sub>1</sub>-  
 C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino,  
 amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-  
 20       C<sub>6</sub>)alkyl;
- R<sub>5</sub> represents hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-  
 C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, phenyl, benzyl, thiophenyl,  
 thiazoyl, pyridyl, imidazolyl, pyrazolyl, or pyrimidinyl;
- R<sub>6</sub> and R<sub>7</sub> independently represent hydrogen, fluorine, or C<sub>1</sub>-C<sub>6</sub>  
 25       alkyl; and
- W represents phenyl, thienyl, thiazoyl, pyridyl, imidazolyl,  
 oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, or

pyrimidinyl, each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

37. A compound or salt according to Claim 35, wherein:

5 W represents a 6-membered aryl or heteroaryl groups, wherein the 6-membered aryl or heteroaryl group is optionally substituted with up to 4 groups independently selected from R<sub>30</sub>, -CO<sub>2</sub>H, -C(=O)OR<sub>E</sub>, -C(=O)NHR<sub>E</sub>, -C(=O)NR<sub>E</sub>R<sub>F</sub>, -C(O)R<sub>E</sub>, -S(O)<sub>m</sub>R<sub>E</sub>, and -OR<sub>E</sub>; and m is 0, 1, or 2.

10

38. A compound or salt according to Claim 35, wherein:

W represents a 5-membered heteroaryl group, wherein the 5-membered heteroaryl group is optionally substituted with up to 4 groups independently selected from R<sub>30</sub>, -CO<sub>2</sub>H, -C(=O)OR<sub>E</sub>,  
15 -C(=O)NHR<sub>E</sub>, -C(=O)NR<sub>E</sub>R<sub>F</sub>, -C(O)R<sub>E</sub>, -S(O)<sub>m</sub>R<sub>E</sub>, and -OR<sub>E</sub>, and m is 0, 1, or 2.

39. A compound or salt according to Claim 35, wherein:

R is independently selected at each occurrence from the group  
20 consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;

R<sub>1</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino,  
25 amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>6</sub> and R<sub>7</sub> are hydrogen; and

W represents phenyl, furanyl, thienyl, thiazoyl, pyridyl,  
30 imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, pyrimidinyl, benzimidazolyl, quinolinyl, isoquinolinyl each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

40. A compound or salt according to Claim 39 wherein  
R<sub>1</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen,  
trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and  
5 W is phenyl, pyridyl, or thiazolyl, each which is optionally  
substituted by one or more substituents independently chosen  
from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-  
C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.


10 41. A compound or salt according to Claim 40, wherein W is  
2-thiazolyl, 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-  
pyridinyl.

15 42. A compound or salt according to Claim 40, wherein R, R<sub>1</sub>,  
and R<sub>4</sub> are hydrogen.

43. A compound or salt according to Claim 40, wherein R<sub>5</sub> is  
ethyl or n-propyl.

20 44. A compound or salt according to Claim 40 wherein  
R<sub>2</sub> is chosen from  
i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-  
C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy,  
ii) C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-  
25 C<sub>6</sub>alkynyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl) C<sub>1</sub>-C<sub>4</sub>alkyl, -NH(R<sub>10</sub>), -  
N(R<sub>10</sub>)(R<sub>11</sub>), (R<sub>10</sub>)NH(C<sub>1</sub>-C<sub>6</sub>)alkyl, (R<sub>10</sub>)(R<sub>11</sub>)N(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
(heterocycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, and heterocycloalkyl, each of  
which is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

30 45. A compound or salt according to Claim 40 wherein  
R<sub>2</sub> is a group of the formula




where J is N, CH, or C-(C<sub>1</sub>-C<sub>6</sub>)alkyl and

R<sub>B</sub> and R<sub>C</sub> are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, and (C<sub>3</sub>-C<sub>8</sub>cycloalkyl) (C<sub>1</sub>-C<sub>4</sub>)alkyl; or

R<sub>B</sub> and R<sub>C</sub> and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

- a) one or more double bonds,
- 10      b) one or more of oxo, O, S, SO, SO<sub>2</sub>, and N-R<sub>D</sub> wherein R<sub>D</sub> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl;
- c) one or more substituents R<sub>20</sub>.

46. A compound or salt according to Claim 40 wherein  
15 R<sub>2</sub> is a group of the formula:



where G is a bond or C<sub>1</sub>-C<sub>2</sub>alkyl; and

R<sub>A</sub> is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

25      47. A compound or salt according to Claim 46 wherein R<sub>A</sub> is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

30      48. A compound or salt according to Claim 40 wherein

R<sub>2</sub> is -HC=N-OH or -HC=N(C<sub>1</sub>-C<sub>6</sub>alkoxy).

49. A compound or salt according to Claim 35, wherein:  
R is independently selected at each occurrence from the group  
5 consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;  
R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen,  
hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy,  
(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-  
C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino,  
10 amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-  
C<sub>6</sub>)alkyl;  
R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
R<sub>6</sub> and R<sub>7</sub> are hydrogen; and  
W represents phenyl, furanyl, thienyl, thiazoyl, pyridyl,  
15 imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl,  
isoxazolyl, pyrimidinyl, benzimidazolyl, quinolinyl,  
isoquinolinyl each of which is optionally substituted with  
up to 4 R<sub>30</sub> groups.

20 50. A compound or salt according to Claim 49 wherein  
R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen,  
trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and  
W is phenyl, pyridyl, or thiazolyl, each which is optionally  
substituted by one or more substituents independently chosen  
25 from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-  
C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

51. A compound or salt according to Claim 50, wherein W is  
2-thiazolyl, 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-  
30 pyridinyl.


52. A compound or salt according to Claim 50, wherein R, R<sub>1</sub>,  
and R<sub>4</sub> are hydrogen.

53. A compound or salt according to Claim 50, wherein  $R_5$  is ethyl or n-propyl.

5 54. A compound or salt according to Claim 50 wherein  $R_3$  is chosen from

- i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo( $C_1-C_6$ )alkyl, and halo( $C_1-C_6$ )alkoxy,
- ii)  $C_1-C_6$ alkyl,  $C_1-C_6$ alkoxy,  $C_3-C_8$ cycloalkyl,  $C_2-C_6$ alkenyl,  $C_2-C_6$ alkynyl, ( $C_3-C_8$ cycloalkyl) $C_1-C_4$ alkyl,  $-NH(R_{10})$ ,  $-N(R_{10})(R_{11})$ ,  $(R_{10})NH(C_1-C_6)alkyl$ ,  $(R_{10})(R_{11})N(C_1-C_6)alkyl$ , (heterocycloalkyl)  $C_1-C_4$ alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

15 55. A compound or salt according to Claim 50 wherein  $R_3$  is a group of the formula




where J is N, CH, or C-( $C_1-C_6$ )alkyl and

$R_B$  and  $R_C$  are independently selected from the group consisting of  
 20 hydrogen, ( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )alkenyl, ( $C_2-C_6$ )alkynyl,  $C_3-C_8$ cycloalkyl, and ( $C_3-C_8$ cycloalkyl) ( $C_1-C_4$ )alkyl; or

$R_B$  and  $R_C$  and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

- 25 a) one or more double bonds,
- b) one or more of oxo, O, S, SO, SO<sub>2</sub>, and N- $R_D$  wherein  $R_D$  is hydrogen or ( $C_1-C_6$ )alkyl;
- c) one or more substituents  $R_{20}$ .

30 56. A compound or salt according to Claim 50 wherein  $R_3$  is a group of the formula:



where G is a bond or C<sub>1</sub>-C<sub>2</sub>alkyl; and

R<sub>A</sub> is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

57. A compound or salt according to Claim 56 wherein R<sub>A</sub> is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

58. A compound or salt according to Claim 50 wherein R<sub>3</sub> is -HC=N-OH or -HC=N(C<sub>1</sub>-C<sub>6</sub>alkoxy).

59. A compound or salt according to Claim 3 wherein:  
Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; Z<sub>4</sub> is CR<sub>4</sub>;  
X<sub>1</sub> is carbon; X<sub>2</sub> is nitrogen; X<sub>3</sub> is CR; X<sub>4</sub> is nitrogen; and Q is C(R<sub>6</sub>)(R<sub>7</sub>).


60. A compound or salt according to Claim 3 wherein  
Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; Z<sub>4</sub> is CR<sub>4</sub>;  
X<sub>1</sub> is carbon; X<sub>2</sub> is nitrogen; X<sub>3</sub> is nitrogen; X<sub>4</sub> is CR; and Q is C(R<sub>6</sub>)(R<sub>7</sub>).

61. A compound or salt according to Claim 3 wherein  
Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; Z<sub>4</sub> is CR<sub>4</sub>;  
X<sub>1</sub> is carbon; X<sub>2</sub> is carbon; X<sub>3</sub> is S; and X<sub>4</sub> is CR.



62. A compound or salt according to Claim 61 wherein Q is C(R<sub>6</sub>) (R<sub>7</sub>) .

63. A compound or salt according to Claim 2, wherein  
 5 Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; Z<sub>4</sub> is CR<sub>4</sub>;  
 and the group




64. A compound or salt according to Claim 63 wherein Q is  
 10 C(R<sub>6</sub>) (R<sub>7</sub>) .

65. A compound or salt according to Claim 3 wherein  
 Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; Z<sub>4</sub> is CR<sub>4</sub>;  
 X<sub>1</sub> is nitrogen; X<sub>2</sub> is carbon; X<sub>3</sub> is nitrogen; and X<sub>4</sub> is CR.  
 15

66. A compound or salt according to Claim 3 wherein  
 Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; Z<sub>4</sub> is CR<sub>4</sub>;  
 X<sub>1</sub> is carbon; X<sub>2</sub> is carbon; X<sub>3</sub> is NH or N(C<sub>1</sub>-C<sub>6</sub>alkyl); and X<sub>4</sub> is  
 CR.  
 20

67. A compound or salt according to Claim 3 wherein  
 Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; Z<sub>4</sub> is CR<sub>4</sub>;  
 X<sub>1</sub> is carbon; X<sub>2</sub> is nitrogen; X<sub>3</sub> is nitrogen; X<sub>4</sub> is nitrogen; and  
 Q is C(R<sub>6</sub>) (R<sub>7</sub>) .  
 25

68. A compound or salt according to Claim 2, wherein  
 Z<sub>1</sub> is CR<sub>1</sub>; Z<sub>2</sub> is CR<sub>2</sub>; Z<sub>3</sub> is CR<sub>3</sub>; Z<sub>4</sub> is CR<sub>4</sub>;  
 and the group



69. A compound or salt according to Claim 3, wherein  
 $Z_1$  is  $CR_1$ ;  $Z_2$  is  $CR_2$ ;  $Z_3$  is  $CR_3$ ;  $Z_4$  is  $CR_4$ ;  
 5  $X_1$  is nitrogen;  $X_2$  is carbon;  $X_3$  is  $CR$ ; and  $X_4$  is nitrogen.

70. A compound or salt according to Claim 69 wherein Q is  
 $C(R_6)(R_7)$ .

- 10 71. A compound or salt according to Claim 3, wherein  
 $Z_1$  is  $CR_1$ ;  $Z_2$  is  $CR_2$ ;  $Z_3$  is  $CR_3$ ;  $Z_4$  is  $CR_4$ ;  
 $X_1$  is nitrogen;  $X_2$  is carbon;  $X_3$  is nitrogen; and  $X_4$  is nitrogen.

- 15 72. A compound or salt according to Claim 71 wherein Q is  
 $C(R_6)(R_7)$ .

73. A compound or salt according to Claim 1 wherein one and  
 only one of  $Z_1$ ,  $Z_2$ ,  $Z_3$ , and  $Z_4$  is nitrogen.

- 20 74. A compound or salt according to Claim 2 wherein one and  
 only one of  $Z_1$ ,  $Z_2$ ,  $Z_3$ , and  $Z_4$  is nitrogen.

75. A compound or salt according to Claim 3 wherein  
 one and only one of  $Z_1$ ,  $Z_2$ ,  $Z_3$ , and  $Z_4$  is nitrogen.

25

76. A compound or salt according to Claim 75 wherein  
 either  $Z_2$  or  $Z_3$  is nitrogen; and  
 W represents a 5-membered heteroaryl group, the 5-membered  
 heteroaryl group is optionally substituted with up to 4  
 30 groups independently selected from  $R_{30}$ ,  $-CO_2H$ ,  $-C(=O)OR_E$ , -

$C(=O)NHR_E$ ,  $-C(=O)NR_ER_F$ ,  $-C(O)R_E$ , and  $-S(O)_mR_E$ ,  $-OR_E$ , where  $R_{30}$  and  $R_E$  are as defined above and  $m$  is 0, 1, or 2.

77. A compound or salt according to Claim 76, wherein  
5  $X_2$  is carbon; and  $Q$  is oxygen.


78. A compound or salt according to Claim 76, wherein  
 $X_2$  is N; and  $Q$  is  $C(R_6)(R_7)$ .

10 79. A compound or salt according to Claim 76, wherein  
 $X_2$  is carbon; and  $Q$  is  $C(R_6)(R_7)$ .

80. A compound or salt according to Claim 76, wherein  $X_1$  is  
carbon;  $X_2$  is N; and  $Q$  is  $C(R_6)(R_7)$ .

15 81. A compound or salt according to Claim 76, wherein  $X_1$  is  
nitrogen;  $X_2$  is carbon; and  $Q$  is  $C(R_6)(R_7)$ .

82. A compound or salt according to Claim 76 of the formula



20 wherein  $R$ ,  $R_1$ ,  $R_2$ ,  $R_4$ ,  $R_5$ ,  $R_6$ ,  $R_7$ , and  $W$  are as defined in Claim  
76.

83. A compound or salt according to Claim 82, wherein:  
25  $R$  is independently selected at each occurrence from the group  
consisting of

- i) hydrogen, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, and
- ii) phenyl and pyridyl each of which is optionally substituted with up to 3 substituents independently chosen from halogen, hydroxy, C<sub>1-4</sub>alkyl, and -O(C<sub>1-4</sub>alkyl);

R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, heterocycloalkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> represents hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, phenyl, benzyl, thiophenyl, thiazoyl, pyridyl, imidazolyl, pyrazolyl, or pyrimidinyl;

R<sub>6</sub> and R<sub>7</sub> independently represent hydrogen, fluorine, or C<sub>1</sub>-C<sub>6</sub>alkyl; and

W represents thienyl, thiazolyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, or isoxazolyl each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

84. A compound or salt according to Claim 82, wherein:

R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;

R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>6</sub> and R<sub>7</sub> are hydrogen; and

W represents furanyl, thienyl, thiazoyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, or isoxazolyl, each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

5        85. A compound or salt according to Claim 84 wherein  
R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen,  
trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and  
W is thiazolyl which is optionally substituted by one or more  
substituents independently chosen from halogen, cyano,  
10        hydroxy, oxo, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

86. A compound or salt according to Claim 85, wherein W is  
2-thiazolyl.


15        87. A compound or salt according to Claim 85, wherein R, R<sub>1</sub>,  
and R<sub>4</sub> are hydrogen.

88. A compound or salt according to Claim 85, wherein R<sub>5</sub> is  
ethyl or n-propyl.  
20

89. A compound or salt according to Claim 85 wherein  
R<sub>2</sub> is chosen from  
i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-  
C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy,  
25        ii) C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>1</sub>-  
C<sub>6</sub>alkynyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, -NH(R<sub>10</sub>), -N(R<sub>10</sub>)(R<sub>11</sub>),  
(R<sub>10</sub>)NH(C<sub>1</sub>-C<sub>6</sub>)alkyl, (R<sub>10</sub>)(R<sub>11</sub>)N(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
(heterocycloalkyl) C<sub>1</sub>-C<sub>4</sub>alkyl, and heterocycloalkyl, each of  
which is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

30        90. A compound or salt according to Claim 85 wherein  
R<sub>2</sub> is chosen from hydrogen, halogen, hydroxy, nitro, cyano,  
amino, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy.

91. A compound or salt according to Claim 85 wherein  $R_2$  is a group of the formula



5 where J is N, CH, or C-(C<sub>1</sub>-C<sub>6</sub>)alkyl and


$R_B$  and  $R_C$  are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, and (C<sub>3</sub>-C<sub>8</sub>cycloalkyl) (C<sub>1</sub>-C<sub>4</sub>)alkyl; or

10  $R_B$  and  $R_C$  and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

- a) one or more double bonds,
- b) one or more of oxo, O, S, SO, SO<sub>2</sub>, and N-R<sub>D</sub> wherein R<sub>D</sub> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and/or
- c) one or more substituents R<sub>20</sub>.

15

92. A compound or salt according to Claim 85 wherein  $R_2$  is a group of the formula:



where G is a bond or C<sub>1</sub>-C<sub>2</sub>alkyl; and

20  $R_A$  is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally


25 substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

93. A compound or salt according to Claim 92 wherein  $R_A$  is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is

30 is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

94. A compound or salt according to Claim 85 wherein  $R_2$  is  $-HC=N-OH$  or  $-HC=N(C_1-C_6\text{alkoxy})$ .

5 95. A compound or salt according to Claim 76 of the formula



wherein R,  $R_1$ ,  $R_2$ ,  $R_4$ ,  $R_5$ ,  $R_6$ ,  $R_7$ , and W are as defined in Claim 76.

10 96. A compound or salt according to Claim 75, wherein:  
R is independently selected at each occurrence from the group consisting of

- i) hydrogen, halogen,  $(C_1-C_6)$ alkyl,  $(C_3-C_8)$ cycloalkyl,  $(C_3-C_8)$ cycloalkyl  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy, halo  $(C_1-C_6)$ alkyl, halo  $(C_1-C_6)$ alkoxy, and
- 15 ii) phenyl and pyridyl each of which is optionally substituted with up to 3 substituents independently chosen from halogen, hydroxy,  $C_{1-4}$ alkyl, and  $-O(C_{1-4})$ alkyl;

$R_1$ ,  $R_3$ , and  $R_4$  are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_3-C_8)$ cycloalkyl,  $(C_3-C_8)$ cycloalkyl  $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl, heterocycloalkyl, halo  $(C_1-C_6)$ alkyl, halo  $(C_1-C_6)$ alkoxy, mono or di  $(C_1-C_6)$ alkylamino, amino  $(C_1-C_6)$ alkyl, and mono- and di  $(C_1-C_6)$ alkylamino  $(C_1-C_6)$ alkyl;

20 25

$R_5$  represents hydrogen,  $(C_1-C_6)$ alkyl,  $(C_3-C_8)$ cycloalkyl,  $(C_3-C_8)$ cycloalkyl  $(C_1-C_6)$ alkyl, phenyl, benzyl, thiophenyl, thiazoyl, pyridyl, imidazolyl, pyrazolyl, or pyrimidinyl;

R<sub>6</sub> and R<sub>7</sub> independently represent hydrogen, fluorine, or C<sub>1</sub>-C<sub>6</sub> alkyl; and

W represents thienyl, thiazoyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, or isoxazolyl each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

97. A compound or salt according to Claim 95, wherein:  
R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;  
10 R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;  
15 R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
R<sub>6</sub> and R<sub>7</sub> are hydrogen; and  
W represents furanyl, thienyl, thiazoyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, or isoxazolyl, each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

98. A compound or salt according to Claim 97 wherein  
R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen, trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and  
25 W is thiazolyl which is optionally substituted by one or more substituents independently chosen from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

99. A compound or salt according to Claim 98, wherein W is  
30 2-thiazolyl.

100. A compound or salt according to Claim 98, wherein R, R<sub>1</sub>, and R<sub>4</sub> are hydrogen.



101. A compound or salt according to Claim 98, wherein  $R_5$  is ethyl or n-propyl.

5


102. A compound or salt according to Claim 98 wherein  $R_3$  is chosen from

- i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo( $C_1$ - $C_6$ )alkyl, and halo( $C_1$ - $C_6$ )alkoxy, and
- 10 ii)  $C_1$ - $C_6$ alkyl,  $C_1$ - $C_6$ alkoxy,  $C_3$ - $C_8$ cycloalkyl,  $C_2$ - $C_6$ alkenyl,  $C_2$ - $C_6$ alkynyl, ( $C_3$ - $C_8$ cycloalkyl)  $C_1$ - $C_4$ alkyl,  $-NH(R_{10})$ ,  $-N(R_{10})(R_{11})$ ,  $(R_{10})NH(C_1-C_6)alkyl$ ,  $(R_{10})(R_{11})N(C_1-C_6)alkyl$ , (heterocycloalkyl) $C_1$ - $C_4$ alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

15

103. A compound or salt according to Claim 102 wherein  $R_3$  is chosen from hydrogen, halogen, hydroxy, nitro, cyano, amino, halo( $C_1$ - $C_6$ )alkyl, and halo( $C_1$ - $C_6$ )alkoxy.

20 104. A compound or salt according to Claim 98 wherein  $R_3$  is a group of the formula



where J is N, CH, or C- $(C_1-C_6)alkyl$  and

$R_B$  and  $R_C$  are independently selected from the group consisting of

25 hydrogen,  $(C_1-C_6)alkyl$ ,  $(C_2-C_6)alkenyl$ ,  $(C_2-C_6)alkynyl$ ,  $C_3$ - $C_8$ cycloalkyl, and  $(C_3-C_8cycloalkyl)(C_1-C_4)alkyl$ ; or


$R_B$  and  $R_C$  and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

a) one or more double bonds,

30 b) one or more of oxo, O, S, SO, SO<sub>2</sub>, and N- $R_D$  wherein  $R_D$  is hydrogen or  $(C_1-C_6)alkyl$ ; and/or

c) one or more substituents  $R_{20}$ .

105. A compound or salt according to Claim 98 wherein  $R_3$  is a group of the formula:



where  $G$  is a bond or  $C_1$ - $C_2$ alkyl; and

$R_A$  is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

106. A compound or salt according to Claim 105 wherein  $R_A$  is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

107. A compound or salt according to Claim 98 wherein  $R_2$  is  $-HC=N-OH$  or  $-HC=N(C_1-C_6\text{alkoxy})$ .

108. A compound or salt according to Claim 76 wherein:  $X_1$  is carbon;  $X_2$  is nitrogen;  $X_3$  is CR; and  $X_4$  is nitrogen; and Q is  $C(R_6)(R_7)$ .

25

109. A compound or salt according to Claim 76 wherein  $X_1$  is carbon;  $X_2$  is nitrogen;  $X_3$  is nitrogen;  $X_4$  is CR; and Q is  $C(R_6)(R_7)$ .

110. A compound or salt according to Claim 76 wherein  $X_1$  is carbon;  $X_2$  is carbon;  $X_3$  is S; and  $X_4$  is CR.

30

111. A compound or salt according to Claim 110 wherein Q is C(R<sub>6</sub>)(R<sub>7</sub>).


112. A compound or salt according to Claim 76 wherein  
 5 X<sub>1</sub> is nitrogen; X<sub>2</sub> is carbon; X<sub>3</sub> is nitrogen; and X<sub>4</sub> is CR.

113. A compound or salt according to Claim 76 wherein  
 X<sub>1</sub> is carbon; X<sub>2</sub> is carbon; X<sub>3</sub> is NH or N(C<sub>1</sub>-C<sub>6</sub>alkyl); and X<sub>4</sub> is  
 CR.

10

114. A compound or salt according to Claim 76 wherein  
 X<sub>1</sub> is carbon; X<sub>2</sub> is nitrogen; X<sub>3</sub> is nitrogen; X<sub>4</sub> is nitrogen; and  
 Q is C(R<sub>6</sub>)(R<sub>7</sub>).

15 115. A compound or salt according to Claim 74, wherein  
 either Z<sub>2</sub> or Z<sub>3</sub> is nitrogen;  
 and the group



W represents a 5-membered heteroaryl group, the 5-membered  
 20 heteroaryl group is optionally substituted with up to 4 groups  
 independently selected from R<sub>30</sub>, -CO<sub>2</sub>H, -C(=O)OR<sub>E</sub>, -C(=O)NHR<sub>E</sub>, -  
 C(=O)NR<sub>E</sub>R<sub>F</sub>, -C(O)R<sub>E</sub>, and -S(O)<sub>m</sub>R<sub>E</sub>, -OR<sub>E</sub>, where R<sub>30</sub> and R<sub>E</sub> are as  
 defined above and m is 0, 1, or 2.

25 116. A compound or salt according to Claim 76, wherein  
 X<sub>1</sub> is nitrogen; X<sub>2</sub> is carbon; X<sub>3</sub> is CR; and X<sub>4</sub> is nitrogen.

117. A compound or salt according to Claim 116 wherein Q is  
 C(R<sub>6</sub>)(R<sub>7</sub>).

30

118. A compound or salt according to Claim 76, wherein

X<sub>1</sub> is carbon; X<sub>2</sub> is carbon; X<sub>3</sub> is NH or NCH<sub>3</sub>; and X<sub>4</sub> is CR.

119. A compound or salt according to Claim 76, wherein  
X<sub>1</sub> is nitrogen; X<sub>2</sub> is carbon; X<sub>3</sub> is nitrogen; and X<sub>4</sub> is nitrogen.

5

120. A compound or salt according to Claim 119 wherein Q is  
C(R<sub>6</sub>)(R<sub>7</sub>).

121. A compound or salt according to Claim 75 wherein  
10 either Z<sub>2</sub> or Z<sub>3</sub> is nitrogen; and  
W represents a 6-membered aryl or heteroaryl group, the 6-  
membered aryl or heteroaryl group is optionally substituted  
with up to 4 groups independently selected from R<sub>30</sub>, -CO<sub>2</sub>H, -  
C(=O)OR<sub>E</sub>, -C(=O)NHR<sub>E</sub>, -C(=O)NR<sub>E</sub>R<sub>F</sub>, -C(O)R<sub>E</sub>, and -S(O)<sub>m</sub>R<sub>E</sub>, -  
15 OR<sub>E</sub>, where R<sub>30</sub> and R<sub>E</sub> are as defined above and m is 0, 1, or  
2.

122. A compound or salt according to Claim 121, wherein  
X<sub>2</sub> is carbon; and Q is oxygen.

20

123. A compound or salt according to Claim 121, wherein  
X<sub>2</sub> is N; and Q is C(R<sub>6</sub>)(R<sub>7</sub>).


124. A compound or salt according to Claim 121, wherein  
25 X<sub>2</sub> is carbon; and Q is C(R<sub>6</sub>)(R<sub>7</sub>).

125. A compound or salt according to Claim 121, wherein X<sub>1</sub>  
is carbon; X<sub>2</sub> is N; and Q is C(R<sub>6</sub>)(R<sub>7</sub>).

30 126. A compound or salt according to Claim 121, wherein X<sub>1</sub>  
is nitrogen; X<sub>2</sub> is carbon; and Q is C(R<sub>6</sub>)(R<sub>7</sub>).

127. A compound or salt according to Claim 121, wherein Q is C(R<sub>6</sub>) (R<sub>7</sub>).

128. A compound or salt according to Claim 121 of the  
5 formula



wherein R, R<sub>1</sub>, R<sub>2</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and W are as defined in Claim 121.

129. A compound or salt according to Claim 128, wherein:  
R is independently selected at each occurrence from the group consisting of
- i) hydrogen, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, and
  - ii) phenyl and pyridyl each of which is optionally substituted with up to 3 substituents independently chosen from halogen, hydroxy, C<sub>1-4</sub>alkyl, and -O(C<sub>1-4</sub>alkyl);
- R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, heterocycloalkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;
- R<sub>5</sub> represents hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, phenyl, benzyl, thiophenyl, thiazoyl, pyridyl, imidazolyl, pyrazolyl, or pyrimidinyl;

R<sub>6</sub> and R<sub>7</sub> independently represent hydrogen, fluorine, or C<sub>1</sub>-C<sub>6</sub> alkyl; and

W represents phenyl, pyrimidinyl, pyridyl, pyridizinyl, or pyrazinyl, each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

130. A compound or salt according to Claim 128, wherein:  
R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;  
10 R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;  
15 R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
R<sub>6</sub> and R<sub>7</sub> are hydrogen; and  
W represents phenyl, pyrimidinyl, pyridyl, pyridizinyl, or pyrazinyl each of which is optionally substituted with up to  
20 4 R<sub>30</sub> groups.

131. A compound or salt according to Claim 130 wherein  
R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen, trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and  
25 W is phenyl or pyridyl, each of which is optionally substituted by one or more substituents independently chosen from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

30 132. A compound or salt according to Claim 131, wherein W is 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

133. A compound or salt according to Claim 131, wherein R, R<sub>1</sub>, and R<sub>4</sub> are hydrogen.

134. A compound or salt according to Claim 131, wherein R<sub>5</sub> is ethyl or n-propyl.


135. A compound or salt according to Claim 131 wherein R<sub>2</sub> is chosen from

- i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, and
- ii) C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, -NH(R<sub>10</sub>), -N(R<sub>10</sub>)(R<sub>11</sub>), (R<sub>10</sub>)NH(C<sub>1</sub>-C<sub>6</sub>)alkyl, (R<sub>10</sub>)(R<sub>11</sub>)N(C<sub>1</sub>-C<sub>6</sub>)alkyl, (heterocycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

136. A compound or salt according to Claim 135 wherein R<sub>2</sub> is chosen from hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy.

20

137. A compound or salt according to Claim 131 wherein R<sub>2</sub> is a group of the formula



where J is N, CH, or C-(C<sub>1</sub>-C<sub>6</sub>)alkyl and

R<sub>B</sub> and R<sub>C</sub> are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, and (C<sub>3</sub>-C<sub>8</sub>cycloalkyl)(C<sub>1</sub>-C<sub>4</sub>)alkyl; or


R<sub>B</sub> and R<sub>C</sub> and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

a) one or more double bonds,

30

- b) one or more of oxo, O, S, SO, SO<sub>2</sub>, and N-R<sub>D</sub> wherein R<sub>D</sub> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
 c) one or more substituents R<sub>20</sub>.

5 138. A compound or salt according to Claim 131 wherein R<sub>2</sub> is a group of the formula:



where G is a bond or C<sub>1</sub>-C<sub>2</sub>alkyl; and

RA is a saturated, partially unsaturated, or aromatic carbocycle,  
 10 consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.


15

139. A compound or salt according to Claim 138 wherein R<sub>A</sub> is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

20

140. A compound or salt according to Claim 131 wherein R<sub>2</sub> is -HC=N-OH or -HC=N(C<sub>1</sub>-C<sub>6</sub>alkoxy).

141. A compound or salt according to Claim 121 of the  
 25 formula





wherein R, R<sub>1</sub>, R<sub>2</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and W are as defined in Claim 141.

142. A compound or salt according to Claim 141, wherein:
- 5 R is independently selected at each occurrence from the group consisting of
- i) hydrogen, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, and
- 10 ii) phenyl and pyridyl each of which is optionally substituted with up to 3 substituents independently chosen from halogen, hydroxy, C<sub>1-4</sub>alkyl, and -O(C<sub>1-4</sub>alkyl);
- R<sub>1</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, heterocycloalkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or
- 15 di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;
- 20 R<sub>5</sub> represents hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, phenyl, benzyl, thiophenyl, thiazoyl, pyridyl, imidazolyl, pyrazolyl, or pyrimidinyl;
- R<sub>6</sub> and R<sub>7</sub> independently represent hydrogen, fluorine, or C<sub>1</sub>-C<sub>6</sub>alkyl; and
- 25 W represents phenyl, pyridyl, pyridiziny, pyrimidinyl, or pyrazinyl, each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

143. A compound or salt according to Claim 142, wherein:
- 30 R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;
- R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy,

(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

5 R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>6</sub> and R<sub>7</sub> are hydrogen; and

W represents phenyl, pyridyl, pyridizynyl, pyrimidinyl, or pyrazinyl, each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

10

144. A compound or salt according to Claim 143 wherein R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen, trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and

15 W is phenyl or pyridyl, each of which is optionally substituted by one or more substituents independently chosen from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

20 145. A compound or salt according to Claim 144, wherein W is 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

146. A compound or salt according to Claim 144, wherein R<sub>1</sub> and R<sub>4</sub> are hydrogen.

25 147. A compound or salt according to Claim 144, wherein R<sub>5</sub> is ethyl or n-propyl.


148. A compound or salt according to Claim 144 wherein R<sub>3</sub> is chosen from

- 30 i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy,  
ii) C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, -NH(R<sub>10</sub>),

-N(R<sub>10</sub>)(R<sub>11</sub>), (R<sub>10</sub>)NH(C<sub>1</sub>-C<sub>6</sub>)alkyl, (R<sub>10</sub>)(R<sub>11</sub>)N(C<sub>1</sub>-C<sub>6</sub>)alkyl, (heterocycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

5 149. A compound or salt according to Claim 148 wherein R<sub>3</sub> is chosen from hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy.

10 150. A compound or salt according to Claim 144 wherein R<sub>3</sub> is a group of the formula



where J is N, CH, or C-(C<sub>1</sub>-C<sub>6</sub>)alkyl and R<sub>B</sub> and R<sub>C</sub> are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, and (C<sub>3</sub>-C<sub>8</sub>cycloalkyl)(C<sub>1</sub>-C<sub>4</sub>)alkyl; or


15 R<sub>B</sub> and R<sub>C</sub> and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

- a) one or more double bonds,
- b) one or more of oxo, O, S, SO, SO<sub>2</sub>, and N-R<sub>D</sub> wherein R<sub>D</sub> is

20 hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

- c) one or more substituents R<sub>20</sub>.

151. A compound or salt according to Claim 144 wherein R<sub>3</sub> is a group of the formula:



25 where G is a bond or C<sub>1</sub>-C<sub>2</sub>alkyl; and R<sub>A</sub> is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently

30 chosen from N, S, and O, said saturated, partially

unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

152. A compound or salt according to Claim 151 wherein R<sub>A</sub> is  
5 chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>.

153. A compound or salt according to Claim 144 wherein  
10 R<sub>2</sub> is -HC=N-OH or -HC=N(C<sub>1</sub>-C<sub>6</sub>alkoxy).

154. A compound or salt according to Claim 121 wherein:  
X<sub>1</sub> is carbon; X<sub>2</sub> is nitrogen; X<sub>3</sub> is CR; X<sub>4</sub> is nitrogen; and Q is  
C(R<sub>6</sub>) (R<sub>7</sub>).  
15

155. A compound or salt according to Claim 121 wherein  
X<sub>1</sub> is carbon; X<sub>2</sub> is nitrogen; X<sub>3</sub> is nitrogen; X<sub>4</sub> is CR; and Q is  
C(R<sub>6</sub>) (R<sub>7</sub>).  
20


156. A compound or salt according to Claim 121 wherein  
X<sub>1</sub> is carbon; X<sub>2</sub> is carbon; X<sub>3</sub> is S; X<sub>4</sub> is CR; and Q is C(R<sub>6</sub>) (R<sub>7</sub>).  
25

157. A compound or salt according to Claim 156 wherein  
X<sub>1</sub> is nitrogen; X<sub>2</sub> is carbon; X<sub>3</sub> is nitrogen; and X<sub>4</sub> is CR.  
30

158. A compound or salt according to Claim 121 wherein  
X<sub>1</sub> is carbon; X<sub>2</sub> is carbon; X<sub>3</sub> is NH or N(C<sub>1</sub>-C<sub>6</sub>alkyl); and X<sub>4</sub> is  
CR.

159. A compound or salt according to Claim 121 wherein  
X<sub>1</sub> is carbon; X<sub>2</sub> is nitrogen; X<sub>3</sub> is nitrogen; X<sub>4</sub> is nitrogen; and  
Q is C(R<sub>6</sub>) (R<sub>7</sub>).  
35

160. A compound or salt according to Claim 119, wherein either  $Z_2$  or  $Z_3$  is nitrogen; and the group



5 W represents a 6-membered aryl or heteroaryl group, wherein the 6-membered aryl or heteroaryl group is optionally substituted with up to 4 groups independently selected from  $R_{30}$ ,  $-CO_2H$ ,  $-C(=O)OR_E$ ,  $-C(=O)NHR_E$ ,  $-C(=O)NR_ER_F$ ,  $-C(O)R_E$ , and  $-S(O)_mR_E$ ,  $-OR_E$ , where  $R_{30}$  and  $R_E$  are as defined above and  $m$  is  
10 0, 1, or 2.

161. A compound or salt according to Claim 121, wherein  $X_1$  is nitrogen;  $X_2$  is carbon;  $X_3$  is CR; and  $X_4$  is nitrogen.

15 162. A compound or salt according to Claim 161 wherein Q is  $C(R_6)(R_7)$ .

163. A compound or salt according to Claim 121, wherein  $X_1$  is nitrogen;  $X_2$  is carbon;  $X_3$  is nitrogen; and  $X_4$  is nitrogen.  
20

164. A compound or salt according to Claim 163 wherein Q is  $C(R_6)(R_7)$ .

165. A pharmaceutical composition comprising a compound or  
25 salt according to Claim 1 combined with at least one pharmaceutically acceptable carrier or excipient.

166. A method for altering the signal-transducing activity of a  $GABA_A$  receptor, said method comprising contacting a cell  
30 expressing such a receptor with an amount of a compound or salt according to Claim 1 sufficient to detectably alter the

electrophysiology of the cell, wherein a detectable alteration of the electrophysiology of the cell indicates an alteration of the signal-transducing activity of GABA<sub>A</sub> receptors.

5        167. A method for altering the signal-transducing activity of a GABA<sub>A</sub> receptor, said method comprising contacting a cell expressing such receptors with an amount of a compound or salt according to Claim 1 to detectably alter the chloride conductance in vitro of cell expressing GABA<sub>A</sub> receptors.

10

168. The method of Claim 167 wherein the cell is recombinantly expresses a heterologous GABA<sub>A</sub> receptor and the alteration of the electrophysiology of the cell is detected by intracellular recording or patch clamp recording.

15

169. The method of Claim 167 wherein the cell is a neuronal cell that is contacted in vivo in an animal, the cell is contacted with the compound or salt dissolved in a body fluid, and the alteration in the electrophysiology of the cell is  
20 detected as a change in the animal's behavior.

170. The method of Claim 169 wherein the animal is a human, the neuronal cell is a brain cell, and the body fluid is cerebrospinal fluid.

25

171. A method for altering the signal-transducing activity of a GABA<sub>A</sub> receptor, the method comprising exposing a cell expressing the GABA<sub>A</sub> receptor to an amount of a compound or salt according to Claim 1 sufficient to inhibit R015-1788 binding in  
30 vitro to cells expressing a human GABA<sub>A</sub> receptor.

172. A method for the treatment of anxiety, depression, a sleep disorder, schizophrenia, attention deficit-hyperactivity

disorder, or for the enhancement of memory, comprising administering an effective amount of a compound or salt of Claim 1 to a patient.

5        173. A method for demonstrating the presence of a GABA<sub>A</sub> receptor in a cell or tissue sample, said method comprising:  
         contacting a cell or tissue sample with a labeled compound  
         or salt according to Claim 1;  
         washing the cell or tissue sample to remove unbound labeled  
10        compound or salt; and  
         detecting the presence of labeled compound or salt in the  
         cell or tissue sample.

         174. The method of Claim 173 in which the cell or tissue  
15        sample is a tissue section.

         175. The method of Claim 173 in which the labeled compound  
         or salt contains a radioactive label or a directly or indirectly  
         luminescent label.  
20

         176. The method of Claim 173 in which the cell or tissue  
         sample is a tissue section, labeled compound or salt contains a  
         radioactive label or a directly or indirectly luminescent label,  
         and the labeled compound or salt is detected autoradiographically  
25        to generate an autoradiogram.

         177. A method for demonstrating the presence of a GABA<sub>A</sub> receptor in a tissue section comprising:  
         contacting the tissue section with a radiolabeled or  
30        luminescently labeled compound or salt according to Claim 1 to  
         yield a contacted tissue section;  
         washing the tissue section to remove unbound labeled  
         compound or salt;

detecting the labeled compound or salt in the tissue section; and

comparing the exposure density of the tissue section with the exposure density of a second tissue section that has not been  
5 contacted with a compound or salt according to Claim 1.

178. A package comprising a pharmaceutical composition of claim 165 in a container and further comprising at least one of:  
instructions for using the composition to treat a patient  
10 suffering from an anxiety disorder, or  
instructions for using the composition to treat a patient suffering from depression, or  
instructions for using the composition to treat a patient suffering from a sleeping disorder,  
15 instructions for using the composition to treat a patient suffering from schizophrenia, or  
instructions for using the composition to treat a patient suffering from attention deficit-hyperactivity disorder.

20 179. A package comprising a pharmaceutical composition of claim 165 in a container and further comprising indicia comprising at least one of: instructions for using the composition to treat a patient suffering from Alzheimer's dementia or instructions for using the composition to enhance  
25 memory in a patient.


180. The use of a compound or salt according to Claim 1 for the manufacture of a medicament.

30 181. The use of a compound or salt according to Claim 1 for the treatment of anxiety, depression, a sleep disorder, schizophrenia, or attention deficit-hyperactivity disorder.



182. The use of a compound or salt according to Claim 1 for the enhancement of memory.


183. A process for preparing a compound of Formula A



5


Formula A

comprising reacting a compound of Formula B



Formula B

10 with a compound of Formula C



Formula C

wherein:

Z<sub>1</sub> is nitrogen or CR<sub>1</sub>;

15 Z<sub>2</sub> is nitrogen or CR<sub>2</sub>;

Z<sub>3</sub> is nitrogen or CR<sub>3</sub>;

Z<sub>4</sub> is nitrogen or CR<sub>4</sub>;

provided that no more than two of Z<sub>1</sub>, Z<sub>2</sub>, Z<sub>3</sub>, and Z<sub>4</sub> are nitrogen;


R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from

20 i) hydrogen, halogen, hydroxy, nitro, cyano, amino, haloalkyl, and haloalkoxy,

ii) alkyl, alkoxy, cycloalkyl, alkenyl, alkynyl, (cycloalkyl)alkyl,  $-NH(R_{10})$ ,  $-N(R_{10})(R_{11})$ , hydroxyalkyl, aminoalkyl,  $(R_{10})NHalkyl$ ,  $(R_{10})(R_{11})Nalkyl$ , alkanoyl, alkoxycarbonyl, (heterocycloalkyl)alkyl, alkylsulfonyl, alkylthio, mono- or dialkylaminocarbonyl, heterocycloalkyl, aryl, and heteroaryl, each of which is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ ,

wherein  $R_{10}$  and  $R_{11}$  are independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, (cycloalkyl)alkyl, aryl, arylalkyl, alkanoyl, and mono and dialkylaminoalkyl; and


iii) a group of the formula:



where G is a bond, alkyl,  $-O-$ ,  $-C(=O)-$ , or  $-CH_2C(=O)-$ , and

$R_A$  is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ , and

iv) a group of the formula



where J is N, CH, or C-alkyl, and

$R_B$  and  $R_C$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, (cycloalkyl)alkyl, heterocycloalkyl, aryl, arylalkyl, alkanoyl, heteroaryl, and mono and dialkylaminoalkyl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, alkoxy, and alkyl;

R<sub>B</sub> and R<sub>C</sub> and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain:

- a) one or more double bonds,
- b) one or more of oxo, O, S, SO, SO<sub>2</sub>, or N-R<sub>D</sub> wherein R<sub>D</sub> is hydrogen, Ar<sub>1</sub>, alkyl, cycloalkyl, heterocycloalkyl, or Ar<sub>1</sub>alkyl; wherein Ar<sub>1</sub> is aryl or heteroaryl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, alkoxy, and alkyl, and/or
- c) one or more substituents R<sub>20</sub>;
- v) -OC(=O)R<sub>E</sub>, -C(=O)OR<sub>E</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NHR<sub>E</sub>, -C(=O)NR<sub>E</sub>R<sub>F</sub>, -S(O)<sub>n</sub>R<sub>E</sub>, -S(O)<sub>n</sub>NH<sub>2</sub>, -S(O)<sub>n</sub>NHR<sub>E</sub>, -S(O)<sub>n</sub>NR<sub>E</sub>R<sub>F</sub>, -NHC(=O)R<sub>E</sub>, -C(=NR<sub>E</sub>)R<sub>F</sub>, -HC=N-OH, -HC=N(alkoxy), -HC=N(alkyl), -NR<sub>E</sub>C(=O)R<sub>F</sub>, -NHS(O)<sub>m</sub>R<sub>E</sub>, and -NR<sub>E</sub>S(O)<sub>m</sub>R<sub>F</sub>, where m is 0, 1 or 2, and

R<sub>E</sub> and R<sub>F</sub> are independently selected at each occurrence from alkyl, cycloalkyl, heterocycloalkyl, alkoxy, mono- or dialkylamino, aryl, or heteroaryl each of which is optionally substituted by 1, 2, or 3 of R<sub>30</sub>;

- R<sub>20</sub> is independently selected at each occurrence from the group consisting of: halogen; hydroxy; nitro; cyano; amino; alkyl; alkoxy optionally substituted with amino or mono- or dialkylamino; cycloalkyl; cycloalkylalkyl; cycloalkylalkoxy; alkenyl; alkynyl; haloalkyl; oxo; haloalkoxy; mono- and dialkylamino; aminoalkyl; and mono- and dialkylaminoalkyl;

- R<sub>30</sub> is independently selected at each occurrence from halogen, hydroxy, nitro, cyano, amino, alkyl, alkoxy optionally substituted with amino or mono- or dialkylamino, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, heterocycloalkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, oxo, mono- and dialkylamino, aminoalkyl, and mono- and dialkylaminoalkyl;

R<sub>5</sub> represents hydrogen or haloalkyl; or

R<sub>5</sub> represents alkyl, cycloalkyl, or (cycloalkyl)alkyl, each of which may contain one or more double or triple bonds, and each of which is optionally substituted with 1, 2, or 3 of

5 R<sub>30</sub>, or

R<sub>5</sub> represents aryl, arylalkyl, heteroaryl, or heteroarylalkyl

each of which is optionally substituted with 1, 2, or 3 substituents selected from the group consisting of

haloalkyl, amino, -NH(R<sub>10</sub>), -N(R<sub>10</sub>)(R<sub>11</sub>), carboxamido,  
10 (R<sub>10</sub>)NHcarbonyl, (R<sub>10</sub>)(R<sub>11</sub>)Ncarbonyl, halogen, hydroxy, nitro, cyano, amino, alkyl, alkoxy optionally substituted with amino or mono- or dialkylamino, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, heterocycloalkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, aminoalkyl, and mono- and  
15 dialkylaminoalkyl;

R<sub>6</sub> and R<sub>7</sub> independently represent hydrogen, fluorine, or alkyl;

R is independently chosen at each occurrence from hydrogen, halogen, amino, C<sub>1</sub>-C<sub>6</sub>alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl)(C<sub>1</sub>-C<sub>4</sub>)alkyl,  
20 halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, haloalkoxy, carboxamido, and 3- to 7-membered carbocyclic or heterocyclic groups which are saturated, unsaturated, or aromatic, which may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, C<sub>1-4</sub>alkyl, and  
25 -O(C<sub>1-4</sub>alkyl); and

W represents aryl or heteroaryl, wherein the aryl or heteroaryl group is optionally substituted with up to 4 groups

independently selected from R<sub>30</sub>, -CO<sub>2</sub>H, -C(=O)OR<sub>E</sub>, -C(=O)NHR<sub>E</sub>, -C(=O)NR<sub>E</sub>R<sub>F</sub>, -C(O)R<sub>E</sub>, and -S(O)<sub>m</sub>R<sub>E</sub>, -OR<sub>E</sub>, where R<sub>30</sub>  
30 and R<sub>E</sub> are as defined above and m is 0, 1, or 2.

184. A process according to Claim 183, wherein:

Z<sub>1</sub> is CR<sub>1</sub>, Z<sub>2</sub> is CR<sub>2</sub>, Z<sub>3</sub> is CR<sub>3</sub>, and Z<sub>4</sub> is CR<sub>4</sub>,

R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;

R<sub>1</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>6</sub> and R<sub>7</sub> are hydrogen; and

W represents phenyl, furanyl, thienyl, thiazolyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, pyrimidinyl, benzimidazolyl, quinolinyl, isoquinolinyl each of which is optionally substituted with up to 4 R<sub>30</sub> groups.

185. A process according to Claim 184, wherein W is 2-thiazolyl, 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

186. A process according to Claim 184, wherein R, R<sub>1</sub>, and R<sub>4</sub> are hydrogen.

187. A process according to Claim 184, wherein R<sub>5</sub> is ethyl or n-propyl.

188. A process according to Claim 184 wherein R<sub>2</sub> is chosen from

i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy,

ii) C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl) C<sub>1</sub>-C<sub>4</sub>alkyl, -NH(R<sub>10</sub>), -N(R<sub>10</sub>)(R<sub>11</sub>), (R<sub>10</sub>)NH(C<sub>1</sub>-C<sub>6</sub>)alkyl, (R<sub>10</sub>)(R<sub>11</sub>)N(C<sub>1</sub>-C<sub>6</sub>)alkyl,

(heterocycloalkyl) $C_1$ - $C_4$ alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of  $R_{20}$ .

189. A process according to Claim 183 wherein

5  $Z_1$  is  $CR_1$ ;

one and only one of  $Z_2$  or  $Z_3$  is nitrogen;

$Z_4$  is  $CR_4$ ; and

$R_2$  or  $R_3$  is chosen from

- 10 i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo( $C_1$ - $C_6$ )alkyl, and halo( $C_1$ - $C_6$ )alkoxy,  
ii)  $C_1$ - $C_6$ alkyl,  $C_1$ - $C_6$ alkoxy,  $C_3$ - $C_8$ cycloalkyl,  $C_2$ - $C_6$ alkenyl,  $C_2$ - $C_6$ alkynyl, ( $C_3$ - $C_8$ cycloalkyl)  $C_1$ - $C_4$ alkyl,  $-NH(R_{10})$ ,  $-N(R_{10})(R_{11})$ ,  $(R_{10})NH(C_1-C_6)alkyl$ ,  $(R_{10})(R_{11})N(C_1-C_6)alkyl$ , (heterocycloalkyl) $C_1$ - $C_4$ alkyl, and heterocycloalkyl, each of which is optionally  
15 substituted with 1, 2, 3, or 4 of  $R_{20}$ ;

$R$  is independently selected at each occurrence from the group consisting of hydrogen, halogen, and ( $C_1$ - $C_2$ )alkyl;

10  $R_1$  and  $R_4$  are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, ( $C_1$ - $C_6$ )alkyl, ( $C_1$ - $C_6$ )alkoxy, ( $C_3$ - $C_8$ )cycloalkyl, ( $C_3$ - $C_8$ )cycloalkyl( $C_1$ - $C_6$ )alkyl, halo( $C_1$ - $C_6$ )alkyl, halo( $C_1$ - $C_6$ )alkoxy, mono or di( $C_1$ - $C_6$ )alkylamino, amino( $C_1$ - $C_6$ )alkyl, and mono- and di( $C_1$ - $C_6$ )alkylamino( $C_1$ - $C_6$ )alkyl;

$R_5$  represents ( $C_1$ - $C_6$ )alkyl;

25  $R_6$  and  $R_7$  are hydrogen;

30  $W$  represents a 5-membered heteroaryl group, the 5-membered heteroaryl group is optionally substituted with up to 4 groups independently selected from  $R_{30}$ ,  $-CO_2H$ ,  $-C(=O)OR_E$ ,  $-C(=O)NHR_E$ ,  $-C(=O)NR_ER_F$ ,  $-C(O)R_E$ , and  $-S(O)_mR_E$ ,  $-OR_E$ , where  $R_{30}$  and  $R_E$  are as defined above and  $m$  is 0, 1, or 2.

190. A process according to Claim 189, wherein  $Z_3$  is nitrogen.

191. A process according to Claim 189 wherein  
R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen,  
trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and  
5 W is thiazolyl, thienyl, imidazolyl, oxazolyl, triazolyl,  
tetrazolyl, pyrazolyl, or isoxazolyl, each of which is  
optionally substituted by one or more substituents  
independently chosen from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-  
C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

10

192. A process according to Claim 191, wherein W is 2-  
thiazolyl.

193. A compound or salt according to Claim 191, wherein R,  
15 R<sub>1</sub> and R<sub>4</sub> are hydrogen.

194. A compound or salt according to Claim 191, wherein R<sub>5</sub>  
is ethyl or n-propyl.

20 195. A process according to Claim 189, wherein Z<sub>2</sub> is  
nitrogen.

196. A process according to Claim 195 wherein  
R<sub>1</sub> and R<sub>4</sub> are independently selected from hydrogen, halogen,  
25 trifluoromethyl, C<sub>1</sub>-C<sub>2</sub> alkyl, and cyano; and  
W is thiazolyl, thienyl, imidazolyl, oxazolyl, triazolyl,  
tetrazolyl, pyrazolyl, or isoxazolyl, each of which is  
optionally substituted by one or more substituents  
independently chosen from halogen, cyano, hydroxy, oxo, C<sub>1</sub>-  
30 C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>alkyl, and C<sub>1</sub>-C<sub>2</sub> alkoxy.

197. A process according to Claim 196, wherein W is 2-  
thiazolyl.

198. A compound or salt according to Claim 196, wherein R, R<sub>1</sub> and R<sub>4</sub> are hydrogen.

5        199. A compound or salt according to Claim 196, wherein R<sub>5</sub> is ethyl or n-propyl.

200. A process according to Claim 183 wherein  
Z<sub>1</sub> is CR<sub>1</sub>;  
10    one and only one of Z<sub>2</sub> or Z<sub>3</sub> is nitrogen;  
Z<sub>4</sub> is CR<sub>4</sub>;  
R<sub>2</sub> or R<sub>3</sub> is chosen from  
i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, and halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy,  
15    ii) C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, (C<sub>3</sub>-C<sub>8</sub>cycloalkyl) C<sub>1</sub>-C<sub>4</sub>alkyl, -NH(R<sub>10</sub>), -N(R<sub>10</sub>)(R<sub>11</sub>), (R<sub>10</sub>)NH(C<sub>1</sub>-C<sub>6</sub>)alkyl, (R<sub>10</sub>)(R<sub>11</sub>)N(C<sub>1</sub>-C<sub>6</sub>)alkyl, (heterocycloalkyl)C<sub>1</sub>-C<sub>4</sub>alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of R<sub>20</sub>;  
20    R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C<sub>1</sub>-C<sub>2</sub>)alkyl;  
R<sub>1</sub>, R<sub>3</sub>, and R<sub>4</sub> are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, mono or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino,  
25    amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and mono- and di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl;  
R<sub>5</sub> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
R<sub>6</sub> and R<sub>7</sub> are hydrogen; and  
30    W represents a 6-membered aryl or heteroaryl group, wherein the 6-membered aryl or heteroaryl group is optionally substituted with up to 4 groups independently selected from R<sub>30</sub>, -CO<sub>2</sub>H, -C(=O)OR<sub>E</sub>, -C(=O)NHR<sub>E</sub>, -C(=O)NR<sub>E</sub>R<sub>F</sub>, -C(O)R<sub>E</sub>, and -



$S(O)_mR_E$ ,  $-OR_E$ , where  $R_{30}$  and  $R_E$  are as defined above and  $m$  is 0, 1, or 2.

201. A process according to Claim 200, wherein  $Z_3$  is  
5 nitrogen.

202. A process according to Claim 201 wherein  
 $R_1$  and  $R_4$  are independently selected from hydrogen, halogen,  
trifluoromethyl,  $C_1$ - $C_2$  alkyl, and cyano; and  
10  $W$  is phenyl, pyrimidinyl, pyridyl, pyrazinyl, or pyridizynyl,  
each of which is optionally substituted by one or more  
substituents independently chosen from halogen, cyano,  
hydroxy, oxo,  $C_1$ - $C_2$ haloalkyl,  $C_1$ - $C_2$ alkyl, and  $C_1$ - $C_2$  alkoxy.

15 203. A process according to Claim 202, wherein  $W$  is 2-  
pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

204. A compound or salt according to Claim 202, wherein  $R$ ,  
 $R_1$  and  $R_4$  are hydrogen.  
20

205. A compound or salt according to Claim 202, wherein  $R_5$   
is ethyl or n-propyl.

206. A process according to Claim 200, wherein  $Z_2$  is  
25 nitrogen.

207. A process according to Claim 206 wherein  
 $R_1$  and  $R_4$  are independently selected from hydrogen, halogen,  
trifluoromethyl,  $C_1$ - $C_2$  alkyl, and cyano; and  
30  $W$  is phenyl, pyrimidinyl, pyridyl, pyrazinyl, or pyridizynyl,  
each of which is optionally substituted by one or more  
substituents independently chosen from halogen, cyano,  
hydroxy, oxo,  $C_1$ - $C_2$ haloalkyl,  $C_1$ - $C_2$ alkyl, and  $C_1$ - $C_2$  alkoxy.

208. A process according to Claim 207, wherein W is 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

5        209. A compound or salt according to Claim 207, wherein R<sub>1</sub> and R<sub>4</sub> are hydrogen.

210. A compound or salt according to Claim 207, wherein R<sub>5</sub> is ethyl or n-propyl.

10